M ESOSCOPIC TRANSPORT AS MANY-BODY PHYSICS

Frederick Green

Centre for Quantum Computer Technology School of Physics, The University of New South Wales Kensington NSW 2052, Australia and Department of Theoretical Physics Research School of Physical Sciences and Engineering

The Australian National University Canberra ACT 0200, Australia

M ukunda P.Das

D epartm ent of T heoretical P hysics Research School of P hysical Sciences and Engineering The Australian N ational University C anberra ACT 0200, A ustralia

1. IN TRODUCTION

M esoscopic transport physics has grown astoundingly in the last decade. This kind of physics continues to astonish by the delicacy of its architectures, and of the experim ents done on them. In every way, m esoscopics provides an ideal test-bed for m any-body theory, since the reduced dim ensionalities and sm all scales naturally am plify the e ects of inter-electron correlations. How ever, there is a deeper reason. It rests with the nature of electronic transport and itsm icroscopic origin in electron-hole uctuations.

The sustained development of this eld depends, therefore, on a matching capacity to understand dynamic electron processes at length scales reaching down to the atom ic. A side from its intrinsic scientic value, such an understanding is crucial to the commercial design and integration of novel electronic devices.

A standard bulk-averaging analysis will break down when actual device sizes approach the typicalm ean free paths for scattering, or the natural screening length.

^{*} To be published in Proceedings of the 25th International W orkshop on Condensed M atter Theories (CM T 25), Canberra 2001.

In such system s, much more care has to be taken. In the last two decades a body of work, widely seen as having its origins in Landauer's early insights [1], has enjoyed great success in mesoscopic transport. The best known result within this approach is the Landauer form ula for the quantized conductance of a one-dimensional wire.

To many practitioners of m esoscopics, the Landauer form ula seem s to possess quasi-magical powers in that it claims to explain a staggering diversity of transport measurements (as witness the authoritative list in Reference [2]). It is a form ula that, in its self-confessed simplicity, bases itself strictly on one-body physics and nothing else. A saresult, it contains none of the important electron correlations that come into prominence for low-dimensional mesoscopic structures. More serious is the fact that Landauer's model ignores a series of indispensable physical bounds on the behavior of the electron gas (see Sections 2 and 3 below for details).

A further important point is that many would-be proofs of Landauer's popular formula insist upon the paradoxical notion that two (or even more) chemical potentials, each thermodynamically distinct, should coexist within a single, closed, electrical circuit when it is out of electro-chemical equilibrium. On that account also, Landauer's prescription falls well short of internal consistency. We will deal with this issue on the way to setting out a proper, microscopic view of small-scale transport.

For a good general reference on m esoscopics, see the book by Ferry and G oodnick [1]. On the status of m esoscopic noise theory as popularly received, refer to B lanter and Buttiker [2]. W e also m ention the recent critiques of Landauer-Buttiker-Im ry theory in the review of m esoscopic transport by A gra t et al. [3].

In the evolution of this new branch of condensed-matter physics, concepts of small-scale current behavior cannot be divorced from their roots within microscopic electron correlations. These roots are im ly grounded in the theory of charged Ferm i liquids [4{6]. W ithin that long-established canon, we have two aim s:

to help m otivate m esoscopics as an inherently m any-body discipline, and

to encourage a reciprocal interest in the fresh opportunities for testing manybody ideas, as o ered by m esoscopic physics.

The relationship between m icroscopic m any-body physics and m esoscopic device physics is indeed two-way. Not only does the former hold the foundations for a true understanding of the latter; m esoscopics, in turn, m anifests (offen quite graphically) the action of the conservation laws. It is the m iscroscopic conservation laws that give m any-body theory its universal reach and its great explanatory power [7].

Attention falls on a prim ary conservation law : gauge invariance. On it, one can build a m and trustworthy picture of dynamic uctuations in an already classic mesoscopic system : the one-dimensional metallic wire, or quantum point contact (QPC). This case study highlights the need for any mesoscopic transport model to be, rst and genuinely, a many-electron theory. It also points to the prospect of researching strong correlations through nonequilibrium current noise, a tool that so far has been unfamiliar to many-body physics.

In Section 2 we review the general conditions applying to an open and externally driven system of mobile electrons, metallic ones in particular. Our discussion recalls a series of insightful, if less often discussed, microscopic analyses. These have been freely available in the literature over the last decade [8{13}. A fter that, in Sec. 3 we

analyze the compressibility of an open conductor. This parameter, closely allied to the electron-hole uctuations, is an exact invariant of the system. It is absolutely unaltered by nonequilibrium transport. This result extends beyond the free electron gas, to system s that are subject to strongly inhom ogeneous C oulom b screening. N onuniform electronic conductors are encountered everywhere in mesoscopic physics, and internal screening leads to the substantial suppression of their compressibility. This e ect too is fully inherited by a system when it is driven out of equilibrium. Last, we foreshadow the elect of exchange-correlation renorm alization on the compressibility.

Charge conservation governs the compressibility of a charged Ferm i liquid. It also governs its response right through the high-eld nonlinear regime. Section 4 retraces our kinetic noise theory with reference to the criticalQPC noise experiment of Reznikov et al. [14], whose features raise some outstanding issues of principle. These are resolved in a natural way within our kinetic many-body perspective, and compressibility is the key to their resolution. We conclude in Sec. 5.

2. TRANSPORT IN OPEN ENVIRONMENTS

A fundamental problem in mesoscopic transport is the explicit consideration of what it means, in microscopic terms, to address \open environments" that no longer seem to admit direct treatment within a closed Ham iltonian formalism. Over the last decade, a number of important comprehensive papers has appeared. Taken as a whole, they contribute substantially towards the realization of a logical and calculable microscopic approach to transport at mesoscopic scales. This is so not just in linear response, but also in truly nonequilibrium situations. The authors responsible for this body of work are:

Sols [8];

Fenton [9,10];

M agnus and Schoenm aker [11,12]; and

Kamenev and Kohn [13].

We rst exam ine the fundam ental principles that they articulate, dissect, and in large measure resolve. It hardly needs saying that their critiques of mesoscopic theory stand upon a time-tested and rigorous analytic tradition, from Maxwell and Boltzmann through Ferm i and Landau.

Neither is it a surprise to nd that such rst-principles descriptions of the mesoscopic electron gas begin { and end { with charge conservation. However, its practical outworking can be subtle. For instance, unlike an ideally closed system of carriers not subject to exchange with the outside, the global charge neutrality of a real { open { conductor is not automatic, so that a nontrivial demonstration is required. Nonetheless, the physics of charge and number conservation applies as strictly as ever.

a.Gauge Invariance

Sols [8] rst established that an open m etallic conductor is globally gauge invariant if, and only if, the m icroscopic equation of m otion for the electrons explicitly includes the contribution from each source and sink of current in the problem. These are located at the device's interfaces with the outlying reservoirs. The reservoirs them selves remain in (local) equilibrium.

Sols' form alism introduces a critical separation of physical roles within the m esoscopic description. Thus, the dynam ical action of charge in jection and rem oval (which brings about the system response) is entirely separate from the equilibrating action of the reservoir leads. The m acroscopic leads act, in the rst instance, not as regulators of the current but to pin and stabilize the nonequilibrium state of the intervening device. The current is not conditioned by the lead equilibria.

The clear implication of Sols' exam ination is that the current sources/sinks operate dynam ically in their own right. They have nothing to do with the locally unchanging state of the reservoirs. A coording to Ref. [8], injection and extraction must act, and be represented, manifestly.

Transport descriptions that take no notice of the entry and exit of ux are necessarily faulty. Ref. [8] shows that the sources and sinks are irreducible elements in a gauge-invariant model. Thus it is theoretically not possible to cut them out of open-system transport merely at convenience, as if these essential ux exchanges could be equated to a remote (and forgettable) background e ect [15].

From Sols, a major point follows:

A microscopically conserving model of transport and uctuations must manifestly account for all in ux and e ux of current at the open boundaries.

Let us now focus on the quiescent reservoirs. They lie outside the conducting sam ple, in which the interesting transport phenom ena all occur. At the same time, they are in intim ate electrical contact with it. As we have remarked, Ref. [8] shows that the reservoirs' microscopic function is distinct from the supply and recovery of the (nonequilibrium) current uxes.

Therefore the reservoir leads have two dom inant properties, which are not dynam ic but, rather, therm odynam ic. The m acroscopic leads are

- (i) always neutral, and
- (ii) always in local equilibrium .

In other words, they pin the carriers asymptotically (via the local Ferm i levels in the respective leads, as well as the therm odynam ic bath tem perature) to their perm anent and locally unchanging equilibrium state. The corollary is that

The nonequilibrium carrier states must connect seam lessly to each invariant equilibrium state that locally and uniquely characterizes each of the leads.

b. Screening

Shortly after Sols' form alproof, Fenton [9,10] m ade a detailed quantum -kinetic analysis of m esoscopic transport. He highlighted the dom inance of charge neutrality of the reservoirs. Through Thom as Ferm i screening, the asymptotic carriers con ne the electrom otive force (EMF) so that it is spatially coextensive with the driven conductor, across whose boundaries it appears.

Fenton drew a deep conclusion from his rst-principles quantum analysis. The quantized conductance observed in one-dimensional (1D) wires is explainable if and

only if its underlying model includes the screening response of the reservoirs at the sample interfaces. This is, of course, a consequence of Gauss' theorem applied to the electron gas (the perfect-screening sum rule [4], also recalled by Sols [8]). It entails the statem ent that

The interface regions of an open mesoscopic sample { the actual sites of the carriers' physical transition to the asymptotic charge-neutral state { must be included as part of the theoretical description of the transport.

As in every electron-gas problem, screening is an integral part of the procedure here. Unless it operates, Landauer's quantized mesoscopic conductance can nd no reconnection with microscopic quantum kinetics [9]. Fenton's analysis requires that { rather than the common practice of demoting metallic-electron screening to some ad hoc perturbative afterthought [15] { a mesoscopic transport description make such collective screening its centerpiece.

c. Ham iltonian Description

So far we have restated the prime roles of gauge invariance, global neutrality, and equilibrium in charged open systems. This is su cient to set up a kinetic theory in the mesoscopic regime, not only in the weak-eld limit but wellaway from equilibrium [16,17].

Since energy dissipation is a real and central e ect in such cases, it is easy to conclude that a canonical H am iltonian approach is not feasible. To the extent that a (dissipative) dynam ical equation cannot be generated from a standard herm itian H am iltonian form, one m ight think that the kinetic description of real carrier dynam ics, with its energy losses, is destined to remain form ally incom plete.

This incom pleteness is not inevitable. A central fact that should be taken into account is the multiply connected topology of a closed circuit. An elegant m icroscopic form alism for nonequilibrium, energy-dissipating and closed m esoscopic circuits has been proposed by M agnus and Schoenm aker [11,12]. They point out that the toroidal topology of the circuit (this includes the battery) adm its valid solutions to M axwell's equations which, though locally conservative, are globally nonconservative along com – plete paths around the hole of the torus.

M agnus and Schoenm aker construct the gauge-invariant H am iltonian for a closed and multiply connected system of driven carriers. The classic m inim al-coupling expression for the electronic H am iltonian is faithfully m aintained. W hat is novel is that the physical solution, resulting from the geom etrical reconnection, w ill dissipate energy from the battery along any full circuit path.

In this way, real processes can be described without altering the locally conservative H am iltonian structure of the interacting electrons. All of the powerful quantum kinetics of K adano -B aym [18], or of K eldysh [19], becomes immediately applicable in principle.

Under very general conditions, the authors form ally prove that the external power P supplied by the source is dissipated according to the standard form ula P = IV, in term s of circuit current I and EM F V. The result is intuitively clear, but challenging to prove. The derivation is fully quantal, and valid at any device scale.

The work of M agnus and Schoenm aker dovetails with that of Sols and Fenton.

The treatments are all equivalent in the end; the latter two stress the governing role of the boundary conditions within a charge-conserving description of a nite mesoscopic device that is electrically open to the outer leads. On the other hand, the form er approach restates the critical fact that any circuit path containing the device must be closed, and that the topology of the closure makes possible (i) long-range neutrality, (ii) dissipation, and hence (iii) dynamical stability within a uni ed H am iltonian description.

d. Uniqueness of the Chem ical Potential

F inally, the recent m any-body derivation of Landauer's conductance form ula by K am enev and K ohn [13] is notew orthy.

- (a) It sets the Landauer result within a solidly orthodox quantum kinetic form ulation, free of the unnecessary phenom enology that has weakened previous such proofs;
- (b) it provides a detailed and self-contained treatm ent of collective screening;
- (c) by applying K ubo response theory in its appropriate form, Ref. [13] naturally preserves the logical structure of them icroscopic uctuation-dissipation theorem;
- (d) it dem onstrates the unphysical character of a purely heuristic scheme now widely accepted in m esoscopics: namely, the practice of \driving" all currents by a kind of pseudodi usive process, generated by a virtual m ism atch in the chemical potential across various leads.

At the two-body level, a consequence of Refs. [8-12] is to rule out that last artice in all its variations, for they all lead inevitably to the violation of gauge invariance [20].


It is K am enev and K ohn's work [13], however, that drives home the ctional nature of pseudodi usive phenom enology, by straightforward ab initio evaluation of K ubo's conductance form ula for a 1D w ire. Just one parameter is needed: the equilibrium chem ical potential. Thus

the microscopic properties of a mesoscopic system are uniquely determined by one chemical potential, and one only: that of the reference equilibrium state.

e. M icroscopics and Phenom enology

Each of the approaches enum erated above counters a view point commonly advanced by other models [2,15,21,22]. As already mentioned, the electrical currents in such phenomenologies are generated and (allegedly) sustained by passive connection of the mesoscopic device to two orm ore large reservoirs at unequal potentials. Instead of being allowed their correct thermodynamic values, measured from the bottom of their occupied local conduction bands, the chemical potentials in the reservoirs are taken from a global zero of energy which itself has no unique physical signi cance. This immediately breaks the canonical linkage that ties together the physical carrier density and the physical compressibility of the collective system [20]. We return to the compressibility below.

In the dom inant conception of mesoscopics, carriers are never driven dynam ically through the conductor, from one lead to another, by the real eld. Rather, they fall passively, and by sheer happenstance, over the associated Ferm i-level drop. All



F igure 1. M esoscopic transport in a nutshell. A closed, i.e. toroidal, loop includes the conductor under test (clear central area). The loop is the essential supplier of the current I, either by a generator as shown or, equivalently, by the EMF from a battery. The source (S) and drain (D) reservoirs ensure (a) inelastic dissipation of the electrical pow er dum ped to the circuit, (b) screening of all elds including the driving eld E (I) across the nonequilibrium region, and (c) continuity of the carrier distribution. These dynam ically competing processes are located at the interfaces (shaded areas). The boundary conditions illustrated here are necessary and su cient for a m icroscopically conserving description. (This F igure is taken from our R ef. [23]).

that can be known of the dynam ics are the asymptotic equilibrium states, and the asymptotic probability for a single carrier to drop over the potential cli.

This picture is of no help in explaining the energy dissipation that has to accompany open-system transport. Pseudodi usive phenom enologies have no access to anything but generic equilibrium properties. From a detailed kinetic perspective, those models cannot contribute to understanding the critical nonequilibrium processes unfolding within the conductor, and at its interfaces with the reservoirs. These processes dissipate energy. D issipation is out of scope.

Figure 1 sum marizes the physical structure of a mesoscopic system open to its macroscopic electrical surroundings, within which all measurements take place. The closed loop supplies the ux, as specified in Ref. [11] while, in accordance with Ref. [8], the ux entry and exit points maintain global conservation as perceived by a test charge within the wire. The large reservoirs maintain therm odynamic stability by pinning the local Ferm i level at the interfaces, by providing energy dissipation (Ref. [11]), and by con ning the electrom otive forces via screening (Ref. [9]).

By simple logic, the phenom enological scenarios cannot allow the reservoirs them selves to reconnect electrically \at in nity" and thus to close the circuit. If they did so, the resulting coexistence of several di erent chem ical potentials (required for pseudodi usive \transport") would pose an imm ediate consistency problem [11,13]. There is no ux generator, and thus no replenishment of the ever-growing charge imbalance between the reservoirs. Therefore an adiabatically sustainable, currentcarrying steady state cannot be described within such an approach.

By contrast, in a standard m any-body description all the dynam ic and dissipative e ects that determ ine both transport and noise conform to the m icroscopic conditions articulated in Refs. [8{13}, and sum m arized in Fig. 1. W ithout those norm ative boundary conditions (which include the batteries and/or current generators), a well-controlled m odel cannot be set up. That is the m essage of the works we have recalled.

3. A CANONICAL SUM RULE:COMPRESSIBILITY

As is well known, the physics of the electron gas is strongly conditioned by a set of canonical theorems, the sum rules [4]. Several of the sum rules are directly linked to number conservation, and thus to gauge invariance. They, if nothing else, should be identically satis ed by any model of transport and uctuations.

The electron-gas sum rules share a common template. On the left-hand side, a physical quantity is given as a function of the therm odynamic and single-particle parameters (temperature; chemical potential; density etc.). This is equated, on the right-hand side, to a two-particle correlation function [24].

Conceptually, the sum rules are closely related to the uctuation-dissipation theorem [25,26]; once again we see a one-body parameter, the electronic conductance, on one side of the relation. On the other side, it is uniquely determined by an irreducible two-body response function: the current-current correlator. Equivalences such as these express the microscopic conservation of energy, momentum, and particle number at each elementary electron-hole vertex in the system 's quantum many-body expansion [5].

W ith this background, let us exam ine the content and m esoscopic consequences of the sim plest of the sum rules, that for compressibility. On its own, that rule is a suprisingly powerful constraint on the behavior of m esoscopic conductors. For the full and form al details, see our Refs. [16,17,20].

a. The Open Electron G as

Consider a driven, m etallic conductor of volum e bounded by equilibrated and charge-neutral reservoirs. Here we invoke the boundary conditions discussed above; see Sec. 2.a. That the well-de ned active region, including the interfaces, must be xed and nite follows from Fenton's m icroscopic analysis; see Sec. 2.b.

W ithin the device, let N be the number of mobile carriers, and let N hN 2 hN i^2 ibe the mean-square uctuation of N, where the averaging will be made precise shortly. Since global neutrality holds,

N is unconditionally invariant for all transport of charge through the system. The total number N can be accounted for in two ways. We can integrate the timedependent nonequilibrium distribution f (t) for the con guration-space label = (r;k) (spin and many-valley indices are implied). A lternatively, we can add up all the electrons in for the equilibrium case, where the one-body distribution is

$$f^{eq} = \frac{1}{1 + \exp[("_{k} + U(r))] = k_{B}T]};$$

The (quasiparticle) kinetic energy is " $_k$; the H artree m ean-eld potential is U (r), and is generally a functional of the density distribution. The global chem ical potential is xed by the outer reservoirs.

Very sim ply,

X
$$f(t) = N = f^{eq}$$
: (1)

The origin and meaning of this equation are clear. We add up all contributions to N over the space of wavevectors k and over all r within the real-space volume . The operation denotes this con gurational integration. Since the region of physical interest is xed and unconditionally neutral, N is always exactly balanced by the unchanging positive background. Equation (1) follows.

We now sketch the corresponding uctuation relation. It is a consequence of Gauss' law, expressed through Eq. (1). The di erence function

$$q(t) = f(t) f^{eq}$$

m easures the local out-of-balance component of the ensemble. Eq. (1) is equivalent to

$$g (t) = 0$$
: (2)

If we take variations of the di erence function, for example with respect to the chem icalpotential proper to the reservoir leads (this is done by changing the leads' asym ptotic electron density and that of its neutralizing positive background), then

$$\frac{\text{@g (t)}}{\text{@}} \qquad \qquad X \qquad X \qquad \frac{\text{g (t)}}{\text{f}_{0}^{\text{eq}}} \frac{\text{@f}_{0}^{\text{eq}}}{\text{@}} = 0:$$
(3)

We have used the variational chain rule to partly unpack the rich inner structure of this uctuation; the functional derivative $g(t) = f_0^{eq}$ is a specialized G reen function whose form is calculable from the kinetic equation for f(t) [17]. It is the centerpiece of our approach.

b. Free-Electron Compressibility

Equation (3) is the cardinal element in deriving the nonequilibrium compressibility sum rule. First, the equilibrium mean-square uctuation in local carrier number is given by

$$f^{eq} = k_B T \frac{\theta f^{eq}}{\theta} = f^{eq} (1 - f^{eq}); \qquad (4a)$$

which lets us compute the total mean-square uctuation in N:

N
$$k_B T \frac{\partial N}{\partial t} = X f^{eq}$$
: (4b)

Second, the exact nonequilibrium m ean-square uctuation is [17]

$$f(t) = f^{eq} + \sum_{0}^{X} \frac{g(t)}{f^{eq}_{0}} f^{eq}_{0}$$
: (5)

From Eq. (3) we get, at last,

X
$$f(t) = N = f^{eq}$$
: (6)

The compressibility for an undisturbed free-electron gas is standard [4]:

$$\frac{0}{N^{2}} \frac{0}{0} = \frac{1}{N k_{B} T} \frac{N}{N}$$
(7)

Note that, so far, we have taken our variations (and the associated uctuation structures) with all internal elds held xed; that is, we freeze the potential U(r). Shortly, we will relax this constraint. At this stage our treatment is precisely analogous to the Lindhard model of the electron gas.

Our result is immediate. Eqs. (1) and (6) are exact within the nonequilibrium kinetic description. Therefore

The invariant compressibility of an open mesoscopic conductor, driven at any current, is given by Eq. (7).

A simple feature of Eq. (7) is seen in its M axwellian limit. In that case, f $^{eq} = f^{eq} / e^{=k_B T}$. The ratio N=N is unity. That renders the right-hand expression equal to the inverse of the pressure P = N $k_B T$ =. Eq. (7) then states the classical, ideal-gas result

$$_{cl} = P^{-1}$$
:

W hen the system is degenerate, then $f^{eq} < f^{eq}$ and

$$free = cl \frac{N}{N} < cl$$
(8)

(we denote the free \Lindhard" compressibility by $_{\rm free}$ in anticipation of the next subsection). This shows the sti ening action of degeneracy on the system, in loose analogy with van der W aals' concept of the excluded hard-core volume. The e ect clearly persists, with no change at all, out of equilibrium. This is despite the fact that the underlying m icroscopic form of the dynam ic f (t) is farm ore com plex than its generic equilibrium limit, f $^{\rm eq}$. The power of the compressibility sum rule com es precisely from its total, and surprising, insensitivity to external driving forces.

We have just exhibited a general result for nonequilibrium electronic conductors. It is the direct consequence of conservation, acting in the context of the boundary conditions of Sec. 2. These boundary conditions are the only ones fully consistent with m icroscopic principles that hold, universally, in a many-body situation. The m esoscopic electron gas is one such.

c. Screened C om pressibility

It is time to focus on the reaction of the internal elds to the carrier uctuations. Intuitively, any such uctuation costs additional electrostatic energy. The system will m inim ize the excess internal energy by self-consistent screening of the initial charge uctuation. This is C oulom b-induced suppression, of which the C oulom b blockade is one of the m ore dram atic instances.

Let us start with the self-consistent equilibrium response:

$$k_{\rm B} T - \frac{f^{\rm eq}}{m} = - U(r) k_{\rm B} T - \frac{\theta f^{\rm eq}}{\theta} = 1 - \frac{dU}{dn} - \frac{n(r)}{r} f^{\rm eq}$$
: (9a)

We have made the usual Ansatz for U (r); one assumes that it depends directly on the local density so that U U [n (r)]. This holds { even mesoscopically { for typical length scales in excess of the inverse Ferm i wavenum ber k_F^{-1} . W ith slightly di erent physics, it applies exactly to quantum -well-con ned systems such as the two-dimensional electron gas formed within heterostructure materials (the basis of very many mesoscopic devices) [27].

The local-density Ansatz provides a constitutive relation to close Eq. (9a). We need another. Denoting by $e^{f} e^{q}$ the left-hand (m ean-square) uctuation in the equation, we have the integral identity

$$\frac{n(r)}{m} = \frac{1}{k_{\rm B}T} h^{\rm ef} \, {}^{\rm eq} {}^{\rm i}{}_{\rm k}$$

where we locally take the trace of e^{f} over the space of wavevectors k. This decouples from the spatial dependence on r, as is appropriate in a local-density treatment.¹ A flerm inor rearrangement, Eq. (9a) goes to

 1 O ur basic screening result does not rely on the local-density assumption. In the wider case, Eq. (9a) reads

$$e_{f} e_{q} = f e_{q} 1 dr_{n} \frac{U(r) n(r^{0})}{n(r^{0})} :$$

Since the mean-eld potential couples only to the density, one can still integrate over k, leaving the real-space (albeit nonlocal) integral relation

$$Z dr^{0} (r r^{0}) + \frac{@n(r)}{@} \frac{U(r)}{n(r^{0})} \frac{n(r^{0})}{-} = \frac{@n(r)}{@};$$

in which the extended response function $U(r) = n(r^0)$ carries the screening. After solving the equation we arrive at $e^{f} e^{q}$, essentially as for Eq. (9b) below:

$$e_{f}e_{q} = \frac{n(r)}{(ln(r))=0} f^{eq}$$

$${}^{e_{f} e_{q}} = \frac{1}{1 + \frac{h f e_{q} i_{k}}{h f} \frac{dU (r)}{dn}} f e_{q} < f e_{q}; \qquad (9b)$$

The leading factor is the inverse static dielectric function. (In its general form it can be seen in the nalequation of the preceding footnote.) The total mean-square number uctuation, in the presence of self-consistent screening, is

$$\Psi = drh^{eq} i_{k}:$$

The screened analog to Eq. (7) can now be stated:

$$\frac{1}{N^2} - \frac{N}{N} = \frac{1}{N k_B T} \frac{q_N}{N}$$
: (10)

Om itting the relevant proof, already fully detailed in Ref. [17], we state the major outcome for nonequilibrium mesoscopic conductors:

In the presence of screening, the invariant compressibility of an open, nonuniform m esoscopic conductor { driven at any current { is given by Eq. (10).


Equation (10) stands in relation to the random -phase approximation (RPA) [4] as Eq. (7) stands in relation to Lindhard. There are di erences, however.

- (a) The conductor must be inhom ogeneous. O there ise U (r) = 0 everywhere, and $= \frac{1}{\text{free}}$. This is merely the rejection of a known fact [4]: in a perfectly uniform system, long-range RPA screening is not manifested.
- (b) The conductor must be degenerate. Classically, the M axwell form of f^{eq} entails equipartition of energy. The C oulom b term in the distribution decouples exactly from the kinetic one, leading to the relation $\P N = N \cdot A \text{ gain}$, $= \frac{1}{\text{free}} = \frac{1}{\text{cl}}$.

A n accessible and striking example of screening-induced suppression occurs in the quasi-two-dimensional electron gas (2DEG) conned within a heterojunction quantum well. In Fig. 2 we display the compressibility of a 2DEG at room temperature, as found in any production-grade A IG aA s/InG aA s/G aA s device. At sheet densities higher than 10^{11} electrons per cm², we see the rapid onset of the C oulom b suppression in where, by comparison, degeneracy acts rather more mildly.

The electrons, Im ly held within the (engineered) quantum well, exhibit their strong mutual repulsion via the density dependence of their Hartree potential U [n (r)]. The consequent reduction of carrier uctuations by self-consistent feedback is an important element of practical hetero junction-device design [27].

Exactly the same C oulom b reduction of f m ust also reveal itself in the nonequilibrium current noise, since the same electron-hole pair processes that underpin the compressibility do so equally for noise [17,20]. We explore this in the next Section. Before doing so, let us add a novel twist to the physics of nonequilibrium compressibility.



F igure 2. Compressibility of a quantum -con ned two-dimensional (2D) electron gas, normalized to that of the 2D classical gas and plotted as a function of sheet electron density. D ashed line: the compressibility of free conduction electrons decreases with increasing density, owing to their degeneracy (see Eq. (8) in the text). Full line: the self-consistent compressibility of interacting 2D electrons (Eq. (10) in text). With increasing density, the mean-eld C oulom b energy of a con ned carrier population will increase. That makes them collectively more resistant to compression over and above free, and provides a prime example of a mesoscopic many-body e ect. Since electron uctuations scale with , mesoscopic noise must directly manifest these (and other) many-body e ects.

d. Exchange and Correlation

So far, we have treated the electron gas regardless of exchange-correlation processes. A lready in this Section, we have seen the many-body nature of uctuation physics come to the fore in the compressibility response of a nonequilibrium conductor. The introduction of exchange-correlation corrections is not a trivial matter; it has many levels of renement. Here, to give the avor of what could be achieved for transport, we will discuss these elects in terms of an additional potential U_{xc} [n (r)].

In density-functional theory, the exchange-correlation potential augments the Hartree term. The total compressibility can be written by inspection from Eq. (9b):

$$tot = \frac{1}{N^2 k_B T} X = \frac{f^{eq}}{1 + \frac{h f^{eq} i_k}{k_B T} \frac{dU}{dn}(r) + \frac{dU_{xc}}{dn}(r)}$$
(11)

At high densities the Hartree, or RPA, term U dom inates. In the low-density limit the counterbalancing exchange-correlation term $U_{\rm xc}$ is expected to dom inate. An interesting density regime should then exist, in which the nonuniform ity of, say, a strongly con ned quantum channel competes against the shorter-ranged part of the electron-electron interaction (with exchange). As far as we know, none of these phe-

nom ena has been discussed system atically in the speci c context of nonequilibrium transport.

We rem ark that the above approach, via an expression for U_{xc} which m ay be quite simple (and certainly static), has much in common with the local-eld correction long familiar in electron-gas theory [28]. A sEq. (11) shows, our procedure amounts to embellishing the RPA with a term that undoes the RPA's over-correction of the internal energy of the system; quasiparticle renorm alization never enters explicitly. Indeed, the sem inalquantum formulation of Boltzm ann transport by K adano and Baym [18] (which informs m any nonequilibrium kinetic theories, including ours) neglects the interaction corrections to single-particle propagation, while keeping particle-particle collision terms that are of comparable order in the coupling strength.

The result of this partial inclusion of exchange-correlation is a fully conserving and calculable description, but one that is conceptually incom plete (m ore obviously so at low carrier densities). In a mean-eld treatment, the Boltzm ann{K adano -Baym collision term swill yield something close to the exchange-correlation potential U_{xc} . However, the corresponding self-energy e ects are not there.²

It is outside our present brief to look into the com plexities of nonequilibrium – transport e ects induced by quasiparticle renorm alization, and how to include them consistently (let alone their yet-to-be-discussed m esoscopic im plications). The inclusion of renorm alization within a kinetic equation of Boltzm ann type, in a m icroscopically gauge-invariant way, is a challenging and still un nished theoretical project. It is the subject of intensive thought by specialists. W e can do no better than to cite a sm all subset of this work. Several excellent sum m aries, with appropriate references, appear in a volum e on m any-body kinetics collated by Bonitz [29]. Out of m any detailed studies, two that address di erent facets of these problems are Refs. [30] and [31].

4. NONEQUILIBRIUM FLUCTUATIONS:NOISE

In the previous Section we reviewed the essentially many-body uctuation origin of a central result: the compressibility sum rule for the electron gas. We have discussed how all the quantum correlations that condition this constraint are faithfully retained when a system of (uctuating) conduction electrons is

² An interim x for this incom pleteness is to set the elective mass of the carriers in the model to that of the renormalized quasiparticles at the Ferm i surface. How – ever, that parameter itself depends on the Landau quasiparticle parameter F_1^s which is, by denition, the p-wave coecient of the two-body exchange-correlation interaction. Again we discern the quite intricate self-consistency between quasiparticle renormalization and collision properties. A beautiful relation, the forward-scattering sum rule [see e.g. W.F.Brinkman, P.M.Platzman, and T.M.Rice, Phys. Rev. 174, 495 (1968)], ties together all of the Landau partial-wave amplitudes, including the determinant of the elective mass F_1^s , and the s-wave term F_0^s which governs the exchange-correlation component of the fermion antisymmetry that quasiparticles inherit from the underlying \free"-electron modes.

- (1) open to large external reservoirs that uniquely x its equilibrium state, and
- (2) driven out of equilibrium by external current sources (or, equally well, by external EMFs) in a way that always preserves global gauge invariance for the open conductor.

The self-consistent structure of the compressibility rule is intimately tied to charge conservation. This internal consistency must therefore be preserved even in an approximate transport description. A model that violates the compressibility sum rule is an incorrect model.

E orts to set up a sum -rule preserving transport theory would be purely academ ic if the nonequilibrium uctuations of the electron gas could not be probed e ectively in the laboratory. W hile conductance is readily measured, the information that it carries on electron-hole uctuations is very coarse-grained. To study the uctuation structure, one must access the electrical noise generated within the device. If our arguments are correct, the noise leads directly to signi cant many-body e ects that are invisible in the conductance.

One of the rst well-controlled measurements of nonequilibrium noise in a 1D mesoscopic channel was the quantum -point-contact experiment of Reznikov et al. [14]. Am ong several phenomenological treatments of this system, the best known is the Landauer-Buttiker model [2]. It predicts a noise-power spectrum S that is dominated by shot noise.


The predicted shot noise has a well developed sequence of peaks and troughs as the channel's carrier density is swept from the \pinch-o " (zero-density) region, upwards through the region where the Ferm i level successively accesses the quantized conduction subbands of the QPC. At the same time, the conductance G increases in a series of more-or-less delineated stepw ise mesas. Ideally, the steps in G come in integer units of the Landauer value $e^2 = h$ (empirically, they hardly ever do so).

It is norm al experim ental practice to measure G and S, as functions of carrier density, for xed values of the source-drain driving voltage. A lthough the t to experim ent is by no means perfect, notably for noise, the standard noise models [2] give a sem iquantitative account of the observations. A major innovation introduced by R eznikov et al. was to obtain an additional series of noise data for xed values of source-drain current. Here, a glaring discrepancy immediately stands out, between the data and the standard noise prediction.

In Fig. 3 we show a single trace of S as published by Reznikov et al. [14], straddling the rst subband. It is evident that the prediction of a strictly monotonic S by the Landauer-Buttiker noise theory totally m isses the strong peak structure in the data. The authors of the experiment note this. However, they make no further comment on what is a wholly unexpected { and unexplained { outcome.

There is another, purely kinetic-theoretical explanation for the Reznikov et al. peak anomalies. Here we sketch it in the formulation of Thakur and the present writers [32]. Our model, which is strictly conserving, obeys the compressibility sum rule by construction (for further details see Refs. [20] and [23]).

From a standard kinetic Boltzm ann equation for the carrier distribution f(t), we generate an associated equation of motion for the dynam ic mean-square uctuation f(t) [16,17]. The full 1D solution, obtained analytically in the collision-time



F igure 3. Low -frequency excess noise spectrum S of a quantum point contact as described in Ref. [14], measured at xed source-drain current I = 300nA and plotted as a function of carrier density in the rst subband. The density is regulated by the gate potential V_g. Full line: S (with threefold enlargement). At the subband threshold, where the Ferm i level rst accesses the subband, S peaks strongly. D ot-dashed line: the outcom e of Landauer-Buttiker noise theory [2] using the measured conductance. This strictly monotonic prediction totally misses the peak. W hen the channel is depleted, the carriers becom e classical; there the plateau in S is qualitatively sim ilar to shot noise, but falls short of the classical asymptotic value 2eI.

approximation, allows us to write the exact nonequilibrium noise.

Norm alized to the classical shot-noise expression 2eI at source-drain current I, the free-electron excess noise (at low frequency) is


$$\frac{S}{2eI} = -\frac{eI}{_{cl}} \frac{eI}{Gm \ L^2} - \frac{2}{_{in}} + 2 \frac{\frac{el \ in}{_{in}}}{_{el} + _{in}} - \frac{2}{_{el} \frac{2}{_{in}}}{(\frac{el \ in}{_{in}})^2} :$$
(12)

The set of parameters is as follows. The classical compressibility $_{cl} = 1 = nk_BT$ is in terms of the subband carrier density n, keeping only the lowest subband of the QPC as in [14]. The device conductance is G, while the electric electron mass is m and L is the operational length of the 1D channel. The inelastic and elastic collision times are, respectively, in and el. Since the sample is ballistic { at least in the neighborhood of equilibrium { the mean free path associated with el is essentially the ballistic device's length, L. Phonon emission, how ever, will cause the inelastic mean free path to become progressively shorter than L as the current intensi es.

Equation (12) has two principal features. First, the compressibility enters directly into the nonequilibrium excess spectrum. At high density,

$$\frac{k_{\rm B}T}{2"_{\rm F}}$$

where $"_{\rm F}$ is the Fermi energy. Even far from equilibrium, the excess noise necessarily



F igure 4. K inetic-theoretical simulation of the excess noise spectrum of a quantum point contact, corresponding to the conditions of Fig. 3. Density is controlled by the chem ical potential . Full line: S (with threefold enlargement). At the subband threshold our strictly conserving solution for S shows a strong peak that is quantitatively close to the observed structure. D ot-dashed line: the outcome of Landauer-Buttiker noise theory [2], using our kinetically computed conductance. This purely m onotonic prediction stillm isses the peak. C lose to depletion of the channel, our sim - ulation does not exhibit a plateau as seen in Fig. 3. W e ascribe this to the absence of the physically quite distinct shot-noise contribution in our present calculation, which addresses only the excess therm al noise contribution to S.

scales with the ambient temperature. The greater the degeneracy, the smaller S becomes owing to the factor above. At low density, the electrons are classical and

In the classical lim it the factor is independent of tem perature.

Second, the structure of the last right-hand term in Eq. (12) m akes S very sensitive to the ratio of the two collision times. If, as expected, in decreases substantially with the applied current I, the magnitude of the excess noise will be correspondingly reduced. This expectation provides the basis for our simulation of the result quoted in Fig. 3. It is displayed in Fig. 4.

Figure 4 shows our direct computation of S from Eq. (12), for environmental variables chosen to m atch the data of Fig. 3. We parametrize the inelastic time as a function of I and n to account for the enhanced rate of phonon release. This is then fed into Eq. (12). The elastic collision time, being set by impurity scattering only, remains una ected throughout.

We see a strong peak structure in the noise. It arises from the competition among , I, and $_{\rm in}$ (I;n). The corresponding Landauer-Buttiker curve holds no trace of any peak behavior.

It is in portant to keep in m ind that the Landauer-Buttiker approach badly violates the compressibility sum rule, and indeed charge conservation [20]. Our standard kinetic m ethod respects both. V iolation of gauge invariance is built right into the Landauer-Buttiker noise form ula. That is known even to its architects [2].

5. SUM M ARY

In this review we have stressed that the understanding of nonequilibrium m esoscopic transport is indivisible from its m icroscopic origins in the electron gas. The degenerate electron gas is, of course, a prime example of a multi-particle system.

We rst discussed the boundary-condition physics of an open mesoscopic conductor. We did this with reference to a series of searching works by a number of authors. They are works that deserve a wider audience { and a far more seriously engaged one { than the present arbiters of mesoscopic fashion seem able, or perhaps willing, to muster. Certainly the papers we have recalled provide, with considerable power, a prelude for the microscopic analysis of open systems which we have applied in our own nonequilibrium investigations.

W ithin the kinetic paradigm, we discussed how the well-known compressibility sum rule for the electron gas is rigidly satis ed, not just at equilibrium but far from it. This result governs the structure of nonequilibrium uctuations in the driven system. Through the action of the compressibility, the noise spectrum of a mesoscopic wire becomes the distinctive signature of Fermi-liquid correlations in low-dimensional, nonequilibrium metallic-electron transport.

From the compressibility, much more might be learned about low-dimensional current correlations. We illustrated our fundamental many-body approach with the graphic example of high-eld noise in a one-dimensional metallic conductor. Its spectrum is remarkable for its sensitivity to many-body elects, and especially for the tight way in which these are orchestrated by microscopic conservation.

It is obvious that any description of mesoscopic noise, especially one that claim s to be well form ulated, must always conserve charge and particle number. There is little here to soften the implications for other approaches to mesoscopics [2,15,21,22]. The absolutely indispensable action of conservation is plain. So is the direct, and by now documented, evidence of its violation [20]. Moreover the speci c predictions of microscopic kinetics, notably for the nonequilibrium noise of a quantum point contact, are a signi cant element in clarifying the discussion through new experiments.

An expanded program to measure nonequilibrium noise in quantum point contacts would be an excellent place to bok for a breakdown of nonconserving models. It would also verify the microscopic alternative, whose conserving properties are necessary but not su cient for validity. The detailed exploration of electron-electron correlations, through renorm alization of the nonequilibrium -noise spectrum, poses an intriguing and (to our know ledge) untouched possibility. M uch has to be done to realize that potential.

ACKNOW LEDGMENT

In loving appreciation of M arjorie Ann O shorne, 1955 $\{2002: sister-in-law and m anuscript facilitator par excellence. FG.$

REFERENCES

- [1] D.K.Ferry and S.M.Goodnick, Transport in Nanostructures (Cambridge University Press, Cambridge, 1997).
- [2] Ya. M. Blanter and M. Buttiker, Phys. Rep. 336, 1 (2000).
- [3] N.Agra t, A.Levy Yeyati, and J.M. van Ruitenbeek, Preprint cond-m at/0208239.
- [4] D.Pines and P.Nozieres, The Theory of Quantum Liquids (Benjamin, New York, 1966).
- [5] P.Nozieres, Theory of Interacting Ferm i Systems (Benjamin, New York, 1963).
- [6] A A. Abrikosov, Fundam entals of the Theory of M etals (North-Holland, Am sterdam, 1988).
- [7] Recall, for example, the W and P itaevski-Takahashi identities which the the renormalization of single-particle dynamics to the two-particle correlation vertex. The identities are the outworking of microscopic conservation [5]; they also entail the sum rules [4,5] that must be satisfied by the electron-hole uctuations, and thus inevitably the current noise, of an electron gas. See also Ref. [20] below.
- [8] F. Sols, Phys. Rev. Lett. 67, 2874 (1991).
- [9] E.W. Fenton, Phys. Rev. B 46, 3754 (1992).
- [10] E.W. Fenton, Superlattices and M icrostruct. 16, 87 (1994).
- [11] W .M agnus and W .Schoenm aker, J.M ath. Phys. 39, 6715 (1998).
- [12] W .M agnus and W .Schoenm aker, Phys. Rev. B 61, 10883 (2000).
- [13] A.Kamenev and W.Kohn, Phys. Rev. B 63, 155304 (2001).
- [14] M.Reznikov, M.Heiblum, H.Shtrikman, and D.Mahalu, Phys. Rev. Lett. 75 3340 (1995).
- [15] Y. Im ry and R. Landauer, Rev. M od. Phys. 71, S306 (1999).
- [16] F.Green and M.P.Das, J.Phys.: Condens. Matter 12, 5233 (2000).
- [17] F.Green and M.P.Das, J.Phys.: Condens. Matter 12, 5251 (2000).
- [18] L.P.Kadano and G.Baym, Quantum Statistical Mechanics (Benjamin, Reading, 1962).
- [19] L.V.Keldysh, Zh. Exp. Teor. Fiz. 47, 1515 (1964) (Sov Phys. JETP 20, 1018 (1965)).
- [20] F.G reen and M.P.Das, in Noise and Fluctuations Control in Electronic Devices, edited by A.A.Balandin (American Scienti c Publishers, New York, 2002), Ch. 3.
- [21] Y. Im ry, Introduction to M esoscopic Physics (O x ford U niversity P ress, O x ford, 1997).
- [22] S. Datta, Electronic Transport in Mesoscopic Systems (Cambridge University

Press, Cambridge, 1997).

- [23] F.Green and M.P.Das, Fluctuation and Noise Letters 1, C21 (2001).
- [24] From this generic structure, we see that no description of transport { m esoscopic or otherwise { can avoid the need to address truly multi-particle correlations. N ot all self-styled m esoscopic theories m eet this requirement [2,15,21,22].
- [25] N.G. van K am pen, Stochastic P rocesses in P hysics and C hem istry, revised edition (N orth-H olland, Am sterdam, 2001).
- [26] R.Kubo, M. Toda, and N.Hashitsum e, Statistical Physics II: Nonequilibrium Statistical Mechanics, second edition (Springer, Berlin, 1991).
- [27] C.W eisbuch and B.V inter, Quantum Sem iconductor Structures: Fundam entals and Applications, (A cadem ic Press, San Diego, 1991).
- [28] G.D.Mahan, Many-Particle Physics (Plenum, New York, 1990).
- [29] M.Bonitz (editor), Progress in Nonequilibrium Green's Functions (World Scientic, Singapore, 2000).
- [30] K.Morawetz, P.Lipavsky, and V.Spicka, Prog. Part. Nucl. Phys. 42, 147 (1999).
- [31] Yu. B. Ivanov, J. Knoll, and D. N. Voskresensky, Nucl. Phys. A 672, 313 (2001).
- [32] J.S.Thakur, M.P.Das, and F.Green, in process.