

Momentum distribution and correlation function of quasicondensates in elongated traps

F. Gerbier,^{*} J. H. Thywissen,[†] S. Richard, M. Hugbart, P. Bouyer, and A. Aspect
Laboratoire Charles Fabry de l'Institut d'Optique[‡], 91403 Orsay, France
 (Dated: March 22, 2022)

We calculate the spatial correlation function and momentum distribution of a phase-fluctuating, elongated three-dimensional condensate, in a trap and in free expansion. We take the inhomogeneous density profile into account via a local density approximation. We find an almost Lorentzian momentum distribution, in stark contrast with a Heisenberg-limited Thomas-Fermi condensate.

PACS numbers: 03.75.Fi, 03.75.-b, 05.30.Jp

Low-dimensional, degenerate Bose gases are expected to have significantly different coherence properties than their three-dimensional (3D) counterparts. In one-dimensional (1D) uniform systems, no true condensate can exist at any temperature T because of a large population of low-lying states that destroys phase coherence (see [1] and references therein). For a trapped gas, the situation is different: the finite size of the sample naturally introduces a low-momentum cutoff, and at sufficiently low temperature $T \ll T_\phi$, a phase coherent sample can exist [1]. Above T_ϕ , the degenerate cloud is a so-called *quasicondensate*: the density has the same smooth profile as a true condensate, but the phase fluctuates in space and time. As shown in [2], this analysis holds also for 3D condensates in elongated traps even if, strictly speaking, radial motion is not frozen. Such 3D, phase-fluctuating condensates have been recently observed experimentally in equilibrium [3] and nonequilibrium [4] samples.

Phase fluctuations of the condensate are caused mainly by long-wavelength (or low-energy) collective excitations [1, 2, 5]. In elongated traps, the lowest energy modes are 1D excitations along the long axis of the trap [6]. Furthermore, in the long-wavelength limit, density fluctuations are small and can be neglected for the calculation of the correlation function [1, 7]. Then, the single-particle density matrix is, assuming cylindrical symmetry,

$$\langle \hat{\Psi}^\dagger(\rho, z) \hat{\Psi}(\rho, z') \rangle \approx \chi(\rho, Z, s) e^{-\frac{1}{2} \Delta \phi^2(Z, s)}. \quad (1)$$

We have introduced $\Delta \phi^2(Z, s) = \langle [\phi(z) - \phi(z')]^2 \rangle$, the variance of the phase difference between two points z, z' on the axis of the trap, with mean coordinate $Z = (z + z')/2$ and relative distance $s = z - z'$, and the overlap function $\chi = \sqrt{n_0(\rho, z)n_0(\rho, z')}$, where n_0 is the (quasi)condensate density. The variance $\Delta \phi^2(Z, s)$, the key quantity to characterize the spatial fluctuations of the phase of the condensate, has been calculated in [2], and an analytical form has been given, which is valid

near the center of the trap (*i.e.* for $Z, s \ll L$, with L the condensate half-length). The first goal of this paper is to find an analytical approximation for the variance $\Delta \phi^2(Z, s)$ valid across the whole sample. This is motivated by the fact that coherence measurements with quasi-condensates [10, 11] are quite sensitive to the inhomogeneity of the sample. In position space, interferometry [12, 13] gives access to the spatial correlation function $\mathcal{C}^{(1)}(s)$ (see *e.g.* [15])

$$\mathcal{C}^{(1)}(s) = \int d^3\mathbf{R} \langle \hat{\Psi}^\dagger(\rho, Z + s/2) \hat{\Psi}(\rho, Z - s/2) \rangle. \quad (2)$$

Equivalently, one can measure the axial (*i.e.* integrated over transverse momenta) momentum distribution $\mathcal{P}(p_z)$, which is the Fourier transform of $\mathcal{C}^{(1)}(s)$ [14, 15, 16, 17]:

$$\mathcal{P}(p_z) = \frac{1}{2\pi\hbar} \int ds \mathcal{C}^{(1)}(s) e^{-ip_z s/\hbar}. \quad (3)$$

A powerful tool to measure $\mathcal{P}(p_z)$ is Bragg spectroscopy with large momentum transfer, as demonstrated in [14] for a 3D condensate, and recently applied in our group to perform the momentum spectroscopy of a quasicondensate [10]. It is clear that both $\mathcal{C}^{(1)}$ and \mathcal{P} are sensitive to the inhomogeneity of the system. Our second goal is to obtain explicit expressions for these two important quantities.

This paper is organized as follows. First, we summarize the results of [2], and give an energetic interpretation of T_ϕ . Next, we discuss in detail a local density approach (LDA) to compute the variance of the phase for any mean position in the trap. This approximation is found to be accurate for $T \geq 8T_\phi$, when applied to a trapped condensate. Using the LDA, we then address the problem of a phase-fluctuating condensate in free expansion. In particular, we point out that at higher temperature, the phase fluctuations dominate over the mean-field release velocity and govern the shape of the momentum distribution.

We consider N_0 condensed atoms, trapped in a cylindrically symmetric, harmonic trap with an aspect ratio $\lambda = \omega_z/\omega_\perp \ll 1$. If $\mu \gg \{\hbar\omega_\perp, \hbar\omega_z\}$, the condensate is in the 3D Thomas-Fermi (TF) regime [18]. The density has the well-known inverted parabola form: $n_0(\mathbf{r}) =$

[‡]UMRA 8501 du CNRS

^{*}e-mail: fabrice.gerbier@iota.u-psud.fr

[†]current address: Department of Physics, University of Toronto, Canada.

$n_{0m}(1 - \tilde{\rho}^2 - \tilde{z}^2)$, with the peak density $n_{0m} = \mu/g$ related to the chemical potential μ . From now on, we use the reduced coordinates $\tilde{\rho} = \rho/R$ and $\tilde{z} = z/L$, with $R^2 = 2\mu/M\omega_{\perp}^2$ and $L^2 = 2\mu/M\omega_z^2$ respectively.

As shown in [1], phase fluctuations in trapped gases are mostly associated with thermally excited, low-energy quasi-particles (the quantum fluctuations are negligible). Under these conditions, the variance $\Delta\phi^2(\tilde{Z}, \tilde{s})$ is

$$\Delta\phi^2(\tilde{Z}, \tilde{s}) \approx \sum_j \frac{2k_B T}{\hbar\omega_j} |\phi_j(\tilde{Z} + \tilde{s}/2) - \phi_j(\tilde{Z} - \tilde{s}/2)|^2, \quad (4)$$

where the sum extends over the 1D axial excitations, with energy $\hbar\omega_j$ and occupation number $N_j \approx k_B T / \hbar\omega_j$ for $N_j \gg 1$. For a 3D condensate in an elongated trap, the amplitude ϕ_j is proportional to a Jacobi polynomial $P_j^{(1,1)}$, and $\omega_j = \omega_z \sqrt{j(j+3)}/2$ [6] for integer j . The explicit result for the variance is then [2]

$$\Delta\phi^2(\tilde{Z}, \tilde{s}) = \frac{T}{T_\phi} f(\tilde{Z}, \tilde{s}), \quad (5)$$

with $f(\tilde{Z}, \tilde{s}) = \sum_j F_j [P_j^{(1,1)}(\tilde{Z} + \tilde{s}/2) - P_j^{(1,1)}(\tilde{Z} - \tilde{s}/2)]^2$, and the coefficients $F_j = (j+2)(j+3/2)/4j(j+1)(j+3)$. Below the characteristic temperature $T_\phi = 15N_0(\hbar\omega_z)^2/32\mu k_B$, the phase profile is almost flat, and the single-particle density matrix (1) is limited by the overlap function χ : therefore the characteristic width of $\mathcal{C}^{(1)}$ (*i.e.* the coherence length) is of order L . On the other hand, if $T \gg T_\phi$ the variance $\Delta\phi^2$ dominates the behavior of $\mathcal{C}^{(1)}$, and the coherence length is substantially smaller than L . Near the center of the trap ($\tilde{Z}, \tilde{s} \ll 1$), Petrov *et al.* [2] have derived the simple law $\Delta\phi^2(\tilde{Z}, \tilde{s}) \approx (T/T_\phi)|\tilde{s}|$, and introduced the characteristic phase coherence length $L_\phi = LT_\phi/T$ that depends implicitly on the temperature, on the number of condensed atoms and on the trapping geometry.

We can understand this expression for L_ϕ from energetic considerations. A random phase gradient of the condensate wavefunction, on a length scale L_ϕ , requires an average kinetic energy $E_\phi \sim N_0 \hbar^2 / ML_\phi^2$. This kinetic energy is supplied by the thermal excitations that drive the fluctuations of the phase [5]. As these excitations are quasi-classical ($N_k \gg 1$), this energy is of order $k_B T$ times the number of relevant modes. In 1D k -space, the distribution of the relevant excitations extends over $\sim 1/L_\phi$, and the spacing between modes is $\sim 1/L$ because of the finite size of the system: this gives L/L_ϕ relevant modes. By equating the two expressions for E_ϕ , we recover finally $L_\phi \sim LN_0(\hbar\omega_z)^2/\mu k_B T$.

As indicated earlier, it is important to take the full spatial dependence of $\Delta\phi^2(\tilde{Z}, \tilde{s})$ into account for quantitative comparison with experiments. In any case, Eq. (5) can be evaluated numerically. However, we gain physical insight with an analytical approach based on the local density approximation (LDA), also used in [3] to calculate the evolution of the density in time-of-flight. This

approximation considers that the condensate is locally equivalent to a homogeneous medium, however with a slowly varying density that depends on the trapping potential. If $T \gg T_\phi$, the coherence length is sufficiently small compared to L , that the LDA is valid for the calculation of correlation properties.

The first step is to consider a finite cylinder of length $2L$, with radial harmonic trapping and periodic boundary conditions along z (and therefore homogeneous axial density). For this geometry, we find in the TF regime: $n_0(\mathbf{r}) = n_{0m}(1 - \tilde{\rho}^2)$ for the condensate wavefunction. Low-lying excitations are found using standard Bogoliubov theory [5], after averaging over the transverse degrees of freedom [6]. The Bogoliubov spectrum for the excitation frequencies is $\omega_k^B = (\omega_k(\omega_k + 2Mc_{1D}^2/\hbar))^{1/2} \approx c_{1D}k$ for small k , with the free particle energy $\hbar\omega_k = \hbar^2 k^2 / 2M$ and the 1D speed of sound $c_{1D} = \sqrt{\mu/2M}$ [19]. The Fourier component for phase fluctuations with wavevector k is

$$\phi_k = \sqrt{\frac{\omega_k^B}{2\omega_k}} \frac{1}{\sqrt{\mathcal{V}}} \approx \sqrt{\frac{Mc_{1D}}{\hbar k}} \frac{1}{\sqrt{\mathcal{V}}}, \quad (6)$$

where the final expression holds for low-lying phonon states ($k \rightarrow 0$), and $\mathcal{V} = 2\pi n_{0m} R^2 L$. In a second step, we take into account the trapping potential by the substitution:

$$\mu \rightarrow \mu - \frac{1}{2} M \omega_z^2 z^2. \quad (7)$$

This implies directly the replacements:

$$\begin{aligned} \text{density :} & \quad n_{0m}(\propto \mu) \rightarrow n_{0m}(1 - \tilde{z}^2) \\ \text{speed of sound :} & \quad c_{1D}(\propto \sqrt{\mu}) \rightarrow c_{1D}\sqrt{1 - \tilde{z}^2} \\ \text{radius :} & \quad R(\propto \sqrt{\mu}) \rightarrow R\sqrt{1 - \tilde{z}^2} \\ \text{half-length :} & \quad L \rightarrow L \end{aligned} \quad (8)$$

With these substitutions, we recover the 3D TF density profile. We require that the excitation frequency $c_{1D}k$ is not modified as well, which implies replacing k with $k(1 - \tilde{z}^2)^{-1/2}$, and using a density of states $\mathcal{N}(k)dk = (L/\pi)(1 - \tilde{z}^2)^{-1/2}dk$. For the position-dependent variance of the phase, we find [20]

$$\Delta\phi^2(\tilde{Z}, \tilde{s}) \approx \frac{T}{T_\phi} \frac{|\tilde{s}|}{(1 - \tilde{Z}^2)^2}. \quad (9)$$

The \tilde{Z} -dependent phase coherence length $L_\phi(1 - \tilde{Z}^2)^2$ appearing in Eq. (9) can be substantially smaller near the edges of the trap than in the center. We will see that this reduces the average coherence length below its value at the center of the trap.

We deduce from Eqs.(1,2,9) the correlation function:

$$\begin{aligned} \mathcal{C}_{\text{trap}, T}^{(1)}(\tilde{s}) \approx & \frac{15N_0}{8} \int_0^{\sqrt{1-\tilde{s}^2/4}} d\tilde{z} (1 - \tilde{z}^2 - \tilde{s}^2/4)^2 \\ & \exp\left(-\frac{T}{2T_\phi} \frac{|\tilde{s}|}{(1 - \tilde{z}^2)^2}\right). \end{aligned} \quad (10)$$

In deriving Eq. (10), we have used the approximation for the overlap function $\chi(\rho, Z, s) \approx (1 - \tilde{\rho}^2 - \tilde{z}^2 - \tilde{s}^2/4)$, valid near the center of the trap. In Fig. 1, we compare the result (10) to the correlation function following from the numerical integration of (5). In the $T = 0$ limit, the correlation function is limited by the overlap χ . Because of the approximate form of χ , our result $\mathcal{C}_{\text{trap}, T=0}^{(1)} \approx N_0(1 - (\tilde{s}/2)^2)^{5/2}$ is about 25% too broad, and one should rather use the gaussian approximation to $\mathcal{C}^{(1)}$ derived in [16]. As T increases, $\mathcal{C}^{(1)}$ turns to an exponential-like function, and our approximation approaches the numerical calculation. For $T > 8T_\phi$, the LDA result is very close to the numerical one (maximum error $\approx 3\%$). For $T \gg T_\phi$, Eq. (10) can be further simplified by keeping the \tilde{s} -dependent term only in the exponential. The Fourier transform then gives the momentum distribution

$$\mathcal{P}_{\text{trap}, T}(p_z) \approx \frac{15N_0 p_\phi}{32\pi} \int_{-1}^1 d\tilde{z} \frac{(1 - \tilde{z}^2)^4}{(1 - \tilde{z}^2)^4 p_z^2 + p_\phi^2/4}, \quad (11)$$

where $p_\phi = \hbar/L_\phi$ is a typical momentum associated with the phase fluctuations. This function is self-similar in p_z/p_ϕ , and approximated to better than 4% by a normalized Lorentzian with a half-width at half-maximum (HWHM) of $\Delta p = 0.67p_\phi$. This Lorentzian shape of the momentum distribution differs qualitatively from the fully coherent case, where it is almost Gaussian and limited by the Heisenberg principle [14]. The increase of the phase fluctuations with increasing T not only broadens the momentum distribution, but also induces the appearance of "wings", that form the "high-energy tail" of the quasicondensate. To quantify the accuracy of our approximation, we have calculated numerically the Fourier transform of the correlation function. We find empirically that the HWHM is accounted for by the formula $\Delta p^2 \approx (2.04\hbar/L)^2 + (0.65\hbar/L_\phi)^2$. The first term corresponds to the Heisenberg-limited momentum width, and the second to the phase fluctuations. For $T \geq 8T_\phi$, the height and width agree to better than 4% with the Lorentzian approximation. For lower T , the overlap function χ still affects the momentum distribution.

Note finally that the momentum distribution is Lorentzian only in the domain $k \ll R^{-1}$. Outside of this region, the 3D nature of the excitations should be taken into account properly. However, this does not affect the quasicondensate peak we are investigating here, but only the much smoother thermal background [14, 16].

The results of the above paragraphs are valid for an equilibrium situation. However, coherence measurements involving Bragg scattering [12, 14] suffer from two major difficulties in a very elongated trap [10]: mean-field broadening of the resonance [14], and elastic scattering from the recoiling atoms and the condensate towards initially empty modes [21]. Both of these problems can be solved by opening the trap abruptly, and letting the BEC expand to decrease its density before measurement. In the remainder of this paper, we discuss how expansion modifies the momentum distribution and the correlation

function, assuming that the expansion time is chosen to be long enough to neglect the collisions.

For a pure, elongated condensate abruptly released from the trap at $t = 0$, the explicit solution was found in [22]. The condensate density keeps its initial Thomas-Fermi shape, with the coordinates re-scaled. The (small) axial momentum from the released mean-field energy is linear in position: $p_z \approx p_{\text{exp}} \tilde{z}$, with $p_{\text{exp}} = (\pi/\sqrt{2})\lambda M c_S$ for $\tau = \omega_\perp t \gg 1$, and $c_S = \sqrt{\mu/M}$ is the 3D speed of sound. The axial momentum distribution mirrors the (integrated) density distribution:

$$\mathcal{P}_{\text{exp}, T=0}(p_z) = \frac{15}{16p_{\text{exp}}} \left(1 - \left(\frac{p_z}{p_{\text{exp}}}\right)^2\right)^2. \quad (12)$$

This expression holds for a pure condensate, at $T = 0$, as indicated. For a phase fluctuating condensate at finite T , it is necessary to consider the time evolution of the fluctuations as well. As shown in [3], the momentum distribution partially converts into density modulations after time of flight. An explicit solution was derived for the density fluctuations in the axially homogeneous case. Using the continuity equation (after radial averaging), we find for $\tau \gg 1$

$$\phi_k(z, \tau) \approx \phi_k(z, 0) \tau^{-(\omega_k^B/\omega_\perp)^2} \cos\left(\frac{\omega_k}{\omega_\perp} \tau\right), \quad (13)$$

If $\omega_z t \ll (\mu/\hbar\omega_z)(T/T_\phi)^2$, then for all $k \lesssim L_\phi^{-1}$, the phase distribution is essentially frozen: $\phi_k(z, \tau) \approx \phi_k(z, 0)$. Physically, this condition states that for such a time of flight, the excitations that have significant contributions to the phase fluctuations have not yet been converted into density modulations. This condition is not at all restrictive for typical experimental parameters [3, 4, 10, 11], and we suppose it is met in the remainder of the paper.

Using the rescaled wavefunction from [22], together with (9), we find the correlation function for the expanding quasicondensate:

$$\mathcal{C}_{\text{exp}, T}^{(1)}(\tilde{s}) \approx \frac{15N_0}{16} \int_{-1}^1 d\tilde{z} (1 - \tilde{z}^2)^2 \exp\left(i \frac{\pi\mu}{\hbar\omega_\perp} \tilde{z}\tilde{s} - \frac{T}{2T_\phi} \frac{|\tilde{s}|}{(1 - \tilde{z}^2)^2}\right). \quad (14)$$

The phase factor in (14) accounts for the local expansion momentum introduced above (recall $\pi\mu/\hbar\omega_\perp \gg 1$). The Fourier transform gives the momentum distribution

$$\mathcal{P}_{\text{exp}, T}(p_z) \approx \frac{N_0}{p_{\text{exp}}} g_{\gamma=p_\phi/p_{\text{exp}}} \left(\frac{p_z}{p_{\text{exp}}}\right). \quad (15)$$

The function g_γ is given by

$$g_\gamma(x) = \frac{15\gamma}{32\pi} \int_{-1}^1 d\tilde{z} \frac{(1 - \tilde{z}^2)^4}{(1 - \tilde{z}^2)^4 (x - \tilde{z})^2 + \frac{\gamma^2}{4}}, \quad (16)$$

and the ratio $\gamma = p_\phi/p_{\text{exp}}$ controls which component of the momentum distribution dominates. In the limit

$T \rightarrow 0$, using $\gamma/(x^2 + \gamma^2) \rightarrow \pi\delta(x)$ as $\gamma \rightarrow 0$, we recover the zero-temperature result Eq. (12). On the other hand, if $\gamma \gg 1$, we expect the momentum distribution to be similar to the distribution in the trap (11). Fig. 2 shows a numerical calculation of g_γ , for various values of γ . We find that already for $p_\phi \gtrsim 2p_{\text{exp}}$, the momentum distribution is almost entirely dominated by phase fluctuations, and, as in the trapped case, is very well approximated by a normalized Lorentzian with HWHM $= 0.67p_\phi$. Here, we note two points: first, that the Heisenberg width $\sim \hbar/L$ is negligible at any temperature, and second that, for large enough condensates, we can have $p_{\text{exp}} \gg p_\phi$ even if the coherence length is smaller than L .

In summary, we have analyzed the measurement of phase fluctuations in elongated Bose condensates. Within a local density approach, we have been able to take the density profile into account, and derived analyt-

tical formulas for the correlation function and the momentum distribution of static and freely expanding quasicondensates. In the regime of interest, the formula compare well to a numerical evaluation based on the results of [2], which are exact in the long-wavelength limit. In particular, we show how the shape of the momentum distribution tends to a Lorentzian with half-width $\approx 0.67\hbar/L_\phi$ as one goes further in the phase-fluctuating regime. We believe that these results may be helpful to understand quantitatively experiments involving quasi-condensates [3, 4, 10, 11].

We gratefully acknowledge D. S. Petrov and G. V. Shlyapnikov for many stimulating discussions, for useful comments on the manuscript and also for providing us with a detailed version of the calculations published in [3]. JHT is supported by a Chateaubriand Fellowship and MH by IXSEA. This work is supported by the EU and DGA.

-
- [1] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Phys. Rev. Lett. **85**, 3745 (2000).
 - [2] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Phys. Rev. Lett. **87**, 050404 (2001).
 - [3] S. Dettmer *et al.*, Phys. Rev. Lett. **87**, 160406 (2001); D. Hellweg *et al.*, Appl. Phys. B **73**, 781 (2001).
 - [4] I. Schvachuck *et al.*, Phys. Rev. Lett. **89**, 270404 (2002).
 - [5] E. M. Lifshitz and L. P. Pitaevskii, *Statistical Physics, part 2* (Butterworth-Heinemann Ltd., London, 1980).
 - [6] S. Stringari, Phys. Rev. A **58**, 2385 (1998).
 - [7] It is necessary to improve Bogolubov theory [8, 9] to obtain the equation of state in any dimensions, in presence of phase fluctuations. For the specific problem we address in this paper, the long wavelength formulation is sufficient.
 - [8] J. O. Andersen, U. Al Khawaja, and H. T. C. Stoof, Phys. Rev. Lett. **88**, 070407 (2002); U. Al Khawaja, J. O. Andersen, N. P. Proukakis, and H. T. C. Stoof, Phys. Rev. A **66**, 013615 (2002).
 - [9] C. Mora and Y. Castin, Phys. Rev. A, 053615 (2003).
 - [10] S. Richard, F. Gerbier, J. H. Thywissen, M. Hugbart, P. Bouyer and A. Aspect, Phys. Rev. Lett **91**, 010405 (2003).
 - [11] D. Hellweg, L. Cacciapuotti, M. Kottke, T. Schulte, K. Sengstock, W. Ertmer and J. J. Arlt, Phys. Rev. Lett **91**, 010406 (2003).
 - [12] E. W. Hagley *et al.*, Phys. Rev. Lett. **83**, 3112 (1999).
 - [13] I. Bloch, T. W. Hänsch, and T. Esslinger, Nature **403**, 166 (2000).
 - [14] J. Stenger, S. Inouye, A. P. Chikkatur, D. M. Stamper-Kurn, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. **82**, 4569 (1999).
 - [15] C. Cohen-Tannoudji and C. Robilliard, C. R. Acad. Sci. Paris, t.2, série IV, 445-477 (2001).
 - [16] F. Zambelli, L. Pitaevskii, D. M. Stamper-Kurn, and S. Stringari, Phys. Rev. A **61**, 063608 (2000).
 - [17] D. Luxat and A. Griffin, Phys. Rev. A **67**, 043603 (2003).
 - [18] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999).
 - [19] E. Zaremba, Phys. Rev. A **57**, 518 (1998); G. M. Kavoulakis and C. J. Pethick, Phys. Rev. A **58**, 1563 (1998).
 - [20] We have also verified that, starting from the large j limit of the Jacobi polynomials, one can recover Eq. (9).
 - [21] A. P. Chikkatur, A. Görlitz, D. M. Stamper-Kurn, S. Inouye, S. Gupta, and W. Ketterle, Phys. Rev. Lett. **87**, 403 (2000).
 - [22] Y. Castin and R. Dum, Phys. Rev. Lett. **77**, 5315 (1996); Yu. Kagan, E. L. Surkov, and G. V. Shlyapnikov, Phys. Rev. A **55**, R18 (1997).




FIG. 1: Spatial correlation function $\mathcal{C}^{(1)}(\tilde{s})$ of a trapped quasicondensate. The solid lines follow from the numerical evaluation of the result (5) derived in [2], for $T = 0$, $T/T_\phi = 4$ and $T/T_\phi = 8$, in order of decreasing width. The dashed lines follow from the local density approximation Eq.(10). Note that for convenience, $\mathcal{C}^{(1)}(\tilde{s})$ as shown here is normalized to 1 rather than to N_0 , differently from what is done in the text.




FIG. 2: The momentum distribution in expansion g_γ (Eq. 16) for $\gamma = p_\phi/p_{\text{exp}} = 0, 1, 3$. As γ increases, g_γ continuously transforms from a quartic profile to a Lorentzian-like profile (see text). For $\gamma = 3$, we find little change from the momentum distribution in the trap (dashed line, a Lorentzian with HWHM 0.67γ , see text). The functions have been rescaled by their maximum values to facilitate comparison.