

# Metal Surface Energy: Persistent Cancellation of Short-Range Correlation Effects beyond the Random-Phase Approximation

J. M. Pitarke<sup>1,2</sup> and J. P. Perdew<sup>3</sup>

<sup>1</sup>Materia Kondentsatuaren Fisika Saila, Zientzi Fakultatea,

Euskal Herriko Unibertsitatea, 644 Posta Kutxatila, E-48080 Bilbo, Basque Country

<sup>2</sup>Donostia International Physics Center (DIPC) and Centro Mixto CSIC-UPV/EHU, Donostia, Basque Country

<sup>3</sup>Department of Physics and Quantum Theory Group, Tulane University, New Orleans, LA 70118

(Dated: March 22, 2024)

The role that non-local short-range correlation plays at metal surfaces is investigated by analyzing the correlation surface energy into contributions from dynamical density fluctuations of various two-dimensional wave vectors. Although short-range correlation is known to yield considerable correction to the ground-state energy of both uniform and non-uniform systems, short-range correlation effects on intermediate and short-wavelength contributions to the surface formation energy are found to compensate one another. As a result, our calculated surface energies, which are based on a non-local exchange-correlation kernel that provides accurate total energies of a uniform electron gas, are found to be very close to those obtained in the random-phase approximation and support the conclusion that the error introduced by the local-density approximation is small.

PACS numbers: 71.15.Mb, 71.45.Gm

## I. INTRODUCTION

The widely-used Kohn-Sham formulation of density-functional theory (DFT)<sup>1</sup> requires approximations to the exchange-correlation (xc) energy  $E^{xc}[n(r)]$ . The simplest approximation to this functional is the so-called local-density approximation (LDA), where  $E^{xc}[n(r)]$  is given at each point by the xc energy of a uniform electron gas at the local density. This approximation was found to be remarkably accurate in some rather inhomogeneous situations,<sup>2</sup> and its widespread use in condensed-matter physics led to the early success of DFT.

Hence, it is important that the LDA be tested against benchmark systems, such as the jellium surface, and that new functionals be developed. Nevertheless, more than 30 years after Lang and Kohn reported the first self-consistent LDA calculation of the jellium surface energy,<sup>3</sup> the question of the impact of non-local xc effects on the surface energy and their interplay with the strong charge inhomogeneity at the surface has remained a puzzle.<sup>4,5</sup> The simple LDA and more advanced density functionals such as generalized gradient approximations (GGA's)<sup>6</sup> and meta-GGA's<sup>7</sup> all predict the same jellium xc surface energy within a few percent, but show no such agreement with the available wave-function based methods: Fermi hypervetted chain (FHNC)<sup>8</sup> and diffusion Monte Carlo (DMC);<sup>9</sup> see Table I of Ref. 10.

An alternative formally exact way to find the xc energy of an arbitrary inhomogeneous system is provided by the adiabatic connection formula and the fluctuation-dissipation theorem.<sup>11</sup> Within this approach, the exchange energy is fully determined from the exact Kohn-Sham (KS) orbitals and the correlation energy is obtained in terms of the xc kernel  $f^{xc}$ .<sup>12</sup> In the random-phase approximation (RPA),  $f^{xc}$  is taken to be zero. Full RPA or corrected-RPA calculations are now feasible not only for bulk jellium<sup>13</sup> but also for jellium surfaces<sup>14</sup> and

molecules.<sup>15,16,17</sup>

In this paper, we take a non-local xc kernel that provides accurate ground-state energies of a uniform electron gas,<sup>13</sup> and evaluate the jellium surface energy through the use of the adiabatic connection formula. We analyze the correlation surface energy into contributions from dynamic density fluctuations of various two-dimensional wave vectors, and find that short-range xc effects on intermediate and short-wavelength fluctuations nearly compensate. Hence, while RPA is known to be a poor approximation for the total correlation energy, our calculations show that it is a surprisingly good approximation for those changes in the correlation energy that arise in surface formation. This is in contrast with FHNC and DMC slab calculations,<sup>8,9</sup> which predict surface energies that are significantly higher than those obtained either in the LDA<sup>3</sup> or in a fully non-local RPA.<sup>14</sup>

The FHNC variational equations have been shown to provide, in the homogeneous limit, reasonable agreement with the known properties of a uniform electron gas,<sup>18</sup> and DMC calculations are often regarded as essentially exact.<sup>19</sup> Furthermore, one may expect that when applied to non-uniform systems these wave-function-based approaches will lead to results whose accuracy is comparable to the high accuracy obtained for uniform systems. Nevertheless, we show that surface formation energies obtained from slab calculations either by a linear fit in the slab thickness<sup>8</sup> or as differences between slab energies and an independently determined bulk energy<sup>9</sup> may result in substantial imprecision, and conclude that wave-function-based estimates need to be reconsidered.

## II. THEORETICAL FRAMEWORK

We consider a jellium slab normal to the  $z$  axis, which is translationally invariant in the surface plane. Subtract-

ing from the slab energy the corresponding energy of a uniform electron gas and using the adiabatic connection formula, one obtains the xc surface energy

$$\epsilon_{xc}^s = \int_0^1 d(\alpha k_F) \epsilon_{q,\alpha}^{xc}; \quad (1)$$

$q$  being a wave vector parallel to the surface.  $k_F$  is the Fermi momentum and  $\epsilon_{q,\alpha}^{xc}$  represents the surface energy associated with the good quantum number  $q$ :

$$\epsilon_{q,\alpha}^{xc} = \frac{k_F}{8} \int_0^1 dz \int_0^1 dz^0 n(z) v_q(z-z^0) \int_0^1 d n_{q,\alpha}^{xc}(z; z^0) n_{q,\alpha}^{xc}(z-z^0); \quad (2)$$

$v_q(z-z^0) = (2\pi)^{-3} \exp(iq(z-z^0))$  being the Fourier transform of the bare Coulomb interaction.  $n_{q,\alpha}^{xc}(z; z^0)$  and  $n_{q,\alpha}^{xc}(z-z^0)$  represent Fourier components of the xc-hole density of a fictitious jellium slab at coupling strength  $\alpha$  and the corresponding xc-hole density of a uniform electron gas of density  $n = k_F^3/3\pi^2$ , respectively. In the LDA, the xc surface energy is obtained by simply replacing  $n_{q,\alpha}^{xc}(z; z^0)$  by the xc-hole density of a uniform electron gas of density  $n(z)$ . A parametrization of the uniform-gas xc-hole density has been given by Perdew and Wang,<sup>20</sup> which yields the DMC ground-state energy of a uniform electron gas.<sup>21</sup>

According to the fluctuation-dissipation theorem,

$$n_{q,\alpha}^{xc}(z; z^0) = \frac{\hbar}{n(z)} \int_0^1 d\omega \chi_{q,\alpha}(z; z^0; i\omega) \quad (z-z^0); \quad (3)$$

where  $\chi_{q,\alpha}(z; z^0; i\omega)$  is the interacting density-response function. Time-dependent DFT (TDDFT) shows that this function obeys the Dyson-type equation<sup>22,23</sup>

$$\chi_{q,\alpha}(z; z^0; i\omega) = \chi_q^0(z; z^0; i\omega) + \int dz_1 \int dz_2 \chi_q^0(z; z_1; i\omega) v_q(z_1 - z_2) + f_{q,\alpha}^{xc}[n](z_1; z_2; i\omega) \chi_{q,\alpha}(z_2; z^0; i\omega); \quad (4)$$

where  $\chi_q^0(z; z^0; i\omega)$  is the density-response function of non-interacting KS electrons<sup>24</sup> and  $f_{q,\alpha}^{xc}[n](z; z^0; i\omega)$  involves

the functional derivative of the KS xc potential at coupling constant  $\alpha$ .

Using the coordinate-scaling relation for the  $\alpha$ -dependence of the xc kernel derived in Ref. 13, we find

$$f_{q,\alpha}^{xc}[n](z; z^0; i\omega) = f_{q,\alpha=1}^{xc}[\alpha^{-3}n(z=\alpha z)](\alpha z; \alpha z^0; i\omega); \quad (5)$$

where  $f_{q,\alpha=1}^{xc}[n](z; z^0; i\omega)$  is the xc kernel at  $\alpha = 1$ . In order to derive an approximation for this quantity, we assume that the density variation  $[n(z) - n(z^0)]$  is small within the short range of  $f_{q,\alpha=1}^{xc}[n](z; z^0; i\omega)$  and write<sup>13</sup>

$$f_{q,\alpha=1}^{xc}[n](z; z^0; i\omega) = f_{q,\alpha=1}^{xc}([n(z) + n(z^0)]/2; z-z^0; i\omega); \quad (6)$$

where  $f_{q,\alpha=1}^{xc}(n; z-z^0; i\omega)$  is the Fourier transform of the xc kernel  $f_{q,\alpha=1}^{xc}(n; k; i\omega)$  of a uniform electron gas of density  $n$ . Here  $k = (q; k_z)$  represents a three-dimensional wave vector.

### III. RESULTS AND DISCUSSION

We have carried out simplified surface-energy calculations with  $f_{q,\alpha}^{xc}(n; z-z^0; i\omega)$  replaced by  $f_{q,\alpha}^{xc}(n; k = q; i\omega) \delta(z-z^0)$  [thus assuming that the dynamic density fluctuation is slowly varying in the direction perpendicular to the surface] and using the parametrization of Richardson and Ashcroft for  $f_{q,\alpha}^{xc}(n; k; i\omega)$ ,<sup>26</sup> and have found that neglect of the frequency dependence of the xc kernel does not introduce significant errors. We have also carried out adiabatic LDA (ALDA) surface-energy calculations with  $f_{q,\alpha}^{xc}(n; z-z^0; i\omega)$  replaced by  $f_{q,\alpha}^{xc}(n; k = 0; i\omega) \delta(z-z^0)$  [thus assuming that the dynamic density fluctuation is slowly varying in all directions], and have found that the spacial range of the xc kernel cannot be neglected. These conclusions also apply to the uniform electron gas.<sup>13</sup>

Hence, we neglect the frequency dependence of the xc kernel and exploit the accurate DMC calculations reported in Ref. 27 for the static xc kernel of a uniform electron gas. A parametrization of this data satisfying the known small- and large-wavelength asymptotic behavior has been reported,<sup>28</sup> which allows us to write

$$f_{q,\alpha=1}^{xc}(n; z-z^0) = \frac{4\pi e^2 C}{k_F^2} (z-z^0) \exp\left[-\frac{2\pi e^2 B}{gk_F^2 + q^2} \exp\left(\frac{P}{gk_F^2 + q^2} |z-z^0|\right)\right] \frac{2\pi e^2}{k_F^3} \left[ \frac{2}{4} \frac{k_F^2 z^2}{k_F^2 + q^2} + q^2 \exp\left[-\frac{k_F^2 z^2}{4} - \frac{q^2}{k_F^2}\right] \right]; \quad (7)$$

where  $C, B, g, P$ , and  $\alpha$  are dimensionless functions of the electron density (see Ref. 28),  $n = k_F^3/3\pi^2$ , and  $z = z - z^0$ . The finite  $q \rightarrow 0$  limit of Eq. (7) will be dominated by the  $q \rightarrow 0$  divergence of  $v_q(z-z^0)$ , making RPA exact in this limit. In the large- $q$  limit, where short-wavelength

excitations tend to be insensitive to the electron-density inhomogeneity, introduction of Eq. (7) into Eq. (6) is expected to yield an xc kernel that is essentially exact.

If the interacting density-response function  $\chi_{q,\alpha}(z; z^0; i\omega)$  entering Eq. (3) is replaced by  $\chi_q^0(z; z^0; i\omega)$ ,




FIG. 1: Wave-vector analysis  $\gamma_q^c$  of the correlation surface energy for a jellium slab of thickness  $a = 7.17r_s$  and  $r_s = 2$ . Thin-solid, dashed, and dotted lines represent Perdew-Wang, RPA and Corradini-based LDA calculations, respectively. The thick-solid line and the open circles represent RPA and Corradini-based non-LDA calculations, respectively. Our 'best' non-local calculation (open circles) provides the exact small- $q$  and large- $q$  limits.

Eqs. (1) and (2) yield the exact exchange surface energy, as obtained in Ref. 14. Here we focus our attention on the correlation surface energy, which for comparison we also calculate in LDA by replacing  $n_{q_i}^c$  in Eq. (2) by the uniform-gas correlation-hole density at the local density  $n(z)$ .

Fig. 1 shows the wave-vector analysis  $\gamma_q^c$  of both LDA and non-LDA correlation surface energies of a jellium slab of thickness  $a = 7.17r_s$  and  $r_s = 2$ .<sup>29</sup> First of all, we focus on our LDA calculations, which have been carried out either by using the uniform-gas correlation-hole density with  $f_q^{xc}(n; \vec{r} = z^0) = 0$  (RPA-based LDA) or the xc kernel of Eq. (7) (Corradini-based LDA), or by using the Perdew-Wang (PW) parametrization of Ref. 20. We observe that in the long-wavelength limit ( $q \rightarrow 0$ ) both RPA and Corradini calculations coincide with the PW parametrization. At shorter wavelengths, the Corradini scheme predicts a substantial correction to the RPA and accurately reproduces the PW wave-vector analysis of the correlation energy. Hence, armed with some confidence in the accuracy of our choice of the xc kernel,

we apply it to the more realistic non-local scheme described above. Fig. 1 shows that our non-LDA beyond-RPA ('best' non-local) calculation coincides in the  $q \rightarrow 0$  limit<sup>30</sup> with the non-LDA RPA, which is exact in this limit. In the large- $q$  limit, local and non-local calculations coincide, and our 'best' non-local calculation accurately reproduces the PW-based LDA (thin solid line), which is expected to be essentially exact in this limit.

Consequently, our 'best' non-local calculation provides both the exact small- $q$  limit, where LDA fails badly, and the exact large- $q$  limit, where RPA is wrong. The LDA largely underestimates our non-local correlation surface

TABLE I: Non-local xc ( $^{xc}$ ) and total ( $^{\text{tot}}$ ) surface energies and their local (LDA) counterparts, as obtained from Eqs. (1) and (2) with the non-local xc kernel of Eq. (7). Also shown are non-local RPA xc surface energies ( $^{xc}_{\text{RPA}}$ ) and PW-LDA total surface energies ( $^{\text{tot}}_{\text{PW-LDA}}$ ). Tiny differences between these RPA xc surface energies and those reported before<sup>14</sup> are entirely due to differences in the parametrization of the xc potential. Units are erg/cm<sup>2</sup>.

$r_s$	$^{xc}_{\text{RPA}}$	$^{xc}$	$^{xc}_{\text{LDA}}$	$^{\text{tot}}$	$^{\text{tot}}_{\text{LDA}}$	$^{\text{tot}}_{\text{PW-LDA}}$
2.00	3467	3466	3369	-752	-849	-862
2.07	3064	3063	2975	-504	-592	-605
2.30	2098	2096	2026	-27	-97	-103
2.66	1242	1239	1193	221	175	171
3.00	803	797	767	258	228	225
3.28	580	577	551	247	221	220
4.00	279	277	262	179	164	164
5.00	120	119	111	106	98	98
6.00	59	58	54	64	59	59

energy, but Fig. 1 shows that the difference between RPA and beyond-RPA  $\gamma_q^c$  (the short-range part of the correlation surface energy) is fairly insensitive to whether the LDA is used or not. This supports the assumption made in Ref. 31 that the short-range part of the correlation energy can be treated within LDA or GGA. However, short-range xc effects on intermediate and short-wavelength contributions to the surface energy tend to compensate, and this cancellation happens to be even more complete than expected from LDA or GGA. Thus, our non-local scheme yields surface energies which are still closer to RPA than is the RPA<sup>+</sup> of Ref. 31.

To extract the surface energy of a semi-infinite medium, we have considered three different values of the slab thickness: the threshold width at which the  $n = 5$  subband for the  $z$  motion is completely occupied and the two widths at which the  $n = 5$  and  $n = 6$  subbands are half occupied, and have followed the extrapolation procedure of Ref. 14. In Table I we show our extrapolated local (LDA) and non-local surface energies, as obtained from Eqs. (1) and (2) either with  $f_q^{xc}(n; \vec{r} = z^0) = 0$  (RPA) or with the xc kernel of Eq. (7). These calculations indicate that the introduction of a plausible non-local xc kernel yields short-range corrections to RPA surface energies that are negligible. For comparison, also shown in Table I are PW-LDA surface energies, as obtained either from Eqs. (1) and (2) with the PW uniform-gas xc-hole density<sup>20</sup> or from the PW parametrization of the uniform-gas xc energy.<sup>25</sup> Corradini and PW-based LDA surface energies ( $^{\text{tot}}_{\text{LDA}}$  and  $^{\text{tot}}_{\text{PW-LDA}}$ ) are found to be very close to each other, and we expect our Corradini-based non-local surface energies ( $^{\text{tot}}$ ) to be close to the exact jellium surface energy, as well.

We close this paper with an analysis of the available wave-function-based surface-energy calculations. Krotscheck et al.<sup>39</sup> considered slabs of four different thickness  $a$ , and obtained both a bulk energy per particle  $\epsilon_1$  and a surface energy  $\gamma$  from the FHNC slab

energy per particle  $\epsilon$  as a function of the particle number per unit area  $n_a$ , by a linear fit  $\epsilon(n_a) = \epsilon_1 + 2\epsilon_2 n_a$  that becomes exact in the limit of infinite thickness. These authors showed that the extrapolated bulk energies ( $\epsilon_1$ ) and those obtained from a separate FHNC bulk calculation ( $\epsilon$ ) agree within about 1%, and claimed that this comparison lent credibility to their numerical treatment. However, these small differences in the bulk calculation yield an uncertainty in the surface energy  $\epsilon_s = (\epsilon_1 - \epsilon_2)n_a = 2$ , which for  $r_s = 2.07$  and  $4.96$  can be as large as  $280$  and  $11$  erg/cm<sup>2</sup>, respectively. Moreover, since for the smallest/largest width under study and due to oscillatory quantum-size effects the quantity  $\epsilon(n_a)$  is larger/smaller than expected for a semi-infinite medium, the extrapolated bulk and surface energies are found to be too negative and too large, respectively.

Liet al.<sup>32</sup> calculated the fixed-node DMC surface energy of a jellium slab with  $r_s = 2.07$  and found  $\epsilon_s = 465$  erg/cm<sup>2</sup>, which is  $40$  erg/cm<sup>2</sup> larger than the RPA value. They also performed LDA calculations with either the Wigner or the Ceperley-Alder form for the uniform-gas xc energy, and found LDA surface energies that are also about  $40$  erg/cm<sup>2</sup> larger than the corresponding LDA surface energies of a semi-infinite jellium, which suggests that finite-size corrections might bring the DMC surface energy into close agreement with RPA. These fixed-node DMC calculations were extended by Acioli and Ceperley to study jellium slabs at various different densities,<sup>9</sup> but these authors extracted the surface energy from release-node bulk energies. Both Liet al.<sup>32</sup> and Acioli and Ceperley<sup>9</sup> claimed that the release-node correction of the uniform electron gas at  $r_s = 2.07$  is  $0.0023$  eV/electron, and argued that this correction would only yield a small error in the surface energy. However, the unpublished uniform-gas fixed-node energy reported and used in Ref. 32 is actually  $0.0123$  eV/electron higher than its release-node counterpart; hence, by combining fixed-node slab and release-node bulk energies Acioli and Ceperley produced for  $r_s = 2.07$  a surface energy that is too large by  $138$  erg/cm<sup>2</sup>. Furthermore, had these authors used fixed-node bulk energies (see, e.g., Ref. 33), they would have obtained surface energies that are close to RPA.

#### IV. SUMMARY AND CONCLUSIONS

We have investigated the role that short-range correlation plays at metal surfaces, on the basis of a wave-vector analysis of the correlation surface energy. Our non-local calculations, which are found to provide the exact small and large- $q$  limits, indicate that a persistent cancellation of short-range correlation effects yields surface energies that are in excellent agreement with RPA, and support the conclusion that the error introduced by the LDA is small. Although this conclusion seems to be in contrast with available wave-function based calculations, we have shown that a careful analysis of these data might

bring them into close agreement with RPA. This is consistent with recent work, where jellium surface energies extracted from DMC calculations for jellium spheres are also found to be close to RPA.<sup>34</sup>

We have found that the RPA xc surface energy displays an error cancellation between short- and intermediate-range correlations. A different (and less complete) error cancellation between long- and intermediate-range xc effects explains<sup>10</sup> why the LDA works for the surface energy. The GGA corrects only the intermediate-range contributions,<sup>10</sup> and so gives surface energies slightly lower and less accurate than those of LDA. Usually GGA works better than LDA, but not for the jellium surface energy where long-range effects are especially important. In the present work, as in four others,<sup>10,31,34,35</sup> we have found that the jellium xc surface energy is only a few percent higher than it is in LDA. These closely-agreeing methods include two different short-range corrections to RPA (present work using a non-local xc kernel and Ref. 31 using an additive GGA correction), a long-range correction to GGA (Ref. 10), extraction of a surface energy from DMC energies for jellium spheres (Ref. 34), and a meta-GGA density functional (Ref. 7). The corresponding correction to the LDA or GGA surface energy has been transferred<sup>36</sup> successfully to the prediction of vacancy formation energies<sup>37</sup> and works of adhesion.<sup>38</sup>

We have almost reached a solution of the surface-energy puzzle, but one piece still does not fit. Krotscheck and Kohn<sup>39</sup> examined a "collective RPA" which approximates our full RPA, and also used several xc kernels to correct for short-range effects. When they used an isotropic xc kernel derived from the uniform gas, in the spirit of our Eq. (6), they found surface energies very close to RPA, as we do. When they used FHNC,<sup>8</sup> corresponding to an anisotropic (but to the eye not very different) kernel constructed explicitly for the jellium surface, they found a large positive correction to the RPA surface energy, amounting at  $r_s = 4$  to as much as 35% of the RPA xc or 60% of the RPA total. The story of the jellium surface energy cannot reach an end until their work is reconciled with the other work on this subject. The DMC energies of jellium slabs should also be re-considered.<sup>40</sup>

Measured surface energies of real metals have been compared with calculations in Refs. 41,42,43. However, experimental and calculational uncertainties and differences between jellium and real metal surfaces seem to preclude a solution to the surface-energy puzzle from these comparisons.

#### Acknowledgments

We wish to acknowledge partial support by the University of the Basque Country, the Spanish Ministerio de Educación y Cultura, and the U.S. National Science Foundation (Grants Nos. DMR-9810620 and DMR-0135678).

- <sup>1</sup> W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
- <sup>2</sup> O. Gunnarsson and B. I. Lundqvist, *Phys. Rev. B* **13**, 4274 (1976).
- <sup>3</sup> N. D. Lang and W. Kohn, *Phys. Rev. B* **1**, 4555 (1970).
- <sup>4</sup> M. Nekovee and J. M. Pitarke, *Comp. Phys. Commun.* **137**, 123 (2001).
- <sup>5</sup> See, e.g., Strong coulomb correlations in electronic structure calculations: beyond the local density approximation, edited by V. I. Anisimov (Gordon and Breach, Amsterdam, 2000).
- <sup>6</sup> J. P. Perdew, K. Burke, and M. Ernzerhof, *Phys. Rev. Lett.* **77**, 3865 (1996).
- <sup>7</sup> J. P. Perdew, S. Kurth, A. Zupan, and P. Blaha, *Phys. Rev. Lett.* **82**, 2544 (1999).
- <sup>8</sup> E. Krotscheck, W. Kohn, and G.-X. Qian, *Phys. Rev. B* **32**, 5693 (1985).
- <sup>9</sup> P. H. Pleschl and D. M. Ceperley, *Phys. Rev. B* **54**, 17199 (1996).
- <sup>10</sup> Z. Yan, J. P. Perdew, S. Kurth, C. Fiolhais, and L. Almeida, *Phys. Rev. B* **61**, 2595 (2000).
- <sup>11</sup> D. C. Langreth and J. P. Perdew, *Solid State Commun.* **17**, 1425 (1975); *Phys. Rev. B* **15**, 2884 (1977).
- <sup>12</sup> E. K. U. Gross and W. Kohn, *Phys. Rev. Lett.* **55**, 2850 (1985).
- <sup>13</sup> M. Lein, E. K. U. Gross, and J. P. Perdew, *Phys. Rev. B* **61**, 13431 (2000).
- <sup>14</sup> J. M. Pitarke and A. G. Eguiluz, *Phys. Rev. B* **57**, 6329 (1998); **63**, 45116 (2001).
- <sup>15</sup> F. Furche, *Phys. Rev. B* **64**, 195120 (2001).
- <sup>16</sup> M. Fuchs and X. Gonze, *Phys. Rev. B* **65**, 235109 (2002).
- <sup>17</sup> F. Aryasetiawan, T. Miyake, and K. Terakura, *Phys. Rev. Lett.* **88**, 166401 (2002).
- <sup>18</sup> E. Krotscheck, *Ann. Phys. (N.Y.)* **155**, 1 (1984).
- <sup>19</sup> W. M. C. Foulkes, L. Mitás, R. J. Needs, and G. Rajagopal, *Rev. Mod. Phys.* **73**, 33 (2001).
- <sup>20</sup> J. P. Perdew and Y. Wang, *Phys. Rev. B* **46**, 12947 (1992).
- <sup>21</sup> D. M. Ceperley and B. J. Alder, *Phys. Rev. Lett.* **45**, 566 (1980).
- <sup>22</sup> M. Petersilka, U. J. Gossmann, and E. K. U. Gross, *Phys. Rev. Lett.* **76**, 1212 (1996).
- <sup>23</sup> J. F. Dobson and J. Wang, *Phys. Rev.* **62**, 10038 (2000).
- <sup>24</sup> Although  $\epsilon_0$  should be determined from the exact KS eigenfunctions and eigenvalues, the error introduced by our use of LDA orbitals is expected to be negligible. Whenever the uniform-gas xc energy is required as input, we use the parameterization by Perdew and Wang<sup>25</sup> of the Ceperley-Alder DMC data.<sup>21</sup>
- <sup>25</sup> J. P. Perdew and Y. Wang, *Phys. Rev. B* **45**, 13244 (1992).
- <sup>26</sup> C. F. Richardson and N. W. Ashcroft, *Phys. Rev. B* **50**, 8170 (1994).
- <sup>27</sup> S. Moroni, D. M. Ceperley, and G. Senatore, *Phys. Rev. Lett.* **75**, 689 (1995).
- <sup>28</sup> M. Corradini, R. Del Sole, G. Onida, and M. Palumbo, *Phys. Rev. B* **57**, 14569 (1998).
- <sup>29</sup> The electron-density parameter  $r_s$  is defined by the relation  $1/n = (4\pi/3) r_s^3$ .
- <sup>30</sup> We have found that as  $q \rightarrow 0$  the positive contribution to the correlation surface energy from the creation of electron-hole pairs is increasingly compensated by the corresponding negative exchange surface energy. As a result, the narrow peak that  $\epsilon_q^c$  exhibits at small  $q$  is not present in  $\epsilon_q^{xc}$ , which in the  $q \rightarrow 0$  limit is largely dominated by the shift in plasmon zero-point energies  $\epsilon_q^{xc} - \epsilon_q^c = \frac{1}{2} \omega_p^2 - \frac{1}{2} \omega_p^2 = 0$  (see, e.g. Ref. 11),  $\omega_p = (4\pi n)^{1/2}$  being the bulk-plasmon frequency. We also note that large and opposite separate long-range (small- $q$ ) corrections to the LDA for exchange and correlation are known to compensate partially, as discussed earlier,<sup>11,14</sup> making the LDA a much better approximation for  $\epsilon^{xc}$  than for  $\epsilon^c$  alone.
- <sup>31</sup> Z. Yan, J. P. Perdew, and S. Kurth, *Phys. Rev. B* **61**, 16430 (2000).
- <sup>32</sup> X.-P. Li, R. J. Needs, R. M. Martin, and D. M. Ceperley, *Phys. Rev. B* **45**, 6124 (1992).
- <sup>33</sup> Y. Kwon, D. M. Ceperley, and R. M. Martin, *Phys. Rev. B* **58**, 6800 (1998).
- <sup>34</sup> L. M. Almeida, J. P. Perdew, and C. Fiolhais, *Phys. Rev. B* **66**, 75115 (2002).
- <sup>35</sup> S. Kurth, J. P. Perdew, and P. Blaha, *Int. J. Quantum Chem.* **75**, 889 (1999).
- <sup>36</sup> A. E. Mattsson and W. Kohn, *J. Chem. Phys.* **115**, 3441 (2001).
- <sup>37</sup> K. Carling, G. Wahnstrom, T. R. Mattsson, A. E. Mattsson, N. Sandberg, and G. Grimvall, *Phys. Rev. Lett.* **85**, 3862 (2000).
- <sup>38</sup> A. E. Mattsson and D. R. Jennison, *Surf. Sci. Lett.* (to appear).
- <sup>39</sup> E. Krotscheck and W. Kohn, *Phys. Rev. Lett.* **57**, 862 (1986).
- <sup>40</sup> W. M. C. Foulkes and B. Wood, private communication.
- <sup>41</sup> L. Vitos, A. V. Ruban, H. L. Skriver, and J. Kollar, *Surf. Sci.* **411**, 186 (1998).
- <sup>42</sup> T. L. Atinsworth and E. Krotscheck, *Phys. Rev. B* **45**, 8779 (1992).
- <sup>43</sup> C. Fiolhais and J. P. Perdew, *Phys. Rev. B* **45**, 6207 (1992).