Russian Doll Renorm alization G roup and Superconductivity

Andre LeClair^y, Jose Mar a Roman and German Sierra Instituto de F sica Teorica, UAM/CSIC, Madrid, Spain. (Dated: November, 2002)

We show that an extension of the standard BCS Ham iltonian leads to an in nite number of condensates with di erent energy gaps and self-sim ilar properties, described by a cyclic RG ow of the BCS coupling constant which returns to its original value after a nite RG time.

PACS num bers: 11.10 H i, 74.20 F g, 71.10 L i

The Renorm alization G roup (RG) continues to be one of the most important tools for studying the qualitative and quantitative properties of quantum eld theories and m any-body problem s in C ondensed M atter physics. The emphasis so far has been mainly on ows toward xed points in the UV or IR. Recently, an entirely novel kind of RG ow has been discovered in a number of systems wherein the RG exhibits a cyclic behavior: after a nite RG transform ation the couplings return to their original values and the cycle repeats itself. Thus if one decreases the size of the system by a specic factor that depends on the coupling constants, one recovers the initial system, much like a Russian doll, or quantum version of the M andelbrot set. Bedaque, Hammer and Van Kolck observed this behavior in a 3-body ham iltonian of interest in nuclear physics [1]. This motivated G lazek and W ilson to de ne a very sim ple quantum -m echanical ham iltonian with similar properties [2]. In the meantime such behavior was proposed for a certain regime of an isotropic current-current interactions in 2 dim ensional quantum eld theory [3].

The models in [1, 2] are problem s in zero-dimensional quantum mechanics, and are thus considerably simpler than the quantum eld theory in β]. In the latter, standard quantum eld theory methods of the renorm alization group were used, however know ledge of the beta function to all orders was necessary to observe the cyclic ow. W hat is somewhat surprising is that the model considered in [3] is not very exotic, and is in fact a wellknown theory that arises in m any physical problem s: at one-loop it is nothing m ore than the fam ous K osterlitz-Thouless RG ow, where the cyclic regime corresponds to $jg_{?} j > jg_{k} j$. This motivated us to nd a simpler m any-body problem that captures the essential features of the cyclic RG behavior. We found that a sim ple extension of the BCS ham iltonian has the desirable properties. Namely, ourm odel is based on the BCS ham iltonian with scattering potential V_{jj^0} equal to g + i for $"_j > "_{j^0}$ and i for $"_{1} < "_{1^{\circ}}$ in units of the energy spacing q

The main features of the spectrum are the following. For large system size, there are an in nite number of BCS condensates, each characterized by an energy gap

 $_{n}$ which depends on g, . The role of these m any condensates becomes clearer when we investigate the RG properties. As in the models considered in [1, 2, 3], the

RG ow possesses jumps from g = +1 to g = -1 and a new cycle begins. Let $L = e^{-s}L_0$ denote the RG scale, which in our problem corresponds to N the number of unperturbed energy levels, and the period of an RG cycle: g(e - L) = g(L). We show that = -.

The model we shall consider is an extension of the reduced BCS model used to describe ultrasmall superconducting grains [4], although our results are valid for more general cases. Let c_{j}^{v} (c_{j} ;) denote creationannihilation operators for electrons in tim e reversal states j i. The index j = 1; ::; N refers to N equally spaced energy levels " $_{i}$ with ! < " $_{i}$ < !. The energy " $_{i}$ represents the energy of a pair of electrons in a given level. The level spacing will be denoted 2 , i.e. $"_{j+1}$ $''_1 = 2$, so that ! = N is twice the Debye energy. The energies in this paper are twice their standard values [5]. Let $b_j = c_{j;} c_{j;+}, b_j^y = c_{j;+}^y c_{j;+}^y$ denote the usual Cooperpair operators. The H ilbert space H $_{\rm N}$ is spanned by the com bination of empty and occupied states. At half-lling the dimension of the Hilbert space is the combinatorial num ber $C_{N=2}^{N}$.

Ourmodel is de ned by the reduced BCS ham iltonian

$$H = \bigcup_{j=1}^{X^{N}} b_{j}^{y} b_{j} \qquad \qquad X^{N} V_{jj^{0}} b_{j}^{y} b_{j^{0}}; \qquad (1)$$

where V_{jj^0} is the scattering potential. In the usualBCS model, V_{jj^0} is taken to be a constant. Here we add an imaginary part which breaks time reversal,

$$V_{jj^{0}} = \begin{cases} G + i & \text{if } "_{j} > "_{j^{0}} \\ G & \text{if } "_{j} = "_{j^{0}}; \\ G & i & \text{if } "_{j} < "_{j^{0}} \end{cases} \qquad G = g = (2)$$

T his ham iltonian is herm it ian since V_{jj^0} = $\,V_{j^0j}$. We consider the positive dimensionless couplings g and $\,$.

The BCS variational ansatz for this model is

$$j_{BCS}i = u_j + v_j b_j^V$$
 (3)

The mean-eld treatment yields the well-known equations

$$u_{j}^{2} = \frac{1}{2} + \frac{j}{E_{j}}; \quad v_{j}^{2} = \frac{1}{2}e^{2ij} + \frac{j}{E_{j}};$$
$$q - \frac{q}{2} + \frac{2}{j}; \quad j = \mathbf{y}, \quad Y_{j}; \quad (4)$$

where j and j satisfy the gap equation:

$$e_{j} = \sum_{\substack{i \neq 0 \\ j \neq 0 \\ j \neq 0}}^{X} V_{jj^{0}} \frac{e_{j^{0}}}{E_{j^{0}}}; \quad e_{j} \quad j e^{i_{j}}: \quad (5)$$

The chem ical potential equation in our case is satis ed with = 0. In the therm odynam ic limit, N $\,!\,$ 1 and $\,!\,$ 0, with xed ! = N , the sum s over "j become the integrals $\,_!\,$ d"=2 . The gap equation turns into

$$e(\mathbf{n}) = g \begin{bmatrix} Z & i \\ i \end{bmatrix} \frac{d\mathbf{n}^{0}}{2} \frac{e(\mathbf{n}^{0})}{E(\mathbf{n}^{0})} + i \begin{bmatrix} Z & i \end{bmatrix} \begin{bmatrix} Z & i \end{bmatrix} \frac{d\mathbf{n}^{0}}{2} \frac{e(\mathbf{n}^{0})}{E(\mathbf{n}^{0})}; (6)$$

where $e(") = () e^{i} (")$.

D i erentiating (6) with respect to "yields

$$\frac{\mathrm{d}}{\mathrm{d}^{\mathbf{u}}} = \frac{1}{\mathrm{E}(\mathbf{u})}; \tag{7}$$

and the condition that (") = is independent of ". The solution to eq. (7) can be taken to be

$$(") = \sinh^{-1} \frac{"}{-1}$$
: (8)

U sing eq. (8) in the gap equation (6) gives

$$1 = \int_{0}^{Z} \frac{(!)}{d} (g\cos + \sin) = \tan (!) = -; (9)$$

Solving eq. (9) for the gap yields an in nite number of solutions n. They can be parameterized as follows:

$$n = \frac{!}{\sinh t_n}; \quad t_n = t_0 + \frac{n}{-}; \quad n = 0;1;2;...; \quad (10)$$

where t_0 is the principal solution to the equation

$$\tan(t_0) = \frac{1}{g}; \quad 0 < t_0 < \frac{1}{2}; \quad (11)$$

The gaps satisfy $_0 > _1 > ...$ Each gap represents a di erent BCS eigenstate j $_{BCS}^{(n)}$ i. One can show that $h _{BCS}^{(n)}$ j $_{BCS}^{(n^0)}$ ij< exp N ($_n _{n^0}$)²=8!², in the lim it where $_n$!. Thus in the large N lim it, these eigenstates are orthogonal and should all appear in the spectrum, together with the usual quasi-particle excitations above them . In the lim it ! 0 the gaps $_{n>0}$! 0, and since $t_0 = 1$ =g, $_0$ 2! e ^{1=g}, in the weak coupling regim e, recovering the standard BCS result.

For weak coupling models $_{n}$!, all the gaps are related by a scale transformation $_{n}$ 2N e^{to n =}. Therefore, de ning the condensation energy of the n-th BCS eigenstate as $E_{c}^{(n)} = h_{BCS}^{(n)}$ $\sharp j_{BCS}^{(n)}$ $i \in E_{FS}$ we get

$$E_{c}^{(n)} = \frac{2}{8} = E_{c}^{(n)} = \frac{1}{2} N^{2} e^{2t_{0}} = (12)$$

Thus the spectrum of condensation energies relects the scaling behavior of the gaps.

Next we derive RG equations for our model. Let g_N , N denote the couplings for the ham iltonian H_N with N energy levels. The idea behind the RG method is to derive an elective ham iltonian H_N 1 depending on renormalized couplings g_N 1, N 1 by integrating out the highest energy levels "N or "1. This can be accomplished by a canonical transformation, which is form ally analogous to the one used to derive the t J model from the Hubbard model at strong coupling [6].

We perform the calculation for general V . The integration of the level ${\tt "}_{\rm N}$ yields

$$V_{jj^{0}}^{(N-1)} = V_{jj^{0}}^{(N)} + \frac{1}{2}V_{jN}^{(N)}V_{Nj^{0}}^{(N)} - \frac{1}{N-j} + \frac{1}{N-j^{0}};$$
(13)

where $j = "_j$ V_{jj} . Integration of the level "_1 gives the same eq. (13) with the replacement N j ! 1 + j.

Specializing to the potential eq. (2) and approximating " $_{N}$ "; or " $_{1}$ +"; by ! = N, the above equation implies

$$g_{N-1} = g_{N} + \frac{1}{N} (g_{N}^{2} + g_{N}^{2}); \qquad N = N :$$
 (14)

Thus is unrenormalized.

In the large N $\,$ lim it one can de ne a variable s = $\log N_{\,0} = \!\! N$, where N $_{0}$ is the initial size of the system . Then the beta function reads

$$\frac{dg}{ds} = (g^2 + {}^2); \quad s \quad \log \frac{N_0}{N}:$$
 (15)

The solution to the above equation is

$$g(s) = \tan^{h} s + \tan^{1} \frac{g_{0}}{2}$$
; $g_{0} = g(N_{0})$: (16)

The main features of this RG ow are the cyclicity

$$g(s +) = g(s)$$
 () $g(e N) = g(N);$ -; (17)

and the jum ps from +1 to 1, when reducing the size.

The cyclicity of the RG has some important implications for the spectrum. Let fE (g; ;N)g denote the energy spectrum of the ham iltonian H_N . The RG analysis implies we can compute this spectrum using the ham iltonian $H_{N^0}(g(N^0))$ if $g(N^0)$ is related to g(N) according the RG equation (16). Moreover, if N^0 and N are related by one RG cycle, $N^0 = e N$, then $g(N^0) = g(N)$. Thus a plot of the spectrum fE (g; ;N)g as a function of N but at xed g, is expected to reveal the cyclicity fE (g; ;e N)g = fE (g; ;N)g. Since our RG procedure is not exact, we expect to observe this signature within the range of our approximations, i.e. for F j !. Indeed, this agrees with the result shown in eq. (12). This can also be observed in g. 1 for the one Cooper pair case, with the cyclicity given by $_1 = 2$ (see below).

E lim inating g_0 in eq. (16) in terms of the mean-eld solution, eqs. (10,11), we observe that the jumps in g(s) from +1 to 1 occur at scales $s = t_h \cdot A s N$ decreases,

g increases steadily to +1 and then jumps to 1. At g = +1, $t_0 = 0$, $t_1 = =$;::; whereas for g = 1, $t_0 = =$, $t_1 = 2 =$;::: Plugging this into eq. (10), one readily sees that

$$_{0}(g = +1) = 1$$

 $_{n+1}(g = +1) = _{n}(g = 1);$ (18)

which indicates that at every jump the lowest condensate disappears from the spectrum , since E $_{\rm C}^{(0)}$ (g = +1) =

1. Eq. (18) in plies, for the remaining condensates, $E_{c}^{(n+1)}(g = +1) = E_{c}^{(n)}(g = -1)$. This result is in agreement with eq. (12). Therefore, the condensate $j_{BCS}^{(n+1)}$ i of one RG cycle plays the same role as $j_{BCS}^{(n)}$ i of the next cycle.

The blow up of $_0$ and $E_c^{(0)}$ at g = +1 is an artifact of the RG scheme used here, since we can only trust the RG for energies below the cuto !. However the disappearance of bound states is correctly described by this RG (see the one Cooper pair problem for a more detailed discussion).

W hen N = 1 the in nite number of condensates are allexpected to appear in the spectrum . However at nite N this is not possible since the Hilbert space is nite dimensional. One can use the RG to estimate the number of condensates $n_{\rm C}$ in the spectrum as a function of N. From the discussion above a condensate disappears from the spectrum for each RG cycle. Thus $n_{\rm C}$ should simply correspond to the number of cycles in log N :

$$n_{\rm C}$$
 – log N : (19)

So far we have found a close relationship between the spectrum of our extended BCS ham iltonian in the meaneld approximation and the RG ow of the coupling constants. In order to get a further con mation of our results we should compute the spectrum for a nite size system. However, it is very di cult to reach intermediate sizes for this model numerically, since the dimension of the Hilbert space grows as $2^{N} = N^{1=2}$. Fortunately, to this end, the similarities between the many-body case and the case of one C opper pair, in the presence of the Ferm i sea, are widely known.

For one C ooper pair in the presence of the Ferm i sea, consider an eigenstate of the form $j = \int_{j}^{j} b_{j}^{y} D i$. The Schrödinger equation reads

$$("_{j} E)_{j} = G_{j} + (G + i)_{l < j} X X_{l} + (G i)_{l < j} X_{l}$$
(20)

with "j 2 (0;!), i.e. the Ferm i sea is not accessible for the pair. Then, in the large N limit, the sum s $_{\rm j}$ are replaced by integrals $_0^{\rm C}$ d"=2 , leading to

$$(\mathbf{m} \quad \mathbf{E} \) \quad (\mathbf{m}) = \mathbf{g}_{0} \quad \frac{\mathbf{d}^{\mathbf{m}0}}{2} \quad (\mathbf{m}^{0}) + \mathbf{i}_{0} \quad \mathbf{m}^{-1} \quad \frac{\mathbf{d}^{\mathbf{m}0}}{2} \quad (\mathbf{m}^{0}):$$
(21)

D i erentiating the above with respect to " and integrating, one obtains

(")
$$\frac{1}{E} e^{i \log (E)}$$
: (22)

This wave function does not have cuts in two cases, i) E < 0 and ii) E > ! . Case i) corresponds to the usual C opperpairproblem where one is looking for bound state solutions (these are the solutions we claim to have a sim - ilar behavior to the m any-body case). P lugging eq. (22) back into eq. (21) one nds

$$\frac{g+i}{g-i} = 1 \frac{!}{E}^{i}$$
: (23)

This equation has an in nite number of solutions given by

$$E_n = \frac{!}{e^{t_n}}; \quad t_n = t_0 + \frac{2 n}{}; \quad n \ 2 \ Z; \quad (24)$$

where t_0 is the principal solution to the equation

$$\tan \frac{1}{2} t_0 = -; \quad 0 < t_0 < -: \quad (25)$$

The n 0 solutions correspond to $E_n < 0$, while those with n < 0 yield $E_n > !$.

As for the many-body case the spectrum has a scaling behavior for weak coupling systems, namely

$$E_n N e^{t_0 2n} = :$$
 (26)

An RG analysis similar to the one in the many-body problem leads to the equation

$$g_{N_{2}} = g_{N} + \frac{g_{N}^{2} + \frac{2}{N}}{N \ 1 \ g}; \qquad N_{1} = N;$$
 (27)

which in the large N lim it becomes

$$\frac{dg}{ds} = \frac{1}{2}(g^2 + {}^2):$$
(28)

The solution to this equation is given by eq. (16) just by replacing s ! s=2. This implies that the period of the cyclicity in $s = \log N_0 = N$ is $_1 = 2 = .$

The factor 1=2 in the above form ulas, as compared to them any-body problem, com es from the non accessibility of the C ooper pair to the states below the Ferm i level.

The discussion leading to eq. (18) can be repeated for the one-C ooper pair problem where the role of the n-th condensate is played by the n-th bound state with energy E_n given in eq. (24), obtaining

$$E_0 (g = +1) = 1$$

 $E_{n+1} (g = +1) = E_n (g = 1)$: (29)

Thus we expect that in each RG cycle a bound state will disappear. The analogue of equation (19) is

$$n_{\rm B} = \frac{1}{2} \log \frac{N}{2}; \qquad (30)$$

where $n_{\scriptscriptstyle\rm B}\,$ is the number of bound states in the spectrum .

This again shows the agreement between the meaneld and the RG results. We con m bellow this picture with numerical calculations.




FIG.1: Exact eigenstates of one-Cooper pair H am iltonian for N levels, from N₀ = 500 down to 50. We depict only the states nearest to zero. The vertical lines are at the values N_n = e^{n 1}N₀. The dotted horizontal lines show the cyclicity of the spectrum .

Fig.1 shows the num erical solution of eq. (20) for g = 1, = 12 and N ranging from 500 down to 50. For each N there are n_B (N) bound states $E_n < 0$, where n_B (N) is in good agreement with eq. (30).

The spectrum shows the self-similarity found in the approaches above: scaling the system by a factor e^{-1} , with $_1 = 2 =$, one recovers the same spectrum for sufciently small energies, i.e.

$$E_{n+1} (N;g;) = E_n (e^{-1}N;g;):$$
 (31)

Fig. 1 also shows the existence of critical values N $_{\rm c;n}$, in the intervals (e ^{n-1}N ; e $^{(n+1)-1}N$), where the bound state closest to the Ferm i level disappears into the \continuum". This e ect leads to the reshu ing of bound states, n + 1 ! n, observed in eq. (31). The critical sizes are also related by scaling, i.e. N $_{\rm c;n}$ =N $_{\rm c;n+1}$ = e $^{-1}$. A llthese phenom ena are in good agreem ent with the RG interpretation we proposed, where the condensates disappear at scales where g (s = t_n) = +1. A llthese points are related by the scaling factor e $^{-1}$.

The RG behavior is presented in g.2, which shows the eigenvalues $E_n (N)$ of the one-C coper pair H am iltonian, with g_N running under eq. (27). The spectrum remains unchanged for $E_n (N) = N$, as shown in g.2b. In g.2a one observes that for the energies $E_n (N) > N$ the result of the RG is not reliable. Nevertheless the RG ow describes qualitatively the disappearance of the lowest bound state and the reshu ing of energy levels after a cycle, and furtherm ore at the predicted scales.

In sum m ary, we have shown that adding to the standard BCS Ham iltonian a time reversal breaking term, parametrized by a coupling constant , generates an in – nite num ber of condensates with energy gaps $_{\rm n}$ related, for weak BCS couplings g, by a scale factor e with = = . This unusual spectrum is explained by the cyclic behavior of the RG ow of g, which reproduces itself after a nite RG time s equal to . We have also solved the nite temperature BCS gap equation, obtaining a critical temperature T_{c,n} for the n-th condensate which is related to the zero temperature gap $_{\rm n}$ (0) exactly as in the BCS theory, i.e. $_{\rm n}$ (0)=T_{c,n} = 3.52 for weak couplings.

The sim plicity of the model proposed in this letter suggests that Russian doll superconductors could perhaps be realized experim entally. Finally, we point out that the critical tem perature can be raised by varying .




FIG.2: Eigenstates of one-C ooper pair H am iltonian with q_N given by eq.(27) with $q_0 = 1$ and = 12. The vertical lines denote the positions at which g jumps from +1 to 1.

W e thank D.Bernard, J.Dukelsky and M A.Mart n-Delgado for discussions. This work has been supported by the Spanish grants SAB2001-0011 (AL), BFM 2000-1320-C 02-01 (JM R and G S), and by the NSF of the USA.

- [y] On leave from Newman Laboratory, Cornell University, Ithaca, NY 14853.
- [1] P.F.Bedaque, H.-W. Hammer, and U.van Kolck, Phys. Rev.Lett. 82 (1999) 463.
- [2] S.D.G lazek and K.G.W ilson, hep-th/0203088.
- [3] D.Bemard and A.LeClair, Phys.Lett.B 512 (2001) 78.
- [4] J. von Delft, D.C. Ralph, Phys. Reps., 345, 61 (2001).
- [5] J. M. Roman, G. Sierra, and J. Dukelsky, Nucl. Phys. B 634 (2002) 483.
- [6] A P. Balachandran, E. Ercolessi, G. Morandi, A M. Srivastava, Lect. Notes in Phys. 38 (World Scientic, 1990).