M easuring the transm ission phase of a quantum dot in a closed interferom eter

Am non Aharony^a, O ra Entin-W ohlm an^{a,b}, and Yoseph Im ry^c

^a School of Physics and Astronom y, Raym ond and Beverly Sackler faculty of Exact Sciences,

TelAviv University, TelAviv 69978, Israel

^b A bert E instein M inerva Center for Theoretical Physics, W eizm ann Institute of Science, Rehovot 76100, Israel

^c Department of Condensed Matter Physics, Weizmann Institute of Science, Rehovot 76100, Israel.

(April 14, 2024)

The electron transm ission through a closed A haronov-B ohm m esoscopic solid-state interferom eter, with a quantum dot (Q D) on one of the paths, is calculated exactly for a sim plem odel. A lihough the conductance is an even function of the magnetic ux (due to 0 nsager's relations), in many cases one can use the measured conductance to extract both the am plitude and the phase of the \intrinsic" transm ission am plitude $t_D = if_D je^{i D}$ through the \bare" Q D. We also propose to compare this indirect measurement with the (hitherto untested) direct relation $sin^2 (D) = j_D f = max (f_D f)$.

PACS numbers: 73.63.-b, 03.75.-b, 85.35Ds

Recent advances in nanoscience raised interest in quantum dots (QDs), which represent articial atoms with experimentally controllable properties [1,2]. Connecting the QD via metallic leads to electron reservoirs yields resonant transmission through the QD, with peaks whenever the Fermi energy in the leads crosses a resonance on the QD. The energies of the latter are varied by controlling the plunger gate voltage on the QD, V. The quantum information on the tunneling of an electron is contained in the complex transmission amplitude, $t_D =$

i $\overline{T_D}e^{i_D}$. The phase $_D$ is particularly interesting, given its relation to the additional electron occupation in the system via the Friedel sum rule [3,4]. This phase is also predicted to exhibit interesting behavior e.g. near a K ondo-like resonance [5]. This motivated experimental attempts to measure $_D$ [6,7], using the A haronov-B ohm interferom eter (ABI) [8].

In the ABI, the QD is placed on one branch, in parallel to a \reference" branch (both connecting the two external leads). A magnetic ux in the area between the two branches creates a phase di erence = e = hcbetween the wave functions in the two branches [9]. In the two-slit lim it, the total ABI transm ission is

 $T = f_{D}^{2} = f_{D} e^{i} + t_{B} f = A + B \cos(+);$ (1)

with = $_{\rm D}$, where contains V -independent contributions from the reference transmission, $t_{\rm B}$ =

it j_{e} $j_{e}^{i_{B}}$, and from the electron \optical" paths on the two branches. However, for the \closed" two-term inalgeom etry, unitarity (conservation of current) and time reversal symmetry in ply the Onsager relations [10], which state that the two-term inal conductance, $G = (e^{2}=h)T$, is an even function of . Therefore, a naive t of the experimental transmission to Eq. (1) must yield = 0 or { with no relation to _D. Indeed, the experimental data [6] for T depend only on cos [11].

A im ing to measure a non-trivial A B phase shift then led to experiments with \open" interferometers [7,12],

which contain additional \leaky" channels, breaking the O nsager symmetry. A t to Eq. (1) then yields a phase which increases (with V) gradually from 0 to through

each resonance. However, the detailed V -dependence of depends on the strength of the coupling to the additional term inals [13]. A though it is possible to optim ize this strength, and reproduce the two-slit conditions [14], this involves large uncertainties.

In the present paper we present exact results for the total transm ission of the closed A B I, T . A lthough T is even in , contradicting the sim ple two-slit Eq. (1), it does depend on both T_D and $_D$. Under appropriate conditions (see below), one can thus extract $_D$ from the measured T, eliminating the need to open the interferom eter. This possible extraction was not noticed in earlier discussions of the closed A B I. Theoretical analyses used the K eldysh technique, combined with the wide-band and related approximations [15,16], or ignored electron-electron interactions [17]. These approximations, which sometimes m iss important features of the results (see below), are avoided in our calculation, which is done in the linear response limit, and at temperature T = 0.

W e dem onstrate our results for a sim ple lattice m odel, shown in Fig. 1: for = 0, each (unit length) segment in the gure represents a real tight-binding hopping matrix element J; **⊥;** J and J, as in-¥; dicated. All the on-site energies are zero, except $_{\rm D}\,$ on the site \det and $_0$ on the site ref (which sits on the reference path and represents a simple point contact, a tunnel junction, etc). The latter two energies can be varied experim entally by the plunger (or point contact) gate voltages D V and $_0$ V_0 [11]. A susual for such models, electron-electron interactions are included only via an on-site Hubbard interaction U on the QD. The AB phase in the triangle, = $_{\rm L}$ + $_{\rm R}$, is included by attaching a factor $e^{i_{L}}$ ($e^{i_{R}}$) to the hopping matrix element J_L (J_R). At T = 0, the electron energy $_k$ = 2J cosk is equal to the Ferm i energy on the leads, $_{\rm F}$, and we calculate the transmission for electrons with spin $\$.

We start by reviewing the \intrinsic" transmission through the QD, without the reference path (e.g. for large $y_0 j = j_0 j$ or with $I_L = I_R = 0$). A dapting the results of R ef. [18], one has

$$t_{D} = i_{D} \sin_{D} e^{i_{D}} \quad 2i \sin \frac{1}{2} J_{R} g_{D} (k) = J; \quad (2)$$

with the QD asymmetry factor $_{\rm D} = 2J_{\rm L} J_{\rm R} = (J_{\rm L}^2 + J_{\rm R}^2)$ and the \intrinsic" G reen function on the QD, $g_{\rm D}$ ($_{\rm k}$) = $l=[_{\rm k} _{\rm D} _{\rm D} (_{\rm k})]$. Here, $_{\rm D}$ ($_{\rm k}$) is the self-energy on the QD, which contains contributions from the leads, $_{\rm D,ext} = e^{j\mathcal{K}j}(J_{\rm L}^2 + J_{\rm R}^2)=J$ (which exists also for the non-interacting case [13]), and from the electron-electron interactions on the QD itself, $_{\rm D,int}$ (!) (which vanishes when U = 0). As $_{\rm D}$ V increases, $_{\rm D}$ grows gradually from zero (far below the resonance), through =2 (at the resonance), towards (far above the resonance).

Interestingly, for this one-dimensional model, norm alizing the measured $T_D = \frac{1}{D_D} \int_D^2 = \frac{2}{D} \sin^2(D_D)$ by its (V-independent) maximum $\frac{2}{D}$ yields the value of D_D . A sum ing coherence, this (hitherto ignored) method for measuring D_D directly from T_D eliminates the need for any complicated interferom etry! [19] In the remainder of this paper we discuss ways of extracting D_D indirectly, from the closed ABI measurements. Comparing results from $\sin^2(D_D) = T_D = \frac{2}{D}$, from the closed ABI (below) and from the open ABI [14] (all with the same QD) should serve as consistency checks for this conclusion.

The same analysis yields the transm ission amplitude through the reference path (when e.g. $J_L = J_R = 0$),

$$t_{B} = i_{B} \sin_{B} e^{i_{B}} 2i \sin \frac{1}{2} I_{R} g_{B} (k) = J \quad (3)$$

with the bare reference site G reen function $g_B = 1 = [_k _0 + e^{i j k \cdot j} (I_L^2 + I_R^2) = J]$, and $_B = 2 I_L I_R = (I_L^2 + I_R^2)$. In the two-slit situation, Eqs. (2) and (3) su ce to determ ine the overall transm ission, as in Eq. (1). However, the situation is more complicated for the closed ABI. The main result of the present paper concerns the exact transm ission amplitude through the closed ABI,

$$t = A_D t_D e^i + A_B t_B; \qquad (4)$$

where we nd $A_D = g_B \begin{pmatrix} k & 0 \end{pmatrix} G_D \begin{pmatrix} k \end{pmatrix} = g_D \begin{pmatrix} k \end{pmatrix}$ and $A_B = 1 + G_D \begin{pmatrix} k \end{pmatrix}_{ext} \begin{pmatrix} k \end{pmatrix}$. Here, $G_D \begin{pmatrix} l \end{pmatrix} = 1 = \begin{bmatrix} l & D \end{pmatrix}$ (!)] is the fully \dressed" Green function on the QD, with the dressed self-energy = $_{int} + _{ext}$. Both terms here di er from their counterparts in the \intrinsic" $_D$, by contributions due to the reference path. Equation (4) looks like the two-slit form ula, $t = t_D e^i + t_B$. How ever, each of the terms is now renorm alized: A_D contains all the additional processes in which the electron \visits" the reference site ($A_D = 1$ when $I_L = I_R = 0$), and A_B contains the corrections to t_B due to \visits" on the dot. In fact, a physical derivation of Eq. (4) am ounts to starting from Eq. (2), and adding an in nite power series in I_L and I_R . We now discuss the -dependence of T f_2^2 , in connection with the Onsager relations and with the possible indirect extraction of $_D$.

We rst note that both parts in ($_{\rm k}$) are even in , due to additive contributions (with equal amplitudes) from clockwise and counterclockwise motions of the electron around the ring (see e.g. Refs. [8,13,17,20]). In order that T also depends only on cos , as required by the Onsager relations, the ratio K $A_{\rm B} \, t_{\rm B} = (A_{\rm D} \, t_{\rm D}) \times [G_{\rm D} \, (_{\rm k})^{-1} + _{\rm ext} \, (_{\rm k})]$, with the real coe cient $\varkappa = I_{\rm L} \, I_{\rm R} = [J_{\rm L} \, J_{\rm R} \, (_{\rm k} \, _{\rm O})]$, must be real, ie.

$$= [G_{D}(k)^{1} + ext(k)] = int 0:$$
(5)

The same relation follows from the unitarity of the 2 2 scattering m atrix of the ring. This relation already appeared for the special case of single in purity scattering, in connection with the Friedel sum rule [4], and was implicitly contained in Eq. (2), where $= D_{jint} = 0$ [18]. Equation (5) implies that (at T = 0 and ! = $_k$) the interaction self-energy $_{int}(k)$ is real, and therefore the width of the resonance, $= G_D(k)^{-1}$, is fully determined by the non-interacting self-energy = ext(k).

Since $_{ext}(!)$ depends only on the (non-interacting) tight-binding terms, it is easy to calculate it explicitly. We nd $_{ext}(_k) = _{D;ext}(_k) + _{ext}$, where

$$ext = e^{2ijkj}g_B (J_L^2 I_L^2 + J_R^2 I_R^2 + 2J_L J_R I_L I_R \cos) = J^2: (6)$$

The term proportional to \cos comes from the electron clock-and counterclockwise motion around the ABI \ring". Similarly, one can write $int(k) = D_{int}(k) + D_{int}(k)$

 $\begin{array}{ll} & \text{int, and thus } G_D\left(\begin{smallmatrix}k\\ k\end{smallmatrix}\right)^1 = g_D\left(\begin{smallmatrix}k\\ k\end{smallmatrix}\right)^1 , \text{ with } = \\ & \text{ext} + & \text{int. Hence, } t = A_D t_D (e^i + K). \text{ W riting also} \\ & A_D = C = [1 \quad g_D\left(\begin{smallmatrix}k\\ k\end{smallmatrix}\right)], \text{ with } C = (\begin{smallmatrix}k\\ k\end{smallmatrix})_0 g_B, \text{ we nd} \end{array}$

$$T = jC j^{2} T_{D} \frac{1 + K^{2} + 2K \cos}{1 - 2 \langle g_{D} \rangle + jg_{D} \rangle j^{2}};$$
 (7)

Equation (7) presents an alternative form of our main result. A though the num erator looks like the two-slit (depending on signK), the Eq. (1), with = 0 or new physics is contained in the denom inator { which becom es im portant in the vicinity of a resonance. The central term in this denominator depends explicitly on the phase of the complex number $g_{\!\scriptscriptstyle D}$. Since this number is directly related to t_D , via Eq. (2), one may expect to extract $_{D}$ from a t to Eq. (7), taking advantage of the dependence of the denom inator on cos . Physically, this dependence originates from the in nite sum over electron paths which circulate the ABI ring. The rest of this paper is devoted to the conditions for such an extraction. Generally, this is not trivial, as one needs the detailed dependence of on cos and on the various param eters. We have presented this dependence for ext, but not for int•

The extraction of $_{\rm D}$ becomes easy when one may neglect $_{\rm int}$. The simplest case for this is for singleelectron scattering, when $_{\rm int} = 0$. Interactions (i.e. U) are also negligible for a relatively open dot, with small barriers at its contacts with the leads [21]. Another e ectively single-electron scattering case arises near a C oulom b blockade resonance, when the e ect of interactions can simply be absorbed into a Hartree-like shift, $_{\rm D}$ + $_{\rm int}$! $_{\rm D}$ + N U, if one assumes that N depends smoothly on the number of electrons on the Q D, and not on [20]. If one may neglect $_{\rm int}$, then $_{\rm ext}$ is given in Eq. (6). Using also Eqs. (2) and (3), we nd

$$T = j C j^{2} T_{D} \frac{1 + K^{2} + 2K \cos}{1 + 2P (z + \cos) + Q (z + \cos)^{2}}; \quad (8)$$

where $z=~(J_L^2\,I_L^2+J_R^2\,I_R^2)=(2J_L\,J_R\,I_L\,I_R\,),\,P=<[vt_B\,t_D\,],\,Q=~jvt_B~j^2T_D$, and $v=~e^{2ijk\,j}=(2\sin^2\,k\,j)$ depends only on the Ferm i wavevector k, independent of any detail of the ABI.A 5-parameter t to the explicit –dependence in Eq. (8) for given values of V and V_0 then yields $j C~j^2T_D$; K; z;P and Q, and thus $\cos(~_D + ~_B + 2jk\,j = P = Q$, from which one can extract the V-dependence of $_D$. The same V-dependence of $_D$ is also contained in K / (cot $_D$ + cot $jk\,j$). As discussed after Eq. (2), ourm odel also in plies that $T_D=~_D^2\,\sin^2(~_D)$. Since the V-dependence of T_D can also be extracted from the t-ted values of either $jC~j^2T_D$ or Q, we end up with several consistency checks for the determ ination of $_D$. Additional checks arise from direct measurements of T_D and $T_B=~j_B~j^2$, by taking the lim its $jV_0\,j=~j_0\,j\,!$ 1 or $jV\,j=~j_D\,j\,!$ 1.

The LHS frame in Fig. 2 shows an example of the Vand -dependence of T for this limit (no interactions), with k = -2 and $J_L = J_R = I_L = I_R = 1$; $V_0 = 4$ (in units of J), im plying $K = D = 0 = V = V_0$. Far away from the resonance T 1,Q Pj 1 and Kj 1, yielding the two-slit-like form T A + B cos , dom inated by its rst harmonic, with B = A2 K ¹ P]. However, close the the resonance T shows a rich structure; the denom inator in Eq. (8) generates higher harm onics, and the two-slit form ula is completely wrong. This rich structure m ay be m issed if one neglects parts of the dependence of , as done in Ref. [16]. Note also the Fano vanishing [22] of T for V 10 at = 2n, with integer n.W ithout interactions, everything can be extended to a QD with many resonances, e.g. due to Coulom b blockade shifts in the electric d with the number of electrons. U sing a generalization to Eq. (8), given in Ref. [14], the RHS frame in Fig. 2 shows results for two resonances, with $_{\rm D}$ = 5. Interestingly, Fig. 2 is qualitatively sim ilar to the experim entally measured transmission in Ref. [11]. How ever, so far there has been no quantitative analysis of the experim ental data.

To treat the general case, we need inform ation on $_{int}$. First of all, we emphasize that a successful t to Eq. (8) justi as the neglect of the -dependence of $_{int}$. If the various procedures to determ ine $_D$ from Eq. (8) yield the same V-dependence, this would also con m that $_{int}$ is negligibly small. A failure of this check, or a more complicated dependence of the measured T on cos , would im ply that $_{int}$ is not negligible.

As seen from Eq. (6), $_{ext}$ is fully determined by a single \visit" of the electron at \ref". For small T_B , or large $j_{V_0} j = j_0 j$, it is reasonable to conjecture that $_{int}$ is also dominated by such processes. In that case, we expect $_{int}$ to be proportional to the same brackets as in Eq. (6), i.e. $_{int}$ w (z + cos), with a real coe cient w. This yields the same dependence of T on cos as in Eq. (8), with a shifted coe cient v. If w depends only weakly on V, then this shift has little e ect on the determination of $_D$. Again, the validity of this approach relies on getting the same V-dependence of $_D$ from all of its di erent determinations.

The situation becomes more complicated near a Kondo-like resonance. Maintaining the (non-trivial) assumption that $G_D = 1 = [!]$ D $_{\rm D}$ (!)], the K ondo peak at the Ferm i energy must be generated by $D_{\rm D}$. For the intrinsic QD, this yields $_{D}$ = =2 and t_{D} = $_{D}$, resulting in a V-independent plateau for T_D . A priori, it is not obvious what happens in the presence of the \reference" path. Hofstetter et al. identi ed the Kondo region by requiring that the phase res of the fully dressed Green function G_D be equal to =2.0 ur result for G_D shows that this is impossible: the phase res depends on , via the -dependence of , and thus cannot be set at the constant value =2. (Apparently, this -dependence was neglected in the analytic parts, and weak for the numerical parameters used in Ref. [16]). Alternatively, one m ight assume that the \bare" QD sticks to the K ondo resonance, and thus $_{D} = = 2$ (independent of V) even in the ABI. Equation (7) then replaces the Kondo plateau by a complicated dependence on (including the rst harm onic), which diers signicantly from that of Ref. [16]. C learly, this lim it requires m ore research.

Finally, we give some m ore details of our derivation. O ur H am iltonian, which simply adds the reference path to that of N g and Lee [18], is

$$H = {}_{D} {}^{X} {}_{d} {}^{y} {}_{d} + \frac{U}{2} {}^{X} {}_{n_{d}} {}_{n_{d}} {}_{-}$$

$$+ {}^{X} {}_{k} {}^{y} {}_{c} {}_{k} + {}^{Y} {}_{k} {}_{d} {}^{y} {}_{c_{k}} + {}^{V} {}_{k} {}^{d} {}^{y} {}_{c_{k}} + {}^{V} {}_{k} {}^{c} {}^{y} {}_{c_{k}} {}^{d} {}_{-}$$

$$+ {}^{k} {}_{0} {}^{x} {}_{0} {}^{y} {}_{c_{0}} {}^{z} {}_{+} {}^{k} {}^{Y} {}_{c_{0}} {}^{z} {}_{c_{0}} {}^{z} {}_{+} {}^{k} {}^{y} {}_{c_{0}} {}^{z} {}_{c_{0}} {}^{z} {}_{+} {}^{z} {}^{y} {}_{c_{0}} {}^{z} {}_{+} {}^{z} {}^{y} {}_{c_{0}} {}^{z} {}_{+} {}^{z} {}^{z} {}^{z} {}^{z} {}_{c_{0}} {}^{z} {}_{+} {}^{z} {}^{z}$$

where c_k^y creates single particle eigenstates (with spin) on the unperturbed \background" chain (with $I_L = I_R = J$, $J_L = J_R = 0$, with eigenenergy $_k = 2J \cos k$, while $c_p = {}_k c_k = \overline{N}$, $U_k = [(I_L J)e^{ik} + (I_R J)e^{ik}] = \overline{N}$, and $V_k = (I_L e^{i + ik} + J_R e^{i + r^{+}ik}) = \overline{N}$. The operators on the dot, d and d^y , anti-com m ute with c_k ; c_k^y . A lso, n_d = d^yd , and $\bar{}$.

As stated above, one can derive Eq. (4) by expanding Eq. (2) in powers of I_L and I_R . A more general approach uses the standard relation between the 2 2 scattering m atrix T_{kk^0} and the m atrix of retarded singleparticle G reen functions, G_{kk^0} (!) = $_{kk^0}g_k^0 + g_k^0T_{kk^0}g_{k^0}^0$, with g_k^0 (!) = 1=(! k), evaluated on the energy shell, $! = F = k = k^{\circ}$ [5]. The equation-of-motion (EOM) method is then used to express (! $_{k}$)G $_{kk^{0}}$ (!) and k)Gkd (!) as linear combinations of each other and (! of G_D (!), allowing us to express each of them (and thus also t / $T_{kj;kj}$ in terms of G_D (!), yielding Eq. (4). Since we do not use an explicit solution for G_D (!) itself, we don't need to deal with the higher order correlation functions (due to U), which appear in its EOM.

We hope that our paper will stimulate attempts to t experimental data to our Eq. (8), and to compare the resulting $_{\rm D}$ with its direct estimate via T. This procedure should work in many cases. We also hope that our paper will stimulate more detailed theoretical calculations of $_{\rm int}$. As explained, the existing approximate calculations miss the crucial -dependence of these interactiondependent term s.

W e thank R.Englm an, M.Heiblum, Y.Levinson, A. Schiller, H.A.W eidenmuller and A.Yacoby for helpful conversations. This project was carried out in a center of excellence supported by the Israel Science Foundation, with additional support from the Germ an-Israeli Foundation (GIF), and from the Germ an Federal M inistry of Education and Research (BMBF) within the Fram ework of the Germ an-Israeli Project Cooperation (DIP).

- [1] L.P.K ouw enhoven et al., M esoscopic E lectron Transport, NATO A dvanced Study Institute, Series E : A pplied Science, Vol. 345, edited by L.L.Sohn, L.P.K ouvwenhoven and G.Schon (K luwer, D ordrecht, 1997), p.105.
- [2] Y. Im ry, Introduction to Mesoscopic Physics (Oxford University Press, Oxford 1997; 2nd edition, 2002).
- [3] J.Friedel, Can. J. Phys. 34, 1190 (1956).
- [4] D.C.Langreth, Phys.Rev. 150, 516 (1966).
- [5] A.C.Hewson, The Kondo Problem for Heavy Fermions (Cambridge University Press, Cambridge 1997).
- [6] A. Yacoby, M. Heiblum, D. Mahalu and H. Shtrikman, Phys. Rev. Lett. 74, 4047 (1995).
- [7] R.Schuster et al, Nature 385, 417 (1997).
- [8] Y.Gefen, Y.Im ry and M.Ya.Azbel, Phys. Rev. Lett. 52, 129 (1984).
- [9] Y.Aharonov and D.Bohm, Phys. Rev. 115, 485 (1959).
- [10] L. Onsager, Phys. Rev. 38, 2265 (1931); H. B. G. Casim ir, Rev. M od. Phys. 17, 343 (1945); M. Buttiker, Phys. Rev. Lett. 57, 1761 (1986).

- [11] See also K .K obayashi, H .A ikawa, S.K atsum oto, and Y . Iye, Phys. Rev. Lett. 88, 256806 (2002).
- [12] Y. Ji, M. Heiblum, D. Sprinzak, D. Mahalu, and H. Shtrikman, Science 290, 779 (2000).
- [13] O. Entin-Wohlman, A. Aharony, Y. Imry, Y. Levinson and A. Schiller, Phys. Rev. Lett. 88, 166801 (2002).
- [14] A.Aharony, O.Entin-Wohlman, B.I.Halperin and Y. Im ry, Phys. Rev. B 66, 115311 (2002).
- [15] B.R.Bulka and P.Stefanski, Phys. Rev. Lett. 86, 5128 (2001).
- [16] W . Hofstetter, J. Konig and H. Schoeller, Phys. Rev. Lett. 87, 156803 (2001).
- [17] O. Entin-W ohlman, C. Hartzstein and Y. Im ry, Phys. Rev.B34, 921 (1986); Z.Y. Zeng, F.C laro and A. Perez, Phys. Rev.B65, 085308 (2002); T.S.K im et al., Phys. Rev.B65, 245307 (2002); B.K ubala and J.K onig, Phys. Rev.B65, 245301 (2002); Ref. [13].
- [18] T.K.Ng and P.A.Lee, Phys.Rev.Lett.61, 1768 (1988).
- [19] However, interferrom etry is still important, since it ensures coherence. Interestingly, this conclusion holds for any Breit-W igner-like resonance, with an energyindependent width. Therefore, it may hold beyond our speci c m odel.
- [20] H.A.W eidenmuller, Phys. Rev. B65, 245322 (2002).
- [21] K.A.Matveev, PhysRev.B51, 1743 (1995).
- [22] U. Fano, Phys. Rev. 124, 1866 (1961); O. Entin-Wohlman, A. Aharony, Y. Im ry, and Y. Levinson, J. Low Tem p. Phys. 126, 1251 (2002) and references there.




FIG.1. M odel for the closed ABI.




FIG.2. AB transmission T versus the AB phase $\$ and the gate voltage V, for one (LHS) and two (RHS) non-interacting resonances.