

Relation between d-density wave of electron and staggered flux of spinon

Koichi Hamada^{a,1}, Daijiro Yoshioka^a

^aDepartment of Basic Science, University of Tokyo, Komaba 3-8-1, Meguro-ku, Tokyo 153-8902, Japan

Abstract

A $d_{x^2-y^2}$ -density wave (ddw) order of electron in two-dimensional t-J model is analyzed in saddle point level using the U(1) slave boson formalism. We considered not only the staggered flux (s-flux) order of spinon but also the s-flux order of holon. This analysis provides the relation between the s-flux order of spinon and the ddw order of electron. We discovered a new phase in the phase diagram. In this phase, there is a s-flux order of spinon, but no ddw order of electron. Our results are that 1) a region of electron ddw exists, 2) there is no coexistence of ddw and $d_{x^2-y^2}$ -wave pairing (singlet-RVB) in all region of phase diagram, and that 3) the ground state is a purely $d_{x^2-y^2}$ -wave superconducting state.

Key words: Superconductivity; Cuprate; Pseudo gap; d-density wave; t-J model

Recently, Chakravarty et al. [1] proposed that the electron $d_{x^2-y^2}$ -density wave (ddw) order exists in the pseudo gap region of high- T_c superconductor. The electron ddw state [2] is the staggered flux (s-flux) state [3,4] of electron coordinate. In this state, $d_{x^2-y^2}$ -wave (d-wave) gap exists, time-reversal-symmetry is broken and the 'real' staggered current of the electron exists. The order parameter of the ddw is $y_e = \frac{1}{2} \sum_k (\cos k_x - \cos k_y) h c_k^y c_{k+Q}^\dagger$, $Q = (\pi; \pi)$. However they didn't discuss its scenario microscopically. There remains a question, can the electron ddw phase exist in highly correlated system?.

The 2-dimensional t-J model is a promising model which includes highly correlated effects. Many phases are proposed in this model [3-9]. Zhang [8] analyzed the competition between s-flux order and $d_{x^2-y^2}$ -wave pairing order at zero temperature by Gutzwiller approximation. The s-flux state is unstable against in nite s-d-wave pairing at nite doping. Ubbens and Lee [9] analyzed this model at nite temperature. The s-flux phase of the spinon exists on the region where doping and temperature are both nite.

The SU(2) s-flux state, in which the Fermi-surface of spinon are always points, is also analyzed in SU(2) slave boson model [10,11]. However, there remains a question, how do electrons behave in nite temperature s-flux phase? The current of the spinon and the electron are not equivalent in nite temperature s-flux phase because there is no Bose condensation of holon. The region of nite temperature s-flux phase exists above the temperature of holon condensation.

In this paper, we revealed the relation between the ddw of electron and the s-flux of spinon. We analyzed it in the 2-dimensional t-J model based on the U(1) slave boson formalism. We introduce order parameters $i_j = h \frac{f_i^y}{f_j} f_j^\dagger$, $i_j = h f_i^\dagger f_{j\#} - f_{i\#} f_j^\dagger$, $B_{ij} = h b_i^\dagger b_j$ to decouple the Hamiltonian.

We considered not only the staggered flux order of the spinon but also the staggered flux order of the holon, $i_{\pm x;i} = x_s + i(1)^\dagger y_s$, $i_{\pm y;i} = x_s - i(1)^\dagger y_s$, $B_{i\pm x;i} = x_h + i(1)^\dagger y_h$, $B_{i\pm y;i} = x_h - i(1)^\dagger y_h$. Here, \hat{x} and \hat{y} are unit vectors in the x and y direction, $x_s = \cos(\phi_s=4)$, $y_s = \sin(\phi_s=4)$, $x_h = B \cos(\phi_h=4)$, $y_h = B \sin(\phi_h=4)$. The order parameters y_s and y_h correspond to the ddw order parameter of spinon and holon, respectively. For the pairing symmetry, we considered $d_{x^2-y^2}$, namely

¹ E-mail: koichi@sol.a.u-tokyo.ac.jp




Fig.1. Staggered flux of spinon s and holon h .

$$i + \hat{x}; i = i + \hat{y}; i = .$$

This formalism is an extension of the study by Ubbens and Lee [9]. The $f_j^y f_i$ term couples not only to $(3J=8)_{ij}$ but also to tB_{ij} , i.e. the spinon feels the spinon s -flux (s) and the holon s -flux (h). The expectation values of the holon, B and h , have finite value for the solution of self-consistency equations. Two advantages exist in our formalism; 1) this is a new saddle point solution whose free energy is lower than the previous one, 2) this solution provides the relation between the ddw of the electron and the s -flux of the spinon. In this formalism, the hopping order parameter of the electron is a product of the hopping order parameters of spinon and holon.

$$\sum_h c_i^y c_j = \sum_h f_i^y f_j \text{ and } \sum_h b_i = \sum_{ij} B_{ij} \quad (1)$$

The electron s -flux order parameter y_e and the electron ddw order parameter y_e are given by $y_e = s - h$, and $y_e = B \sin(\phi_e/4)$.




Fig.2. MF phase diagram for $t/J=1$. The electron ddw order exists only in region 2. Here, x is hole concentration. The phase diagram for $t/J=2$ is quantitatively similar to the phase diagram for $t/J=1$. With the boson order parameter B_{ij} , the s -flux phase and s -flux phase of spinon and holon extends to the higher-doped region compared to the previous work [9], where B_{ij} were not considered.

We solved the self-consistency equations numerically, and obtained the phase diagram (Fig. 2). At half-filling, the holon order parameter B_{ij} is zero and the degeneracy of spinon between the s -flux state and the d -wave pairing state exists due to the local $SU(2)$ symmetry; $\phi_s \neq 0; B_{ij} = 0; y_s^2 + y_h^2 = \text{const} \neq 0$. In region 1, spinon s -flux exists but electron ddw doesn't exist. The spinon and holon state are s -flux

order state respectively. In electron picture, the s -flux is canceled completely; $\phi_s \neq 0; B \neq 0; s = h = 0; y_e = 0$. The electron ddw order parameter, $y_e = B \sin(\phi_e/4) \neq 0$. The electron ddw order exists only in region 2. The staggered current of electron can be observed experimentally. The spinon s -flux s and holon s -flux h are not equal to zero, and $s \neq h; \phi_s \neq 0; B \neq 0; s \neq 0; h \neq 0; y_e = 0$, and $y_e = B \sin(\phi_s - \phi_h)/4 \neq 0$. In region 3, $d_{x^2-y^2}$ -wave pairing exists; $\phi_s \neq 0$ and $B \neq 0; s = h = 0; \phi_e \neq 0$, and $y_e = 0$. In region 4, there exists only uniform hopping order; $\phi_s \neq 0; B \neq 0; s = h = 0$, and $y_e = 0$. In region 5, all order parameters are zero. Spinon and holon cannot hop; $s = B = s = h = 0$, and $y_e = 0$.

The transition between region 1 and region 2 is a 2nd order transition in our theory. (If one only focuses on spinon degree of freedom, this doesn't look like phase transition [9][11].) There exists an order parameter which characterizes this transition. It is the electron ddw order parameter.

In conclusion we have found a possibility for the electron ddw phase in the t - J model at finite temperature, which does not coexist with the singlet-RVB state.

Acknowledgements

K.H. thanks A.Himeda, T.Koretsune, Y.Yanase, M.Ogata, Y.Ueno, K.Nomura, R.Shindou, S.Ryu, Y.Morita, J.Kishine, and N.Nagaosa for their useful discussions. Numerical computation in this work was partially carried out at the Yukawa Institute Computer Facility.

References

- [1] S.Chakravarty, R.B.Laughlin, D.K.Morr, C.Nayak, Phys. Rev. B 63 (2001) 94503.
- [2] C.Nayak, Phys. Rev. B 62 (2000) 4880.
- [3] I.Aeck, J.B.Marston, Phys. Rev. B 37 (1988) 3774.
- [4] J.B.Marston, I.Aeck, Phys. Rev. B 39 (1989) 11538.
- [5] P.W.Anderson, Science 235 (1987) 1996.
- [6] I.Aeck, Z.Zou, T.Hsu, P.W.Anderson, Phys. Rev. B 38 (1988) 745.
- [7] G.Kotliar, J.Liu, Phys. Rev. B 38 (1988) 5142.
- [8] F.C.Zhang, Phys. Rev. Lett. 64 (1990) 974.
- [9] M.J.Ubbens, P.A.Lee, Phys. Rev. B 46 (1992) 8434.
- [10] X.-G.Wen, P.A.Lee, Phys. Rev. Lett. 76 (1996) 503.
- [11] P.A.Lee, N.Nagaosa, T.K.Ng, X.-G.Wen, Phys. Rev. B 57 (1998) 6003.