A Modied Random Phase Approximation of Polyelectrolyte Solutions

A.V.Emoshkin, M.Olvera de la Cruz Department of Materials Science and Engineering, Northwestern University, Evanston, IL, USA 60208-3108 (March 22, 2024)

We compute the phase diagram of salt-free polyelectrolyte solutions using a modi ed Debye-Huckel Approach. We introduce the chain connectivity via the Random Phase Approximation with two important modi cations. We modify the electrostatic potential at short distances to include a bound on the electrostatic attractions at the distance of closest approach between charges. This modi cation is shown to act as a hard core in the phase diagram of electrolyte solutions. We also introduce a cut-o on the integration of the modes of wave length smaller than the size over which the chains are strongly perturbed by the electrostatic interactions. This cut-o is shown to be essential to predict physical phase diagram in long chain solutions.

I. IN TRODUCTION

D ebye-H uckel (D H) theory derived nearly 80 years ago successfully explains m any therm odynam ic properties of dilute sym m etric electrolyte solutions.¹ R ecent com puter simulations of hard-sphere ionic uids² show that the D H approach for nite size ion solutions provides a rather accurate estim ate of the critical tem perature, although it underestim ates the critical density. A conceptually simple modi cation of the D H approach, based on accounting for the interactions of ionic pairs introduced by B jerrum ³ w ith residual ionic uid, how ever, gives better estim ates for the critical param eters.⁴

Though a large scientic e orthas been made towards understanding the therm odynam ics of sim ple electrolyte solutions, exible charged chains solutions are not understood. The Random Phase Approximation (RPA), a DH approach which includes the structure function of the chains, is widely used to describe polyelectrolytes solutions.^{5;6} RPA, how ever, provides unphysical coexisting curves in salt free polyelectrolyte solutions.⁷ In particular, RPA predicts that the critical density decreases with increasing the chains degrees of polym erization N, and it goes to zero as N goes to in nite. M oreover, the critical tem perature increases rapidly as N increases suggesting that strongly charged chains are not water soluble in salt-free solutions. Recent com puter simulations of charged chains reveala critical density rather insensitive to the value of N, and a critical tem perature that does increases as N increases but not as rapidly as predicted by the RPA approach.⁸ A one component plasm a approach to polyelectrolytes solutions gives both a critical density and a critical tem perature insensitive to the the degree of polym erization N⁹. The one component plasm a approach ignores the contribution from the uctuations of the monomers. That is, the one component plasm a assum es that the chains provide a constant charge density support over which the counterions uctuate. It is di cult to dem onstrate and understand why and when it is possible to neglect the chain density uctuations. M oreover, the value of the critical tem perature is strongly underestin ated in the one component plasm a approach to polyelectrolytes salt-free solutions.

Here we describe a modied polyelectrolyte RPA m odel that predicts realistic values of the critical tem perature, and a critical density rather insensitive to N in agreement with the simulations. The RPA approach described here includes two modi cations. The rstmodi cation is related to the cut-o in RPA. In salt free polyelectrolyte solution the electrostatic contribution to the free energy of charged G aussian chains does not reduces to the DH lim iting law.⁵ Instead, a m ore strong electrostatic contribution results in salt-free solutions that generates instabilities in the free energy as the monom er concentration goes to zero at room tem peratures in strongly charged chains.⁷ Scaling argum ents 10 ^{{12} and computer simulations, 13;14 however, show that charged chains are stretched in monovalent ionic solutions on length scales sm aller and of the order of the inverse screening length. Therefore, it is not possible to include the monomer density uctuations within RPA, which assumes that the electrostatic interactions do not perturbed the chain statistics (a linear response theory). Here we introduce a concentration dependent cut o proportional to the inversive screening length in the RPA electrostatic contribution to include only the contribution from the unperturbed Gaussian chains. W e also discuss other possibilities for the cut-o that yield similar modi cations in the phase diagram . The second and less im portant m odi cation involves including the contribution from the hard core of the ions and the monom ers. In simple electrolyte solutions the hard core of the ions is included in the electrostatic contribution to the free energy to avoid spurious results as the concentration of ions increases. In this paper we modify the interaction potential and show that ourm odi ed potential describes the polyelectrolyte solution at the same level of approximation as the modied DH for hard-sphere ionic uids.

In Section II we describe the modi ed RPA model for salt-free exible strongly charged chains. In Section III we discuss the electrostatic interaction potential used in the model. In Section IV we state the problem of RPA describing Gaussian chains and the required modi cations. In Section V we outline and discuss the results, and give some conclusions.

II.M ODEL AND FREE ENERGY

Let us consider a mixture of negatively charged polyelectrolyte chains and a residual ionic uid. We assume that every chain consists of N m onom ers each carrying a negative charged q and we denote the concentration of m onom ers as . Every ion in the system carries the charge + q and their total concentration is $_+$. D ue to electroneutrality the following condition holds

Wewrite free energy of the system in the form

$$F = F_{ref} + F_{el}$$
 (2)

Here F_{ref} is the free energy of the system without any electrostatic interactions and in the lim it of good solvent it could be written as

$$\frac{F_{ref}}{TV} = + \ln_{+} + \frac{1}{N} \ln_{-} + \frac{1}{a^{3}} (1) \ln_{-} (1) \ln_{-} (3)$$

where T and V is the tem perature (in units of Boltzm ann constant $k_{\rm B}$) and the volume of the system respectively. We also assumed that monomers of the chains and dissociated ions have the same sizes a, so we can de ne the reduced density as follows

$$= a^{3}(++)$$
 (4)

The second term on the right hand side of Eq. (2) is the electrostatic contribution to the free energy. U sing RPA approach we write it in the form 5

$$\frac{F_{el}}{TV} = \frac{1}{4^{2}} \int_{0}^{Z^{k}} \ln (1 + (r_{+} + r_{2}(k))) \frac{U(k)}{T} (r_{+} + r_{+}) \frac{U(k)}{T} k^{2} dk \quad (5)$$

where the structure function of the chain g(k) has the following denition

$$g(k) = \frac{1}{N} \sum_{i;j}^{X D} e^{ik(r_{i} r_{j})}$$
(6)

Indices i and j in Eq. (6) run over all m onom ers of the chain and the average < ::: > is taken over all possible chain con m ations. U (k) in Eq. (5) is the Fourier transform of the interaction potential between charges and the parameter K is used to cut the integration over the large values of k. The purpose of the present work is to discuss the RPA approximation, and determ ine appropriate form s of U (k) and K.

III. IN TERACTIONS BETW EEN CHARGES

For dilute solutions of electrolytes the potential of interaction between charges can be taken in the form

$$U(\mathbf{r}) = \frac{q^2}{\mathbf{r''}}$$
(7)

where r is the distance between charges and " is the dielectric constant of the media. W ith this potential Eq. (5) calculated for a simple case of N = 1 and K = 1 reduces to 3 =12 . That is, the electrostatic contribution reduces to the limiting case (a 1) of the DH expression for the free energy of hard-sphere ionic uid,

$$\frac{F_{el}^{DH}}{TV} = \frac{3}{12}^{DH} (a)$$
(8)

Here the inverse screening length is given by $^2 = 4 q^2 l_B (+ +)$, the B jerrum lenght $l_B = e^2 = "T$ and

$$D^{DH}(\mathbf{x}) = \frac{3}{\mathbf{x}^3} \ln(1+\mathbf{x}) + \frac{\mathbf{x}^2}{2}$$
 (9)

C learly, Eq. (7) for U (r) can not be used in the RPA approach if a 1. A sthe value of a increases the hardsphere interactions between the charges become more and more important and they should be taken into account. For this purpose we introduce the following form of U (r)

U (r) =
$$\frac{q^2}{r''}$$
 (1 exp(r=a)) (10)

which suppresses electrostatic interactions on the scale of the size of the monomer. The Fourier transform of the above potential can be easily calculated

$$U(k) = \frac{4}{k^2 (1 + k^2 a^2)}$$
(11)

and for the electrostatic free energy given by Eq. (5) we get

$$\frac{F_{el}}{TV} = \frac{1}{4^2} \int_{0}^{2K} \ln 1 + \frac{2(1 + g(k))}{2k^2(1 + k^2a^2)} \frac{2}{k^2(1 + k^2a^2)} k^2 dk$$
(12)

The integral in Eq. (12) can be done analytically for the case of N = 1 and K = 1 which results in the following expression for the free energy

$$\frac{F_{el}^{1}}{TV} = \frac{3}{12} (a)$$
(13)

where

¹ (x) =
$$\frac{1}{x^3}$$
 1 + $\frac{3x^2}{2}$ + $\frac{1+x}{(1+2x)^{1=2}}$ (14)

A nother way to account for the hard sphere nature of the charges is to use the potential (7) but cut integration in (5) at K = 1=a. This results in the following form of the free energy

$$\frac{F_{el}^{\kappa}}{TV} = \frac{3}{12}^{\kappa} (a)$$
(15)

where

^K (x) =
$$\frac{1}{2} \arctan(x^{1}) \frac{\ln(1+x^{2})}{x^{3}}$$
 (16)

In Figure 1 we compare the results for [DH](x), [1](x), and κ (x), given by Eqs. (9), (14), and (16) respectively. The curves presented in the gure show similar behavior as parameter x = a increases. In Figure 2 we also show the phase diagram s obtained on the plane $(;T = a=l_{B})$ using different form of the electrostatic free energy given by Eqs. (8), (13), and (15). The gure clearly shows that the phase diagram obtained using Eq. (13) is in reasonable agreem ent with Debye-Huckel theory where as the one obtained using Eq. (15) di ers signi cantly. This fact leads us to the conclusion that the hard-core interactions between the charges could be correctly taken into account through the potential U (r) given by Eq. (10), but not through the large k cut-o integration parameter K = 1=a. However, in the next section we show that a dierent form of the cut-o K is very important for the proper discription of polyelectrolyte solutions.

IV . POLYELECTROLYTE SOLUTIONS

We rst exam ine the dilute case of in nitely long polyelectrolyte chains. In order to calculate the integral (5)analytically we assume that the macromolecules obey Gaussian statistics. For g(k) de ned by Eq. (6) we use the following approximation

$$g(k) = 1 + \frac{N}{1 + N b^2 k^2 = 12} / 1 + \frac{12}{b^2 k^2}$$
 (17)

where b^2 is the mean-squared distance between the neighboring monom ers of the chain. For simplicity we assume here that b = a. We also use U (r) in the form (7) valid for the dilute case a 1 and we rewrite Eq. (12) with K = 1 in the form

$$\frac{F_{e1}}{TV} = \frac{1}{4^2} \int_{0}^{24} \ln 1 + \frac{2}{k^2} + \frac{6^2}{a^2k^4} \int_{0}^{2} k^2 dk \quad (18)$$

Evaluation of the above integral leads to

$$\frac{F_{e1}}{TV} = \frac{3=2}{12 a^{3=2}} (a)$$
(19)

where

$$(\mathbf{x}) = \frac{12}{(\frac{p}{24} + \mathbf{x})^{1-2}}$$
(20)

The result F_{el} ³⁼² for in nitely long Gaussian chains was previously obtained in a number of papers (for example see Refs. 5,17).

W ith the use of Eq. (19) we take the second derivative of the free energy (2) with respect to the density = $_{+}$ + and in the lim it of ! 0 we nd

$$\frac{\theta^2}{\theta^2} = \frac{F_{el}}{TV} + \frac{1}{2} = \frac{3^{3-2}}{64 b^{3-2-2}} < 0$$
(21)

The negative sign of the second derivative of the free energy shows that dilute solution of in nitely long polyelectrolyte chains are unstable in good solvent conditions.⁷ In Figure 3 we show the phase diagrams on the plane ($;T = a = l_B$) obtained for di erent chain lengths N. The electrostatic contribution to the free energy used to obtain the diagram s is

$$\frac{F_{el}}{TV} = \frac{1}{4^2 a^3} \int_{0}^{2^2} \ln 1 + \frac{(a)^2 (3=2+N k^2=12)}{k^2 (1+k^2) (1+N k^2=12)}$$
$$\frac{(a)^2}{k^2 (1+k^2)} k^2 dk \qquad (22)$$

and it is evaluated num erically. The gure clearly dem onstrates that as N increases the region of instability becomes broader and broader even at very dilute concentrations. The same results can be obtain if we introduce a constant cut-of value K = Ka = 2.

We suggest here that the unphysical behavior described above arises from the inappropriate contribution to the RPA electrostatic free energy at large k. At large k values the chains are perturbed due to the electrostatic interactions. RPA, however, is only valid if the interactions do not modify the chain conformations. Therefore, the cut-o has to be chosen such that we only include the electrostatic contributions that can be accounted by RPA. It is straight forward to determ ine the cut-o given that the major feature of polyelectrolyte solutions is the existence of a concentration dependent correlation length $.^{10}$ On the scales smaller then the chain is stretched and on larger scales it obeys G aussian statistics. For strongly charged polyelectrolytes can be estim ated as^{11;12}

$$=\frac{1}{()^{1=2}}$$
 (23)

In Fourier space scales smaller than are given by ka > 2 = . Since the contributions to the free energy on these scales can not be accounted within the RPA approach, we cut integral (22)

$$K = 2 ()^{1=2}$$
 (24)

V.RESULTS, DISCUSSION AND CONCLUSIONS

In Figure 4 we show the modi ed RPA phase diagram s obtained by evaluating the integral (22) up to ka = K with K given in (24). The modi ed RPA phase diagram has a much lower value of the critical temperature for large N than the standard RPA phase diagram Figure 3. The criticaltem perature initially increases as N increases in the modi ed RPA phase diagram, and it plateaus at very high N values at about $l_{\rm B}$ =b 73. Since most polymers have values of $l_{\rm B}$ =b < 42 at room temperature, salt-free solutions of polyelectrolytes with m onovalent counterions are water soluble at room temperatures in agreem ent with the experiments.

Another im portant feature of the modi ed RPA phase diagram in Figure 4 is the prediction that the critical concentration remains nearly constant as N increases in agreem ent with the simulations. This is only true if the contributions from uctuations of k > K are completely ignored. That is, if we add the RPA electrostatic contribution from the 2 = a > k > K in the free energy using the structure function for rod like units, though the critical tem perature only increases slightly, the critical concentration goes to zero as N increases. This suggests that the charge uctuations at lower length scales do not contribute to the free energy or that they are strongly suppressed. The suppression of charge uctuations on shorter length scales that can be explained by the fact that the electrostatic interactions are not screened at these length scales. Therefore, the counterions and m onom ers are strongly correlated. Indeed, these correlations stretched the chains.

W e argue above that in polyelectrolyte solutions the cut-of is a natural inverse length over which RPA breaks down. It is important to point out, how ever, that a concentration dependent integration cut-o is also obtained in the one component plasm a_{\prime}^{18} in which counterions uctuate over a non-uctuating charged medium. The physical reason and the concentration dependence of the cut-o in the one component plasm a explained in Ref. 18 is di erent than in polyelectrolytes. In a one com ponent plasm a the integration in the electrostatic free energy given by the RPA approach should be carried over a nite number of the wave-vectors k in the same way as in the Debye theory of the speci c heat of solids.¹⁹ Namely, the total number of degrees of freedom in the system 3N = 3V should be equal to the total number of physically di erent m odes with wave-vectors k within the spherical shell of radius K . The num ber of m odes is twice the number of the wave-vectors since each k has a sine a cosine m ode. Therefore, one obtains

$$2V \int_{0}^{ZK} \frac{d^{3}k}{(2)^{3}} = 3N$$
 (25)

which leads to

$$K = (9^{2})^{1=3}$$
(26)

The above argument assumes that the important length scale is the average distance between ions d 2 = K $^{1=3}$, and that this length scales imposes a "periodicity" even though the non uctuating support for the ions is a solid with no structure.

 $1=^{1=3}$ is an impor-The distance between charges d tant length scale to determ ine the break down of DH in simple electrolyte solutions.²⁰ Therefore, in simple electrolytes d is an important length scale as the system gets concentrated when the electrostatic energy between two charges separated by d, $e^2 = (4 \text{ "d}) > K_B T \text{ or } \frac{1}{16} = d > 1$. Interestingly, the integration to a concentration dependent cut-o K ¹⁼³ versus integrating to the inverse hard core size of the ions K 1=a does not a ect significantly the RPA phase diagrams in simple electrolytes. Instead, in salt-free polyelectrolyte solutions a concentration dependent cut-o is essential to recover reasonable values of the critical tem perature as N increases and a N independent critical concentration.

Though we argue that in salt-free polyelectrolyte solutions the natural length scale that determ ines the cut-o is , we obtain very similar phase diagram s if we instead use d as shown in Figure 5. Notice that the importance of a concentration dependent cut-o is at dilute solutions where > d. Therefore, there should be counterion uctuations at length scales d < l <that we did not include in our RPA approach. If we add a RPA contribution from these counterion uctuations (only the counterions) we get negligible corrections to the phase diagram . Our argum ents are in agreem ent with the results in R ef. 21 where a self-consistent polyelectrolyte RPA where the screening length from polyions is wavevector-dependent due to chain connectivity shows that at high wavevectors (short length scales), only counterions contribute to screening. At low erw avevectors, both polyions and counterions contribute to screening.

In sum m ary, we show that in salt-free solutions a wave vector dependent cut-o is essential to recover physical phase diagram s. The cut-o is determ ined by the applicability lim it of the RPA. In the presence of salt the electrostatic contribution to the free energy reduces to the wellknown DH lim iting law.⁵ In that case reasonable phase diagram are obtain without a concentration dependent cut-o, ²² although in principle one should only include uctuations from the monom ers on the length scale where the RPA approach is valid (i.e., larger than).

W eacknow ledge the nancial support of the N IH grant number G M 62109-02 and of the Institute for B ioengineering and N anosciences in A dvanced M edicine (IB – NAM) at N orthwestern U niversity.

¹ C onway, B.E. "Ionic Interactions and A ctivity B ehavior of E lectrolyte Solutions" in B.E.Conway, J.O M. Bockris, E.

Yeager: Comprehensive Treatise of Electrochemistry; v.5. New York: Plenum Press (1983).

- ² Orkoulas, G; Panagiotopoulos, A. Z. J. Chem. Phys. 1999, 110, 1581. Yan, Q.; De Pablo, J.J. J. Chem. Phys. 1999, 111, 9509. Luiten, E.; Panagiotopoulos, A. Z.; Fisher, M. E. Phys. Rev. Lett. 2002, 88, 045701.
- ³ B jerrum, N. Koninklinge Dans. Vidensk. Selsk. 1926 7, 9.
- ⁴ Fisher, M.E.; Levin, Y.Phys. Rev. Lett. 1993, 71, 3826.
- ⁵ Borue, V.Yu.; Erukhim ovich, I.Ya.M acrom olecules 1988, 21, 3240.
- ⁶ Joanny, J. F.; Leibler, L. J. Phys. France, 1990, 51(6), 545.
- ⁷ Mahdi, K. A.; Olvera de la Cruz, M. Macromolecules, 2000, 33, 7649.
- 8 O rkoulas G .; K um ar Sanat K .preprint 2002.
- ⁹ M .M uthukum ar; M acrom olecules, 2002 35, 9142.
- ¹⁰ de G ennes, P G. Scaling C oncepts in P olym er P hysics; C ornell U niv. P ress: 1979.
- ¹¹ Schiessel, H.; Pincus, P. Macrom olecules 1998 31, 7953. Schiessel, H.; Macrom olecules, 1999, 32, 5673.
- ¹² D obrynin, A.V.; Rubinstein, M.M. acromolecules, 1999, 32, 915.
- ¹³ StevensM .J.;, K rem er K .J. C hem . P hys. 1995, 103, 1669.
- ¹⁴ U.M icka, C.Holm, K.K rem er Langm uir 1999, 15, 4033.
- Stevens, M .; K rem er, K .J. Phys. II France, 1997, 6, 1607.
- ¹⁵ M anning G.S.J. Phys. Chem . 88 (1984) 6654
- ¹⁶ P. Gonzales-M ozuelos, M. Olvera de la Cruz: J. Chem. Phys. 1995, 103, 3145.
- ¹⁷ O lvera de la C nuz, M .; B elloni, L .; D elsanti, M .; D albiez, J.; Spalla, O .; D ri ord, M . J. Chem . Phys., 1995, 103, 5781.
- ¹⁸ Brilliantov, N.V.; Malinin, V.V.; Netz, R.R.Eur. Phys. J.D 2002, 18, 339.
- ¹⁹ Z im an, J. M . P rinciples of the T heory of Solids; C laredon P ress: O x ford, 1964.
- ²⁰ B rilliantov, N.V.; K uznetsov, D.V; K lein, R.P hys. Rev. Lett., 1998, 81, 1433.
- ²¹ Donley, J. P.; Rudnick, J.; Liu, A. J. M acrom olecules, 1997, 30, 1188.
- ²² M ahdi,K. A. PhD thesis (N orthwestern University, USA, 2000).




FIG.1. The plots for (x) given by Eq. (9) (solid line), Eq. (14) (dash line), and Eq. (16) (dot line).




FIG.2. Phase diagrams of simple electrolyte solutions obtained using di erent forms of electrostatic free energy given by Eq. (8) (solid line), Eq. (13) (dash line), and Eq. (15) (dot line).




FIG. 3. Phase diagrams of salt free polyelectrolyte solutions obtained within classical RPA approach for di erent lengths of the polymers N. From buttom to top N = 10;20;50;100;1000.





FIG. 4. Phase diagrams of salt free polyelectrolyte solutions obtained within modi ed RPA approach with K given by Eq. (24) for di erent lengths of the polymers N. From buttom to top N = 50;100;1000;10000.



 $\begin{array}{c} Concentration \\ FIG.5. Phase diagram s of salt free polyelectrolyte solutions obtained within modi ed RPA approach with K given by Eq. (26) for di erent lengths of the polymers N. From buttom to top N = 50;100;1000;10000. \end{array}$