## N on-K ram ers degeneracy and oscillatory tunnel splittings in the biaxial Spin System Degang Zhang<sup>1;2;3</sup> and BambiHu<sup>2;1</sup>

<sup>1</sup>D epartm ent of Physics, University of Houston, Houston, TX 77204, USA <sup>2</sup>D epartm ent of Physics and Centre for Nonlinear Studies, Hong Kong Baptist University, Kow loon Tong, Hong Kong, China <sup>3</sup>Institute of Solid State Physics, Sichuan Norm al University, Chengdu 610068, China

## Abstract

We have investigated analytically quantum tunneling of large spin in the biaxial spin system with the magnetic eld applied along the hard and medium an isotropy axes by using a purely quantum -mechanical approach. When the magnetic eld parallels the hard axis, the tunnel splittings of all the energy level pairs are oscillatory as a function of the magnetic eld. The quenching points are completely determined by the coexistence of solutions of the Ince's equation. When the magnetic eld points the medium axis, the tunnel splitting oscillations disappear due to no coexistence of solutions. These results coincide the recent experimental observations in the nanom agnet  $Fe_8$ .

In recent years much attention has been paid to quantum tunneling of magnetization (large spin) in nanom agnets, both from experiment and from theory [1]. Several magnetic particles have been identified as promising candidates for the observation of such macroscopic quantum phenomena, where the magnetization (or the Neel vector) tunnels from one potential minimum to another one. The excellent examples that have widely studied are the molecular nanomagnets  $Fe_8$  [2-5] and  $Mn_{12}$  [6-8], which have the well-de ned structures and magnetic properties. On the one hand, these phenomena are very interesting from a fundamental point of view because they extend our understanding of the transition between quantum and classical behavior. On the other hand, tunneling of the magnetization changes the magnetic properties of the nanomagnets, which has the potential application for the data storage technology, e.g., making qubits – the elements of quantum computers.

Recently W emsdorfer and Sessoli [9] have observed a novel phenom enon, i.e. oscillations of tunnel splittings of the ground and excited states in the nanom agnet Fe<sub>8</sub> described well by the spin H am iltonian [2-5]

$$H = AS_x^2 BS_y^2 g_BS H;$$
(1)

where S is a spin operator, H is the magnetic eld applied in the x-z plane, the spin quantum number s = 10, A 0.092K, B 0.229K, g 2 is the g-factor, <sub>B</sub> is the Bohr magneton. The zero-eld H am iltonian has a biaxial symmetry with hard, easy and medium axes along x, y and z, respectively. W hen H rotates from x to z direction, the oscillations of tunnel splittings of all the level pairs gradually disappear.

In fact, oscillation of the ground state tunnel splitting  $E_s$  of the model (1) with the magnetic eld  $H_x$  (H kx) was predicted by using instanton technique [10]. The tunneling of spin is quenched when

$$h = h (1 \frac{n}{s} \frac{1}{2s}); n = 0;1; ;2s 1:$$
 (2)

Here,  $h = \frac{H_x}{H_c}$ ,  $H_c = \frac{2s(A+B)}{g_B}$  is the critical eld at which the energy barrier vanishes,  $h < h < h = \frac{A}{A+B}$ , and s is an integer or half odd integer. This kind of topological quenching is the result of quantum interference of di erent instanton paths within the context of acroscopic quantum tunneling [11, 12], and need not be related to K ram ers's degeneracy. Up to the rst order of the rate  $\frac{B}{A+B}$ , the form ula (2) was rederived by quantum -m echanical perturbation theory [13]. Very recently, G arg extended his previous work [10] to the excited states by using a discrete W entzel-K ram ers-B rilbuin approach [14]. To order s  $^{1}$ , the quenching points for the excited state pairs are the same as those for the ground state pair. W ith increasing the magnetic ed, the quenching points gradually decrease and nally disappear. These results were also obtained by the potential eld description of spin systems with exact spin-coordinate correspondence [15] and the instant technique [16, 17]. Because quantum tunneling of spin in the nanom agnets was observed at very low tem peratures, the quantization of spin levels becom es very im portant to explain well the experiments. In essence, the energy spectrum of the spin systems can help us to understand the mechanism of spin tunneling thoroughly. In this paper, we have analytically diagonalized the H am iltonian (1) in the fram ework of the Schrodinger's picture of quantum mechanics. The energy spectrum of the spin model is obtained in the large-s limit. It is clearly shown that when H k x, the tunnel splittings of the ground and excited state pairs are oscillatory as a function of H x and the quenching points agree with the num erical simulation of model (1). When H k z, the tunnel splitting oscillations of all the energy level pairs disappear, which coincided

with those obtained by the phase space path integral [18]. These phenom ena have also been observed in the experiments [9, 19].

Let E and  $_{\rm m}$  be the eigenenergies and eigenstates of H , respectively, then the eigenvalue equation in the basis js;m  $>_{\rm z}$  reads

$$u_{m 1 m 2} + u_{m+1 m+2} t_{m \frac{1}{2} m 1} t_{m+\frac{1}{2} m+1} + f E + \frac{1}{2} (A B) [s(s+1) m^{2}] g_{B} H_{z} m g_{m} = 0;$$
(3)

where

$$u_{m 1} = \frac{1}{4} (A + B)^{q} \overline{[s(s+1) (m 1)^{2}]^{2}} (m 1)^{2};$$

$$t_{m \frac{1}{2}} = \frac{1}{2} g_{B} H_{x} s(s+1) (m \frac{1}{2})^{2} + \frac{1}{4}$$

. O by iously, it is very di cult to strictly solve the equation (3) for arbitrary s. However, in the large-s lim it, Eq. (3) becomes [20]

$$(1 \quad x^{2})\frac{d^{2}}{dx^{2}} \quad 2x\frac{d}{dx} + \left[\frac{E}{A+B} \quad \frac{1}{4} \quad \frac{1}{4}\frac{1}{1}\frac{1}{x^{2}} + \frac{As(s+1)}{A+B}(1 \quad x^{2}) \quad \frac{g_{B}H_{z}}{S(s+1)}\frac{s(s+1)}{A+B}x - \frac{g_{B}H_{x}}{S(s+1)}p\frac{1}{1}\frac{1}{x^{2}} = 0; \qquad (4)$$

Here,  $x = \frac{p}{s} \frac{m}{s(s+1)}$  and only the leading term s are remained, i.e. 0 (s<sup>-1</sup>). In deriving Eq. (4), we have used

$$u_{m \ 1 \ m \ 2} = u_{m \ 1} \left[ u_{m \ 1} - u_{m \ 2} - u_{m \ 1} \right] + \frac{q}{\frac{m \ 1}{s \ (s + 1)}} + \frac{m \ 1}{2s \ (s + 1)} + \frac{1}{2s \ (s + 1)} + \frac{1}{2$$

$$+ \frac{u}{2s(s+1)} \frac{d^{2}}{dx^{2}} + ;$$

$$t_{m \frac{1}{2} m+1} = t_{m \frac{1}{2}} \left[ \begin{array}{c} m \frac{1}{2} \end{array} - \frac{q}{2} \frac{\frac{m}{m \frac{1}{2}}}{\frac{1}{2} s(s+1)} + \end{array} \right]$$

$$= t \frac{q}{2} \frac{1}{s(s+1)} \frac{d}{dx} (t) + \frac{q}{2} \frac{t}{s(s+1)} \frac{d}{dx} + ;$$

where

$$u = \frac{1}{4}(A + B) \begin{bmatrix} q \\ s^{2}(s + 1)^{2}(1 + x^{2})^{2} & s(s + 1)x^{2} \\ \frac{1}{4}(A + B) \begin{bmatrix} s(s + 1)(1 + x^{2}) & x^{2} \end{bmatrix} \begin{bmatrix} 2(1 + x^{2}) \end{bmatrix}$$

and t  $\frac{1}{2}$ g <sub>B</sub>H<sub>x</sub>  $\frac{q}{s(s+1)}$   $\frac{p}{1-x^2}$  for large s.

Taking the transformations: =  $(1 \times 2)^{\frac{1}{4}} y(x)$  and  $x = \sin(2t)$ , and substituting them into Eq. (4), we nally obtain the Hill's equation [21]

$$\frac{d^2y}{dt^2} + [ + (a \ b)\cos(2t) + (c \ b)\sin(2t) + \frac{b^2}{8}\cos(4t)]y = 0;$$
 (5)

where

$$= \frac{4E + 2A s (s + 1)}{A + B}; a = b \quad \frac{4g_{B} H_{x} \overline{s(s + 1)}}{A + B}; b = 4 \quad \frac{A s (s + 1)}{A + B}; c = b \quad \frac{4g_{B} H_{z} \overline{s(s + 1)}}{A + B}:$$
(6)

Up to now, we have mapped the spin problem (1) onto a particle problem (5). The energy spectrum of H is completely determ ined by the characteristic levels of the H ill's equation. We note that when A = 0 and H = 0,  $= m^2$ . So  $E = B \left(\frac{m}{2}\right)^2$  are nothing but the eigenvalues of the H am iltonian (1) with integers for even m or with half odd integers for odd m.

For Eq. (5), there exist two monotonically increasing sequences of real numbers  $a_0;a_{2i};b_{2i};a_{2i}^0$  and  $b_{2i}^0$  (i = 1;2; ) such that Eq. (5) has a solution with period if and only if =  $a_0;a_{2i}$  or  $b_{2i}$ , and a solution with

period 2 if and only if  $= a_{2i}^0$  or  $b_{2i}^0$ . The  $a_0; a_{2i}; b_{2i}; a_{2i}^0$  and  $b_{2i}^0$  satisfy inequalities:

$$a_0 < b_1^0 \quad a_1^0 < b_2 \quad a_2 < b_3^0 \quad a_3^0 < b_4 \quad a_4 <$$
 : (7)

A coording to the relation between E and in Eq. (6), it is easy to see that the ground state of H corresponds to the allowed highest characteristic level of the H ill's equation with period or 2, depending on integer or half odd integer spin s. The lower the characteristic level of Eq. (5) is, the higher the associated eigenstate of the H am iltonian (1) is. The tunnel splitting of the characteristic level pair of Eq. (5) with period is

$$a_{2i} = a_{2i} \quad b_{2i};$$
 (8)

and the tunnel splitting of the characteristic level pair of Eq. (5) with period 2 is

$${}^{0}_{2i\ 1} = a^{0}_{2i\ 1} \qquad b^{0}_{2i\ 1}:$$
(9)

The tunnel splitting of the energy level pair of the H am iltonian (1) can be evaluated by the tunnel splitting of its associated characteristic level pair using Eq.(6). We note that when two of three parameters A, H<sub>x</sub> and H<sub>z</sub> vanish, the Hill's equation (5) becomes the well-known M athieu equation [22]. So the tunnel splittings of all the energy level pairs of the H am iltonian (1) are monotonous increasing rather than oscillatory in the three cases [20]. For arbitrary parameters A, B, H<sub>x</sub> and H<sub>z</sub>, it is di cult to solve analytically Eq. (5). Here we only consider the following two special cases.

(i)  $H_z = 0$  (i.e. c = b). In this case, Eq. (5) reduces to the Ince's equation, which has been studied in a set of literature due to its physically basic in portance [21]. To observe the oscillations of tunnel splittings of model (1), we must not some vanishing points at which its eigenstates are degenerate, i.e.  $a_{2i} = b_{2i}$  or  $a_{2i \ 1}^0 = b_{2i \ 1}^0$ . This is equivalent to not the coexistence of solutions of the Ince's equation, which means that there exist two linearly independent solutions (one even and one odd) with period or 2. Very fortunately, due to the positive coe cient of the last term cos(4t), the Ince's equation has the coexistence of solutions [21] with period when

$$a = 2nb \tag{10}$$

and with period 2 when

$$a = (2n \ 1)b;$$
 (11)

where n = 0; 1; 2; . O by iously, the quenching points of tunnel splittings of the characteristic level pairs with period are exactly shifted by half a period relative to those with period 2. Under the conditions (10) and (11), the Ince's equation becomes the W hittaker equation [21]

$$\frac{d^2 y}{dt^2} + [pbcos(2t) + \frac{b^2}{8}cos(4t)]y = 0:$$
(12)

Here, p = 2n + 1 or 2n for Eq. (10) or (11), respectively. For Eq. (12), when b ! 0, then  $a_0 ! 0$ ,  $a_{2i}$  and  $b_{2i} ! (2i)^2$ , and  $a_{2i 1}^0$  and  $b_{2i 1}^0 ! (2i 1)^2$  [23].

Integer spin s. Due to Eq. (6), the eigenstates of the Ham iltonian (1) correspond to the characteristic levels of the Ince's equation with period , i.e.  $E_0^a = \frac{1}{4}(A + B)a_0 + E_0; E_1^a = \frac{1}{4}(A + B)a_{2i} + E_0$  and  $E_1^b = \frac{1}{4}(A + B)b_{2i} + E_0; E_0 = \frac{1}{2}As(s+1); i = 1; 2;$ ; s. So the tunnel splitting of the energy level pair  $(E_1^a; E_1^b)$  is

$$E_{i} = E_{i}^{b} \quad E_{i}^{a} = \frac{1}{4}(A + B) \quad _{2i}$$
 (13)

From Eq. (10), we have  $E_i = 0$  when

$$H_{x} = \frac{(2n+1)^{q} \overline{A(A+B)}}{g_{B}};$$
 (14)

The existence of the quenching elds (14) clearly shows that the tunnel splittings of all the energy levels of H are oscillatory as a function of H<sub>x</sub> and the period of oscillations  $H = \frac{2^{\frac{1}{A}} \frac{A+B}{A+B}}{g_{B}}$ . This coincides with G arg's result (2) [10, 14]. For Eq. (12), when  $\dot{p}j=21+1$ , then the even intervals of instability on the axis disappear with at most 1+ 1 exceptions [21]. In other words, the characteristic values a<sub>0</sub>; a<sub>21</sub> and b<sub>21</sub> satisfy

$$a_0 < b_2 < a_2 < f_1 \ b \ a_{21}$$
  
 $< b_{2(l+1)} = a_{2(l+1)} < f_2 \ b \ a_{2s}$ : (15)

Thism eans that the tunnel splittings of the lhighest excited state pairs of the H am iltonian (1) do not vanish while those of the others lenergy level pairs vanish. With increasing the magnetic eld  $H_x$  (i.e. jp), the quenching points gradually decrease and nally disappear when 1 s. The con guration of

the quenching points agree with that from the numerical simulation of the Ham iltonian (1) (see Fig. 1).

Halfodd integer spin s. The eigenstates of the Ham iltonian (1) correspond to the characteristic levels of Eq. (12) with period 2 , i.e.  $E_{i}^{a^{0}} = \frac{1}{4} (A + B)a_{2i}^{0} + E_{0}$  and  $E_{i}^{b^{0}} = \frac{1}{4} (A + B)b_{2i}^{0} + E_{0}$ ; i = 1; 2;;  $\frac{1}{2}$  + The tunnel splitting of the energy level pair ( $E_{i}^{a^{0}} = \frac{1}{2}; E_{i}^{b^{0}} = \frac{1}{2}$ ) reads

$$E_{i}^{0} = E_{i}^{b^{0}} = E_{i}^{b^{0}} = \frac{1}{2} = \frac{1}{4} (A + B) = \frac{0}{2i} = \frac{1}{2} (A + B)$$
(16)

According to Eq. (11), we obtain  $E_{i \frac{1}{2}}^{0} = 0$  when

$$H_{x}^{0} = \frac{2n^{q} \overline{A(A+B)}}{g_{B}}; \qquad (17)$$

Obviously, the period of oscillations of tunnel splittings for all the energy level pairs  $(E_{i}^{a^{0}}, E_{i}^{b^{0}}, E_{i}^{b^{0}})$  is  $H^{0} = \frac{2^{b} A(A+B)}{g_{B}}$ , which is the same with that of the energy level pairs  $(E_{i}^{a}; E_{i}^{b})$ . However, the quenching points for half odd integer spin s are shifted half a period (i. e.  $\frac{A(A+B)}{g_{B}}$ ) relative to those for integer spin s. For Eq. (12), if pj = 21, then at most 1+ 1 odd interval of instability on the axis remain, i.e.

$$b_{1}^{0} < a_{1}^{0} < b_{3}^{0} < a_{3}^{0} < \qquad \stackrel{0}{21} b < a_{21 \ 1}^{0}$$
$$< b_{21+1}^{0} = a_{21+1}^{0} < \qquad \stackrel{0}{25} b a_{25}^{0} : \qquad (18)$$

It is easy to see that the tunnel splittings of the s  $1+\frac{1}{2}$  by est energy level pairs vanish, but the tunnel splittings of the other l highest energy level pairs do not. When  $1 > s = \frac{1}{2}$ , there do not exist the quenching points. These results also coincide with those obtained by the discrete W KB approach [14].

(ii)  $H_x = 0$  (i.e. a = b). Let  $t + \frac{1}{4}$ , then Eq. (5) becomes the Ince's equation

$$\frac{d^2y}{dt^2} + [ + (c \ b)\cos(2t) \ \frac{b^2}{8}\cos(4t)]y = 0:$$
(19)

Because the coe cient of the last term  $\cos(4t)$  is negative, Eq. (19) does not possess the coexistence of solutions, i.e. two linearly independent solutions

[21]. This means that the sign of equality in the inequalities (7) cannot hold and each energy level of H is singlet in the parameter space. So the tunnel splittings of all the level pairs are not oscillatory with H<sub>z</sub>, which coincide the experiment [9, 19].

In conclusion, we studied quantum tunneling of large spin in the biaxial spin systems by using quantum mechanics. The energy spectrum of the spin model (1) is obtained by solving H ill's equation (5), which is derived directly from the eigenvalue equation of the spin problem in the large-s limit. It is surprising that when H k x, the vanishing points of the tunnel splittings of all the energy level pairs obtained here by the coexistence of solutions of the Ince's equation coincide those given by the W K B method [10, 14] and other approaches [15-17]. However, our approach is a natural way of explaining the oscillations of tunnel splitting in the biaxial spin system, which is also applied in other spin H am iltonians.

This work was supported in part by grants from the Hong Kong Research G rants Council (RGC), the Hong Kong Baptist University Faculty Research G rant (FRG), the Sichuan Youth Science and Technology Foundation and the NSF of the Sichuan Educational Commission.

## References

- For an overview, see e.g., Quantum tunneling of Magnetization, edited by L.Gunther and B.Barbara (Kluwer, Dordrecht, The Netherlands, 1994), and references therein.
- [2] A.-L.Barra, P.Debrunner, D.Gatteschi, C.E.Schulz, and R.Sessoli, Europhys. Lett. 35, 133 (1996).
- [3] C.Sangregorio, T.Ohm, C.Paulsen, R.Sessoli, and D.Gatteschi, Phys. Rev.Lett. 78, 4645 (1997).
- [4] R. Caciu o, G. Amoretti, A. Murani, R. Sessoli, A. Caneschi, and D. Gatteschi, Phys. Rev. Lett. 81, 4744 (1998).
- [5] W. Wernsdorfer, T. Ohm, C. Sangregorio, R. Sessoli, D. Mailly, and C. Paulsen, Phys. Rev. Lett. 82, 3903 (1999).

- [6] J.R.Friedman, M.P.Sarachik, T.Tejada, and R.Ziolo, Phys. Rev. Lett. 76, 3830 (1996).
- [7] S. Hill, J. A. A. J. Perenboom, N. S. Dalal, T. Hathaway, T. Stakup, and J.S. Brooks, Phys. Rev. Lett. 80, 2453 (1998).
- [8] G.Bellessa, N.Vemier, B.Barbara, and D.Gatteschi, Phys. Rev. Lett. 83, 416 (1999).
- [9] W. Wernsdorfer and R. Sessoli, Science, 284, 133 (1999).
- [10] A.Garg, Europhys. Lett. 22, 205 (1993).
- [11] D.Loss, D.P.D iV incenzo, and G.G rinstein, Phys. Rev. Lett. 69, 3232 (1992).
- [12] J. von Delft and C. L. Henley, Phys. Rev. Lett. 69, 3236 (1992).
- [13] S.W eigert, Europhys. Lett. 26, 561 (1994).
- [14] A.Garg, Phys. Rev. Lett. 83, 4385 (1999).
- [15] J.-Q. Liang, H. J.W. Muller, D.K. Park and P.-C. Pu, Phys. Rev. B 61, 8856 (2000).
- [16] Lu Rong, Zhu Jia-Lin, Zhou Yi and Gu Bing-Lin, Phys. Rev. B 62, 11661 (2000).
- [17] E.M. Chudnovsky and X.Martinez-hidalgo, Europhys. Lett. 50, 395 (2000).
- [18] M. Enz and R. Schilling, J. Phys. C: Solid State Phys. 19, 1765, L711 (1986).
- [19] E. Barco, N. Vernier, J. M. Hernandez, J. Tejada, E. M. Chudnovsky, E. Molins and G. Bellessa, Europhys. Lett. 47, 722 (1999).
- [20] Bambi Hu, De-gang Zhang and Bo-zang Li, Phys. Rev. B 61, 8639 (2000).
- [21] W .M agnus and S.W inkler, Hill's Equation, (D over Publications, Inc., New York, 1979).

- [22] M.Abram ow itz and I.A.Stegun, Handbook of Mathematical Functions, (Now York: Dover, 1965).
- [23] E.L. Ince, Proc. Lond. M ath. Soc. 25, 53 (1926).
- Fig. 1 The exact quenching points of all the level pairs of the H am iltonian (1) with s = 10; A = 0.092K; B = 0.229K; and  $H_z = 0$ .

