Description of Quantum Systems by Random Matrix Ensembles of High Dimensions
"Proceedings of ICSSUR'6, Sixth
International Conference on Squeezed States and Uncertainty Relations, 24th May 1999—29th May 1999, Universita degli Studidi Napoli 'Federico II', Naples, Italy", NASA, Greenbelt, Maryland, USA (1999).

Maciej M. Duras
Institute of Physics, Cracow University of Technology, ulica Podchorazych 1, 30-084 Cracow,
Poland
AD 1999, July 26th

Abstract

The new Theorem on location of maximum of probability density functions of dimensionless second dierence of the three adjacent energy levels for N-dimensional Gaussian orthogonal ensemble GOE(N), N-dimensional Gaussian unitary ensemble GUE(N), N-dimensional Gaussian symplectic ensemble GSE(N), and Poisson ensemble PE, is formulated: The probability density functions of the dimensionless second dierence of the three adjacent energy levels take on maximum at the origin for the following ensembles: GOE(N), GUE(N), GSE(N), and PE, where N 3. The notions of level homogenization with level clustering and level homogenization with level repulsion are introduced.

M any com plex N -level quantum systems exhibiting universal behaviour depending only on sym metry of H am iltonian matrix of the system are divided into: G aussian orthogonal ensemble GOE (N), or G aussian unitary ensemble GUE (N), or G aussian sym plectic ensemble GSE (N). The G aussian ensembles are used in study of quantum systems whose classical limit analogs are chaotic. The Poisson ensemble PE (Poisson random -sequence spectrum) is composed of uncorrelated and random ly distributed energy levels and it describes quantum systems whose classical-limit analogs are integrable. The standard statistical measure is Wigner's distribution of the ith nearest neighbour spacing:

$$s_i = {}^{1}E_i = E_{i+1} \quad E_i; \quad i=1,...,N-1$$
: (1)

For ith second di erence (the ith second di erential quotient) of the three adjacent energy levels:

$${}^{2}E_{i} = {}^{1}E_{i+1}$$
 ${}^{1}E_{i} = E_{i} + E_{i+2}$ $2E_{i+1}; i = 1,...,N-2;$ (2)

we calculated distributions for GOE (3), GUE (3), GSE (3), and PE Refs [1{3]. We form ulate the following

Theorem: The probability density functions of the dimensionless second dierence of the three adjacent energy levels take on maximum at the origin for the following ensembles: GOE (N), GUE (N), GSE (N), and PE, where N 3.

We present the idea of proof. For Gaussian ensembles it can be shown that second dierence distributions are symmetrical functions for N 3. Hence, the rst derivatives of the distributions at the origin vanish. For Poisson ensemble the second dierence distribution is Laplace one for N 3. Therefore, the distribution takes on maximum at zero.

The inferences are the following:

- 1. The quantum systems show tendency towards the homogeneity of levels (equal distance between adjacent levels). We call it homogeneization of energy levels.
- 2. There are two generic hom ogeneizations: the rst is typical for G aussian ensembles, the second one for Poisson ensemble. For the former ensembles we de ne level homogenization with level repulsion as follows. Energy levels are so distributed that the situation that both the spacings and second di erence vanish:

$$^{2}E_{i} = s_{i} = s_{i+1} = 0;$$
 (3)

is the most probable one. For the latter ensemble level homogenization with level clustering is described below. Now it is the most probable that only the second dierence is equal to zero but the two nearest neighbour spacings are nonzero:

$${}^{2}E_{i} = 0; s_{i} = s_{i+1}; s_{i} \in 0:$$
 (4)

- 3. The assumption of non-zero value by the second dierence is less probable than the assumption of zero value. Equivalently, the inequality of the two nearest neighbour spacings is less probable than their equality.
- 4. The predictions of the Theorem are corroborated by numerical and experimental data Refs [1{3].

The theorem could be extended to other ensembles, e.g. circular ones, and it is a direction of future development.

REFERENCES

- [1] M.M. Duras and K. Sokalski, Phys. Rev. E 54, 3142 (1996).
- [2] Maciej M. Duras, Finite di erence and nite element distributions in statistical theory of energy levels in quantum systems, Ph.D. thesis, Jagellonian University, Cracow, July 1996.
- [3] M.M. Duras and K. Sokalski, Physica D 125, 260 (1999).