# Slow Relaxation in a Constrained Ising Spin Chain: a Toy M odel for G ranular Com paction

Satya N.Majum dar and David S.Dean

Laboratoire de Physique Quantique (UMR 5626 du CNRS), Universite Paul Sabatier, 31062 Toulouse Cedex, France

We present detailed analytical studies on the zero tem perature coarsening dynam ics in an Ising spin chain in presence of a dynam ically induced eld that favors locally the ' ' phase com pared to the ' ' phase. We show that the presence of such a local kinetic bias drives the system into a late time state with average magnetization m = 1. However the magnetization relaxes into this nal value extremely slow ly in an inverse logarithm ic fashion. We further map this spin model exactly onto a simple lattice model of granular compaction that includes the minimal microscopic moves needed for compaction. This toy model then predicts analytically an inverse logarithm ic law for the growth of density of granular particles, as seen in recent experiments and thereby provides a new mechanism for the inverse logarithm ic relaxation. Our analysis utilizes an independent interval approximation for the particle and the hole clusters and is argued to be exact at late times (supported also by numerical simulations).

PACS num bers: 05.40.-a, 82.20 M j

I. IN TRODUCTION

Slow relaxation dynam ics naturally occurs in system s with quenched disorder such as spin glasses and has remained a subject of long standing interest [1]. However, systems without quenched disorder such as structural glasses also exhibit slow dynamics. It is believed that the slow relaxation in the latter systems is due to kinetic disorder, induced by the dynam ics itself [2]. Another in portant class of system s without quenched disorder is granular m aterial, where once again kinetic disorders are responsible for slow relaxation. In a recent experim ent  $[\beta]$ , a cylinder packed loosely with glass beads was tapped mechanically and it was found that the system gets more and more compact with time. However, the density (t) com pacti ed rather slow ly with time as, 1=log(t). How robust is this inverse loga-(1)(t) rithm ic relaxation? Is it only speci c to granular system s ordoes this also occur in other out of equilibrium system s in presence of kinetic disorders?

In this paper we study, in detail, the e ect of kinetic disorders on the relaxation dynam ics of an Ising system quenched from a high tem perature disordered phase into the low temperature ordered phase. In absence of kinetic disorders, the dynamics of such a system is well. understood [4]. As time progresses, dom ains of equilibrium low-tem perature ordered phases (consisting predom inantly of up and down spins respectively) form and grow. The average linear size of a dom ain grows with t for zero tem perature nonconserved dissitimeasl(t) pative dynam ics. How does disorder, quenched or kinetic, a ect this simple dynamics? The e ect of quenched ferrom agnetic disorder on this phase ordering kinetics has been studied extensively [4]. Essentially the quenched disorder tends to pin the dom ain walls leading to com plete freezing at T = 0. However at sm all nonzero tem - peratures, the dom ains still coarsen via activated dynamics but extrem ely slow ly as, l(t)  $(\log t)^{1-4}$  [4]. The purpose of this paper is to explore the e ect of kinetic disorder on the phase ordering dynam ics. A shorter version of this paper with the main analytical results and their num erical con mation has appeared elsewhere [5]. Here we explore the dynam ics in more detail and also elaborate its connection to other system s such as granular material and reaction-di usion system s.

In this paper, we study the zero tem perature dynam ics of an Ising chain in presence of a specic type of kinetic disorder namely a dynamically induced magnetic eld. If a sm all uniform external eld (say in the down direction) is put on in an Ising system following a rapid quench from in nite temperature to T = 0, the system rapidly relaxes into a pure state with all spins down. In this case, the symm etry between the two ordered pure states is broken globally. Interesting physics happens when, instead of a global external bias, the symmetry between the pure states is broken locally by the dynamics itself. In this paper, we investigate the e ect of this particular kinetic disorder and show that this system also gives rise to a very slow dynam ics. In particular, it gives rise to inverse logarithm ic relaxation (ILR) of magnetization, very sim ilar to the density compaction in granular system s. Our study therefore suggests that the ILR is a very robust phenom enon and not just lim ited to granular system s or speci c types of kinetic disorders.

The paper is organized as follows. In Section-II, we dene ourm odel precisely and sum marize the main results. In Section-III, we establish the connection between our spin model and a lattice model of granular compaction. We also show that ourm odel can be viewed as a new one dimensional reaction-di usion model when the dynam ics is described in terms of the kinks between domains of opposite phases. In Section-IV, we derive some exact results. In section-V, we analyze the dynam ics via an independent interval approximation (IIA). In Section-VI, we argue that the IIA results become exact in the limit when the initial volume fraction of one of the phases is small. In Section-VII, we extend this argument to other volume fractions as well. In Section-VIII, we present a heuristic approach to support the IIA results. Finally we conclude in Section-IX.

# II. THE MODEL AND THE OVERVIEW OF RESULTS

We consider a simple Ising spin chain with spins  $S_i = 1$ . Starting from a given initial conguration the system evolves by single spin ip continuous time dynamics. The rate of ipping of a given spin depends on the its neighboring spins. We denote the rate of spin ip  $S_i$ !  $S_i$  by W ( $S_i; S_{i-1}; S_{i+1}$ ) where  $S_{i-1}$  and  $S_{i+1}$  are the two neighboring spins. In our model the rates are specied as follow s:

W (+;++) = W (; ) = 0  $W (+; +) = W (+;+) = W (; +) = W (; +) \frac{1}{2}$  W (+; ) = 1 W (;++) = (1)

Note that the case = 1 corresponds to the usual zero tem perature G lauber dynam ics [6] which preserves the symmetry between the up and down phases. However, if < 1, the ipping of a down spin sandwiched between two up spins is not as likely as the ipping of an up spin sandwiched between two down spins. Thus < 1 clearly breaks the symmetry between the up and down phases. However this symmetry is broken only dynamically, i.e. not everywhere but only at at the location of the triplets (+; ;+). Thus the isolated ` ' spins (surrounded on both sides by a `+') tend to block the coalescence of `+' domains and locally favor the ` ' spins. One can argue that the asymptotic dynam ics at late times is similar for any < 1. In other words, = 0 is an attractive xed point. We therefore restrict ourselves only to the case

= 0. To see the e ect of the local dynam ical constraint m ore

precisely, we derive (following G lauber's calculation for = 1 [6]) the exact evolution equation for the magnetization m (t) = hS<sub>1</sub>i for the = 0 case:

$$\frac{d}{dt}hS_{i}i = 2P(1; 1;1);$$
(2)

where P ( i 1; i; i+1)(t) = h(1+ i 1Si 1)(1+ iSi)(1+ i+1Si+1)i=8 is the three point probability to nd the sequence of spins ( i 1; i; i+1) about the site i and we have used translational invariance. Hence P (1; 1;1)(t) denotes the probability of the occurrence of the triplet + + ' at time t. We note that for the case = 1,  $dhS_i i=dt = 0$  [6], indicating that the magnetization does not evolve with time. In our case, due to the triplet defects 4 + ', the average m agnetization decays with time. If L (t) denote the fractions of +' and  $\cdot'$  spins, m)=2 we nd from Eq. (2), then using L = (1) $dL = dt = R_1$  (t) where  $R_1$  (t) is the number density of the triplets of type '(+ +)' per unit length, clearly exhibiting the asymmetry generated by the '(+ +)' triplets. We also note that unlike the = 1 case, the evolution equation (2) for the single point correlation function involves two and three point correlations (via  $R_1$  (t)). This hierarchy makes an exact solution di cult for = 0.

It is useful at this point to sum marize the main results obtained in this paper. Let us rst highlight the contrast between the = 1 (no kinetic disorder) and the = 0(with kinetic disorder) cases. For = 1, due to the preserved symmetry between the up and down phases at all times, the average domain size of both "+" and " " do $t^{=2}$  at late times [7]. Thus the mains grow as 1 (t) average m agnetization m (t) = (1,1)=(1 + 1) is a constant of m otion [6]) and stays xed at its initial value. In particular, if we start from an initial state where the spins are random (in nite tem perature), the initial average magnetization is zero and stays 0 at all subsequent times. In contrast, for = 0 case where the symmetry is dynam ically broken, we nd the following results.

(1) For = 0, while domains of both phases continue to grow with time, they have dimensional growth laws. The average domain sizes of the + phases (denoted by 1, (t)) and phases (denoted by 1 (t)) coarsen at late times in the following manner: (i) 1, (t) T at late times and (ii) 1 (t)  $t^{=2}\log(bt)$  where b is a number depending on the initial volume fraction of the '+' spins which we will calculate explicitly (see below). In fact, the main result we show below is that the ratio of the two length scales behaves at late times as,

$$\frac{1}{L} = \frac{1}{L} \frac{\log (bt=t_0)}{\log b} \qquad 1 \tag{3}$$

where is the initial volume fraction (need not be sm all) and  $t_0$  is some initial time after which scaling starts holding. The equation (3) explicitly rejects the eject of broken symmetry. Thus due to the dynamically generated local bias, the ' ' domains grow slightly faster than the ' ' domains. We also point out that in contrast to the spin m odels studied in the context of glassy systems [2,8], the dynamics in our m odel does not freeze at zero tem perature, rather the domains coarsen inde nitely in an in nite system.

(2) Consequently, the magnetization m (t) = (1, 1)=(1, +1) decays at late times as,

$$m(t) = 1 + \frac{2 \log b}{\log (bt=t_0)}$$
: (4)

(3) Evidently the number of 4' domains is same as the number of of ' domains since they alternate on the 1-d lattice. The density of domains of either 4' or ' ' type per unit length: N (t) = 1=[1 + 1, ] decays as,

$$\frac{N(t)}{N(t_0)} = \frac{t_0}{t} \frac{\log b}{\log (bt)};$$
(5)

We also compute explicitly two other densities that play a som ewhat central role in our analysis:

(4)  $r_1$  (t): G iven that a ' ' domain has occurred, the probability that it is of length 1. We show below that

$$r_{1}(t) = \frac{p}{t \log(bt=t_{0})}$$
(6)

Note that the amplitude  $p^{p}$  is universal and independent of volume fraction.

(5)  $p_1$  (t): G iven that a +' dom ain has occurred, the probability that it is of length 1. W e show that

$$p_1(t) = \frac{1}{2t} + \frac{1}{t \log(bt=t_0)}$$
 (7)

Once again the amplitude 1 of the correction term is also universal and independent of volume fraction.

The constant b in the above equations can also be computed exactly, and we show that,

$$b = \exp \frac{p}{r_1(t_0)} \frac{p}{t_0}$$
 (8)

# III. CONNECTION TO GRANULAR COMPACTION AND REACTION DIFFUSION SYSTEM S

We now establish a one to one mapping between our spin model (with = 0) and a simple lattice model of granular compaction. Let us consider a (1 + 1)dim ensional granular packing where the grains are represented by square blocks. The pack consists of horizontal layers consisting of blocks and voids (see Fig 1). We focus on the active' layer, i.e., the sst horizontal layer that is not fully compact with blocks as we go up from the bottom of the pile. Below this active layer, all layers are com pact and rem ain com pact under vertical tapping, ie, their dynam ics is completely frozen. All the activities take place in or above the 'active' layer. W e identify this 'active' layer with the one dimensional lattice of the spin model. This active layer consists of sequences of blocks (particles) and voids (holes). We identify a unit block or a particle in the active layer as a ' ' spin of our spin model. Similarly a hole is identied as a "+ ' spin of the spin model.

As the system is tapped vertically, the particles in the active layer can undergo the following moves: (1) In the

interior of a row of consecutive particles in the active layer, there is no e ect of tapping as the system is com – pletely jammed there. The e ect of vertical tapping is felt only at the edges of a particle cluster. The particles at the edges, under tapping, can move up or 'tap' up from the active layer to the layers above the active layer (see Fig. 1). However if the cluster consists of an isolated particle sandwiched between two holes, it has no place to move up, so it stays at its original location. (2) Tapping also can 'roll' o a particle residing in a layer above the active layer, into the active layer, at the edges of the supporting cluster (see Fig. 1).

Let us now see what the rates of spin ips in Eq. (1) in the spin m odel in ply for the com paction m odel. The rate W (+;++) = 0 in plies that if we have three consecutive holes in the active layer, then a particle can not be deposited in the middle site under tapping. This is because a new particle can appear into the active layer only at the edges of the clusters of particles, but not in the middle of a cluster of voids. The rate W (; ) = 0 in plies that if we have three consecutive particles in the active layer, the middle particle can not move up under tapping since it is completely jammed. The rate W (+; +) = 1=2 signi es that if we have three consecutive sites in the active layer consisting respectively of a particle, hole and a hole, then a particle can appear in the middle hole with rate 1=2. This is 'rolling' o move to the right at the edge of a cluster, as discussed in the previous paragraph. Sim ilarly W (+;+ ) = 1=2 signies the 'rolling' o move to the left. The rate W (;+) = 1=2 im plies that when we have three consecutive sites in the active layer consisting respectively of a hole, particle and a particle, the middle particle can disappear (into the layers above) with rate 1=2. This is the 'tapping' up move to the right from the edge of a cluster as discussed in the previous paragraph. Similarly W (; +) = 1=2 signies the rate of tapping'up to the left from the edge of a cluster. The rate W (+; ) = 1 indicates that if we have a sequence of particle, hole, particle in the active layer, the middle hole can be led up with a particle with rate 1. This is simply the addition of two 'rolling' o rates from the left and the right of the middle hole. Finally the rate W(; + +) = = 0 im plies that if we have a sequence of hole, particle, hole in the active layer, the middle particle can not disappear. This is because an isolated particle has no place to move or 'tap' up as discussed in the previous paragraph. This last rate indeed breaks the particle hole symmetry.

Note that the rates  $W(+; +) = W(+; +) = W(+; +) = W(; +) = W(; +) = W(; +) = \frac{1}{2}$  correspond to the di usion of dom ain walls in the spin model. In the compaction model, these are indeed the moves induced by the mechanical tapping. If these rates were zero, i.e., no tapping, then the dynamics in the active layer would freeze after all the isolated holes are led up and the system will be stuck in a metastable con guration and hence the

com paction will stop. It is these di usion moves that lift the system out of a metastable con guration and the system continues to compactify, though extrem ely slow ly. Identifying the  $\Upsilon$  (+) spins with particles (holes) in the granular model, it is easy to see that the average magnetization m in the spin model is related to the average particle density (t) in the granular model via the sim ple relation, m (t) = 1 2 (t). Thus the result in Eq. (4) for the magnetization indicates that the density will grow to its fully compact value 1 as

$$(t) = 1 \quad \frac{\log b}{\log (bt=t_0)}; \tag{9}$$

at very late times, as observed in experiments on mechanically tapped granular media [3].

O ther one dimensional lattice models, notably the car parking' model, has been previously used to explain the logarithm ic relaxation in granular m aterials [9]. The local rules for the dynam ics of particles in the car parking modelare, how ever, quite di erent from those in our model. In the parking model, the rules for the particle motion are: (i) a new particle can be absorbed only at sites containing a hole at a rate proportional to the num ber of neighboring particles and (ii) a particle evaporates with a small rate (thus leaving behind a hole) if it has one and only one neighboring hole. W hile this model also exhibits an inverse logarithm ic growth of the density, the dynam icalm oves (in particular the fact that the desorption rate is in nitesimally small) are chosen in a som ew hat ad hoc m anner. In contrast, as explained via the mapping detailed above, our model incorporates the basicm in im alm icroscopicm oves of the particles that are observed in the compaction process.

In terms of the motion of the domain walls between "+ " and " " phases, our model can also be considered a new model of two species reaction-di usion in one dimension. We note that in the case = 1, the domain walls di use and annihilate upon contact. This corresponds to the process A + A ! 0 [7]. In the case = 0, we need to distinguish between the two types of dom ain walls + A and + B. Note that by de nition (originating from a spin con guration) the A's and B's always occur alternately. Here both A's and B's di use as before; however when an A and a B meet, they annihilate only if A is to the left of B, otherwise there is hard core repulsion between them . We will show below that this hard core repulsion between the particles is a relevant interaction that changes the late time dynam ics considerably. Recently the relevance of hard core repulsion between particles in reaction-di usion systems have been explored in a number of contexts [10].

## IV . SOM E EXACT RESULTS

To start with we write down two exact relations which are derived directly from the G lauber dynam ics (we shall see later that these exact relations are in fact respected by the IIA approximation).

Let N (t) be the number of domains of either + 'or ' 'spins per unit length. The density of kinks is therefore 2N (t). Let  $P_1$  be the density of the triplets ' + 'per unit length, then clearly  $p_1 = P_1 = N \cdot W$  enote also that the only way in which a domain can be destroyed in an in nitesimal time step is by ipping an isolated + 'spin in a triplet ' + '. This gives

$$\frac{dN}{dt} = P_1:$$
(10)

If P ( $S_i$ ) denotes the probability that the spin at site i takes the value  $S_i$ , the evolution of P ( $S_i$ ) depends only on the rates in Eq. (1) and the three point probability distribution P ( $S_i$  1; $S_i$ ; $S_{i+1}$ ). The evolution of P ( $S_i$ ) is then given by

$$\frac{dP(S_{i})}{dt} = X \\ S_{i+1};S_{i-1} \\ X \\ W(S_{i};S_{i-1};S_{i+1})P(S_{i-1};S_{i};S_{i+1}) \\ W(S_{i};S_{i-1};S_{i+1})P(S_{i-1};S_{i};S_{i+1}) (11) \\ S_{i+1};S_{i-1}$$

Substituting  $S_i = 1$  in the above equation we nd

$$\frac{dP(1)}{dt} = P(1; 1; 1) P(1; 1; 1) P(1; 1; 1)$$
(12)

where we have used the evident left to right symmetries P (1; 1;1) = P (1; 1; 1) and P (1;1; 1) = P (1;1;1). We now observe that P (1; 1;1) + P (1; 1;1) = P (1;1) and P (1;1; 1) + P (1;1; 1) = P (1; 1) = P (1;1) and thus nd

$$\frac{dP(1)}{dt} = P(1; 1;1):$$
(13)

We note that P (1; 1;1) is simply the probability that at a given site the spin is a 'spin and its two neighbors are +' spins, that is to say that their is a + +' defect at the site considered. If we now sum this equation over each site on an interval of unit length on the lattice and recall that R<sub>1</sub> is the density of the + +' triplets per unit length, we obtain

$$\frac{dL_{+}}{dt} = R_1:$$
(14)

where  $L_+$  is the fraction of the +' spins. Note that  $L_+ + L_- = 1$ , where L\_ is the fraction of the ' spins. From the relation Eq. (14) we obtain Eq. (2) for the evolution of the average m agnetization m. Physically it is easy to see the origin of Eq. (14) as, on an average, the fraction of +' spins can decrease only due to the blockage by + +' triplets. Writing Eq. (14) in term  $sofr_1 = R_1 = N$  and the average length of the + ' dom ains  $l_{+} = L_{+} = N$  we obtain

$$\frac{N}{N} = \frac{l_{\pm} + r_1}{l_{\pm}}; \qquad (15)$$

where  $\underline{\mathbf{x}} = d\mathbf{x} = d\mathbf{t}$ . This equation can be integrated starting at some arbitrary time  $t_0$  to give

$$\frac{N(t)}{N(t_0)} = \frac{l_{+}(t_0)}{l_{+}(t)} \exp \left( \begin{array}{c} 2 \\ t \\ t_0 \end{array} \right) \frac{r_1(t^0)}{l_{+}(t^0)} dt^0 \quad : \quad (16)$$

Furtherm ore, if the volume fraction of the + ' phase is,  $L_+$   $(t_0)=-$  , then, using the relation, N (t)= 1=[1 (t)+ 1, (t)] in Eq. (16), we nd,

$$\frac{1}{l_{+}(t)} = \frac{1}{l_{+}} \exp \left( \frac{z_{+}}{t_{0}} \frac{r_{1}(t^{0})}{l_{+}(t^{0})} dt^{0} \right)$$
(17)

clearly showing that the ratio 1 (t)=1, (t) is growing due to the presence of the triplets + +'. Note that the asymmetry between the growth of ' ' and '+' domains is evident due to the presence of the triplet defects '+ +' with density  $R_1$  =  $r_1N$ .

All the results presented above are exact. To derive the late time behavior of the model we rst consider below the IIA.We solve the IIA equations self-consistently and show that the IIA precisely predicts the results mentioned in Section II. Besides we shall argue that in the case where the initial volum e fraction of the + ' domains is small, i.e., 1, correlations do not develop between the domains if no correlations are present in the initial conditions and hence the IIA is exact to leading order in

. At the end we present a very sim ple heuristic argum ent which is also in agreem ent with these results.

#### V.IIA ANALYSIS

In this section we consider the IIA where correlations between neighboring dom ains are neglected. The IIA was used previously for the = 1 case [11] yielding results in agreement, qualitatively as well as quantitatively to a fair degree of accuracy, with the exact results available [6,12]. Let  $P_n$  (t) and  $R_n$  (t) denote respectively the num – ber density of + ' and ` ' dom ains of length n at time t. Note that  $R_1$  (t) is the density pof the triplet + +' as before. Let N (t) =  $_n P_n = _n R_n$  denote the dom ain density of + ' or ` ' spins. A loo the fractions L (t) of + ' and ` ' spins are given by,  $L_+$  (t) =  $_n nP_n$  and L (t) =  $_n nR_n$  with  $L_+$  (t) + L (t) = 1. During an in nitesimal time step t,  $P_n$  (t) evolves as:

$$P_{n} (t + t) = P_{n} (t) tP_{n} (t) tP_{n} (t) 1 \frac{R_{1} (t)}{N (t)} + tP_{n+1} (t) + tP_{n-1} (t) 1 \frac{R_{1} (t)}{N (t)} : (18)$$

The right hand side of the above equation includes the various loss and gain term s. The second and the third term s describe respectively the loss due to the hopping inward and hopping outward of the domain walls at the two ends of a '+' dom ain of size n. An outward hop can occurprovided the neighboring dom ain in the direction of the hop is not an isolated ' ' spin and this is ensured by the prefactor (1  $R_1=N$ ) in the third term . The fourth and the last term describe sim ilarly the corresponding gains. One can similarly write down the evolution equation for the  $R_n$  (t)'s. During an in nitesimal time step t,  $R_n$  (t) evolves as:

$$R_{n}(t+t) = R_{n}(t) \quad tR_{n}(t) \quad tR_{n}(t) \quad 1 \quad \frac{P_{1}(t)}{N(t)}$$

$$2 \ tP_{1}(t) \frac{R_{n}(t)}{N(t)} + \ tR_{n+1}(t)$$

$$+ \ tR_{n-1}(t) \quad 1 \quad \frac{P_{1}(t)}{N(t)}$$

$$+ \frac{P_{1}(t)}{N^{2}(t)} \frac{X^{2}}{I_{i=1}} R_{i}(t)R_{n-i-1}(t); n \quad 2 \quad (19)$$

and

$$R_{1}(t+t) = R_{1}(t) tR_{1}(t) 1 \frac{P_{1}(t)}{N(t)} 2 tP_{1}(t) \frac{R_{1}(t)}{N(t)} + tR_{2}(t): (20)$$

The negative (loss) term s in Eq. (19) for dom ains of size 2 m ay be understood as follows. A dom ain of length n n may be lost by the dom ain wall at either end jum ping inwards with rate 1=2. This term is the second term in Eq. (19) and as there are two dom ain walls we have a factor of 2. A dom ain of length n m ay also be lost by a dom ain wall hopping outwards. This happens with rate 1=2 if the neighboring dom ain is not a triplet ' + '. The third term of Eq. (19) corresponds to this event, the factor  $(1 P_1(t)=N(t))$  is the probability of the absence of a triplet ' + ' as a neighboring dom ain. There is again a factor of 2 coming from the fact that there are two domain walls. However if a neighboring domain is of type ' + ' the outward jum p towards this dom ain occurs with rate 1 (as the central + spin ips with rate 1). The term corresponding to these two events (from the right and left dom ain walls) is the fourth term in Eq. (19). The two rst gain terms come from identical arguments and the last convolution term represents domain coalescence, where a dom ain of length n is form ed with rate 1 from two ' ' dom ains of length i and n i 1 (where 1 i n 2) if they are separated by a ' + triplet. Eq. (20) is obtained in a similar fashion with the exception that the hard core repulsion generates a re ecting boundary condition. Taking the limit t! 0 in the above equations we obtain the IIA equations for the evolution of the dom ain densities

$$\frac{dP_n}{dt} = P_{n+1} + P_{n-1} \quad 2P_n + \frac{R_1}{N} (P_n - P_{n-1}) \quad (21)$$

for all n = 0 (absorbing boundary condition) and

$$\frac{dR_{n}}{dt} = R_{n+1} + R_{n-1} \quad 2R_{n} \quad \frac{P_{1}}{N} (R_{n} + R_{n-1}) + \frac{P_{1}}{N^{2}} R_{1}R_{n-1} R_{1}; \quad n = 2$$

$$\frac{dR_{1}}{dt} = R_{2} \quad R_{1} \quad \frac{P_{1}}{N}R_{1}; \quad (22)$$

It is somewhat convenient to use the normalized variables  $p_n = P_n = N$  and  $r_n = R_n = N$ . The average domain lengths are then given by  $l_{+}(t) = np_n$  and  $l_{-}(t) = P_n = N$ .

 $nr_n$  and the dom ain density N (t) = 1=(l, (t) + 1 (t)). In term s of these normalized variables the IIA equations are given by,

$$\frac{dp_n}{dt} = p_{n+1} + p_{n-1} \quad 2p_n + r_1 (p_n \quad p_{n-1}) + p_1 p_n \quad (23)$$

for all  $1 \text{ with } p_0 = 0$  (absorbing boundary condition) and

$$\frac{dr_{n}}{dt} = r_{n+1} + r_{n-1} + 2r_{n} + p_{n-1} + r_{n-1} +$$

It is easy to check that the normalization condition,  $p_n = r_n = 1$  is satisfied by these two equations. We note that Eq. (23) or more clearly its unnormalized version in Eq. (21), i.e.,  $P_n = P_{n+1} + P_{n-1} - 2P_n + r_1 (P_n - P_{n-1})$ , just represents the motion of a random walker on the positive side of a 1-d lattice with a sink at the origin ( $P_0 = 0$ ) and a time dependent drift term (proportional to  $r_1$ ). To calculate N (t) using Eq. (16), we need to evaluate two quantities from the IIA equations: (i)  $r_1$  (t) =  $R_1$ =N and (ii) 1, (t) = np\_n.

The two IIA equations above are coupled nonlinear equations with in nite number of variables and hence exact solution of Eqs. (23) and (24) are di cult. Our approach will be a combination of a scaling assumption and then rechecking this assumption for self-consistency. Consider rst the  $r_n$  equation, i.e. Eq. (24). On the right hand side, we will rst ignore the di usion term, solve for the rest and show that indeed neglecting the di usion term was justiled in the rst place. This is self-consistency. Ignoring the di usion term, we have the following equation,

$$\frac{dr_n}{dt} = p_1 \qquad \begin{array}{c} & & \\ & X & ^2 \\ & r_i r_{n \ i \ 1} & r_{n \ 1} \end{array} \qquad (25)$$

with the rejecting boundary condition,  $r_2 = r_1$ . It is now easy to see that Eq. (25) admits a scaling solu-(t)  $\exp[n(t)]$ , where -(t) = p(t)(t). tion,  $r_n$  (t) Using the exact relation, dN = dt = pN, we get,  $(t) = (t_0)N(t)=N(t_0)$ . Note that we still do not know what N (t) is. Now let us substitute this solution to estimate the diusion term that had been neglected in the st place. Clearly, the diusion term,  $T_{diff} = r_{n+1} + r_{n-1} - 2r_n - 0$  (<sup>3</sup> (t)), whereas the other term s (for example the left hand side of Eq. (24)) typically scale as O(-(t)) O(p(t)(t)). Thus, in order to be self-consistent in neglecting the di usion term, we <sup>2</sup> (t) N<sup>2</sup> (t). Wew ill see that this need to have  $p_1$  (t) condition is actually satis ed once we derive the expression for N (t). This just means that the di usion terms only contribute to the corrections to the leading scaling behavior.

From the above analysis, we nd to leading order for large t,  $r_1$  (t) = (t<sub>0</sub>)N (t)=N (t<sub>0</sub>), i.e.,  $r_1$  (t)  $r_1$  (t<sub>0</sub>)N (t)=N (t<sub>0</sub>). We now preed to evaluate the other remaining quantity, l<sub>+</sub> (t) =  $np_n$ . For this we now turn to the  $p_n$  equation, Eq. (23). In this equation, we will again rst ignore the drift term  $r_1$  ( $p_n$   $p_{h-1}$ ) solve for the rest and check that indeed the neglect of the drift term was justi ed. Ignoring the drift term, we get,

$$\frac{dp_n}{dt} = p_{n+1} + p_{n-1} + 2p_n + p_1 p_n :$$
 (26)

with the absorbing boundary condition,  $p_0 = 0$ . This equation can be solved exactly. Indeed it also adm its a scaling solution,  $p_n$  (t) = t  $^{1=2}f$  (nt  $^{1=2}$ ), where the scaling function (normalized to unity) is given by, f(x) = $\frac{x}{2}$  exp(  $x^2=4$ ). Now let us estimate the drift term that was neglected. C learly the drift term ,  $r_1 (p_n)$  $p_{1}$  1) 0 (N (t)=t) since n 0 (r<sub>1</sub>=t) N (t) from previous paragraph. The other terms in the Eq. (23) (for example the left hand side of Eq. (23)) is of order, t  $^{3=2}$ at late times. Thus, for self-consistency in neglecting the drift term, we need to show that t  $^{3=2}$ N (t)=t. We will again see that this condition is indeed also satis ed once we obtain the expression for N (t). From this form of  $p_n$  (t), we thus obtain, to leading order  $t^{2} t^{1} t^{1} xf(x) dx$ . Using for large  $t_{r} \downarrow_{t} (t) = np_{n}$  $f(x) = \frac{x}{2} \exp((x^2 = 4))$ , and doing the integral we nally t for large t. nd, l<sub>+</sub> (t)

Using these two results (i)  $r_1(t) = r_1(t_0)N(t)=N(t_0)$ and (ii) 1, (t) = t in the exact equation Eq. (16) and di erentiating with respect to t, we nd a di erential equation for N(t),

$$\frac{d}{dt} \stackrel{P}{=} \frac{r_{1}(t_{0})}{N(t_{0})} N^{2}(t): \qquad (27)$$

Introducing the dim ensionless variable

$$S(t) = \int_{-t}^{t} \frac{T_{0}}{t} \frac{N(t)}{N(t_{0})}$$
 (28)

in Eq. (27) we nd

$$\frac{dS}{dt} = \log(b)\frac{S^2}{t}$$
(29)

where log (b) =  $\frac{p_{-}}{r_1(t_0)^p t_0}$ . Integrating Eq. (29) we nd

$$S(t) = \frac{\log(b)}{\log(bt=t_0)};$$
 (30)

we thus obtain the result

$$\frac{N(t)}{N(t_0)} = \frac{r(t_0)}{t(t_0)} \frac{\log(b)}{\log(bt=t_0)};$$
(31)

Substituting this result in the expression,  $r_1$  (t)  $r_1$  (t<sub>0</sub>)N (t)=N (t<sub>0</sub>), we get

$$r_1 (t) = \frac{p_{-}}{t \log (bt=t_0)}$$
: (32)

Next we use the late time result Eq. (31) in the exact relation Eq. (10) and nd,

$$p_1 = \frac{1}{2t} + \frac{1}{t \log(bt=t_0)};$$
(33)

Let us check the two self-consistency conditions, (a)  $p_1(t) = (t) = N^2(t)$  and (b)  $t^{3=2} = N(t)=t$ . Using the expression for  $p_1$  from Eq. (33) and that of N(t) from Eq. (31), it is immediately evident that indeed these two conditions are satisticed for large t. Thus our whole approach has been completely self-consistent and the IIA results are precisely those mentioned in the abstract. Note also that  $t_0$  must be su ciently large such that both scaling laws,  $r_1(t) = N(t)$  and 1(t) = t start holding for  $t > t_0$ .

#### VI.ZERO VOLUME FRACTION LIM IT

In this section, we show that the IIA results essentially become exact in the zero volume fraction lim it of the + ' phase, i.e., in the lim it ! 0. Suppose we start <sup>n</sup>) and from an initial condition such that,  $r_n(0) = (1)$  $p_n(0) = n_{11}$ . Thism eansthat in the initial condition, the average length of the ' ' dom ains, 1 (0) 1= , whereas  $l_{+}(0) = 1$ . Thus the ' ' domains are typically much larger than the \+' domains, in the limit ! 0. Also initially all the dom ains are completely uncorrelated. So the picture is as follows. We have little droplets of +'phase in a sea of ' ' phase. Besides, one can also com pute the initial density of dom ains of either "+ ' or ' ' types. It is given by, N (0) = 1 = (1) 1 = to leadingorder in .

Now let us consider the time evolution of the system starting from this initial condition. As time increases, the +' domains will certainly grow in size. But a typical +' domain will disappear (via the absorbing boundary condition) much before encountering other +' domains, i.e., before feeling the presence of the constraint due to triplets + +'. The probability of such an event is of order O (). Thus e ectively, the dynam ics of the system will proceed via eating up of the +' domains. Hence, if there is no correlation between domains in the initial condition, the dynam ics is not going to generate correlations between them. This is precisely what happens in the zero tem perature dynam ics of the q state P ottsm odel in 1-d in the limit q ! 1<sup>+</sup> [13,11].

Thus in this lim it, the evolution of the ' ' domains is governed by the exact equation,

which is same as the IIA equation Eq. (24) without the di usion term. Starting from the initial condition,  $r_n(0) = (1 \ ^n)$ , one can solve the above equation for any texactly to leading order in . It turns out that, to leading order in , Eq. (34) admits a solution,  $r_n(t) = (t) [1 \ (t) ]^n \ ^n$ , where  $(t) = \exp[ \begin{array}{c} R_t \\ 0 p_1(t^0) dt^0 ]. Using once again, the exact equation <math>dN = dt = pN$ , we nd,  $(t) = N \ (t) = N \ (0) = 1 = + O \ (^2)$ . Thus we get,

$$r_1(t) = (t) = N(t) + O(2)$$
: (35)

Now, let us consider the evolution of the + 'dom ains. Since a typical + 'dom ain never encounters (to leading order in ) any other + 'dom ain and hence does not feel the constraint due to  $r_1$  's, the e ective dynam ics of a + 'dom ain is that of a single + 'dom ain immersed in a sea of ' 'phase. Let  $P_n$  denote the probability that such a dom ain is of length n. Then, to leading order in ,  $P_n$  's clearly evolve by the simple di usion equation,

$$\frac{dP_{n}}{dt} = P_{n+1} + P_{n-1} \quad 2P_{n};$$
(36)

with the absorbing boundary condition,  $P_0 = 0$ . The normalized conditional probability,  $p_n = P_n = N$  with  $N = P_n$ , then satisfies the equation,

$$\frac{dp_n}{dt} = p_{n+1} + p_{n-1} + 2p_n + p_1 p_n;$$
(37)

same as Eq. (26). This equation has to be solved with the initial condition,  $p_n \ (0) = 1$ . It can be solved exactly. W ithout writing the explicit solution, we just mention the result for  $\ np_n$ . We not that for large t and leading order in ,

$$L_{+}(t) = np_{n} t + 0 ():$$
 (38)

Using the results from Eqs. (35) and (38), i.e., (i)  $r_1(t) = N(t) + O(^2)$  and (ii) 1, (t) t + O() in the exact equation Eq. (16), we once again recover all the IIA results of the previous section, with  $b = \frac{1}{2} = 1$ .

Hence IIA becomes exact in the ! 0 limit. This is not surprising as the dynamics in this limit does not generate correlations if there are none in the initial condition.

#### VII. OTHER VOLUME FRACTIONS

For nite initial volume fraction of the 4 ' phase, the IIA can not be exact since the di usion of kinks correlate the dom ains as time progresses, even if the dom ains had no correlations to start with. However, the volume fraction of the "+" phase decreases monotonically with time according to the exact equation Eq. (14). Thus at very late times when the volume fraction  $L_+$  (t) is very small, the e ective xed point picture of the system is very similar to the ! 0 limit picture, i.e., sm all + ' dom ains immersed in the sea of ' ' dom ains. The only di erence is that the big ` ' dom ains m ay now be correlated. However, it is very likely (though we can not prove this rigorously) that the correlations between dom ains are very sm all at very late tim es and therefore the IIA results ((1)-(4) in the abstract) become asymptotically exact. The num erical results reported in [5] appear to con m this fact.

A ctually if this late time xed point picture is correct, then one can derive all the results from a very simple heuristic argument, presented below.

#### VIII.HEURISTIC APPROACH

The heuristic picture is as follows. Due to the facts that +' dom ains can grow only by di usion and the +' dom ains grow by di usion and coalescence, one expects that at late times the ratio  $\downarrow_{+}(t)=1$  (t) ! 0; this is also clear from Eq. (14). Therefore at late times one expects to nd +' dom ains sandwiched between much larger +' dom ains. Consequently the +' dom ains, to a rst approximation, never encounter the +' + ' triplets at late times and hence di use freely and annihilate via the disappearance of +' + ' triplets. Therefore the length of a + ' dom ain behaves as a di usion with a killing boundary condition at the origin. Solving the discrete di usion equation corresponding to this picture (see Eq. (37)), one nds that at late times

If one now views the system at the length scale of the 'domains l(t)  $l_{+}$  (t), one sees long stretches of 'domains occasionally interrupted by 'domains which

are now shrunk to a single point when viewed from the length scale of the ' 'dom ains. The rate of occurrence of these points per unit length (t) is clearly proportional to N (t), i.e., the kink density. Note that  $r_n$  is simply the conditional probability: G iven that a ' 'dom ain has occurred, what is the probability that it is of length n. Now if one assumes that these punctual ' 'dom ains (or the points) are distributed random ly, one nds that  $r_n$  is simply given by the geom etric distribution,  $r_n = (t) (1 (t))^{n-1}$  where (t) = dN (t) for some constant c. We therefore nd

$$r_1$$
 (t)  $dN$  (t); (40)

for late times. Furthermore, if we denote by  $t_0$  a large time after which this picture becomes valid we may write

$$r_{1}$$
 (t)  $r_{1}$  (t<sub>0</sub>)  $\frac{N(t)}{N(t_{0})}$ : (41)

These two expressions for  $r_1$  and  $l_{+}$  (t) once again are same as obtained by a more careful analysis of the IIA equations and when substituted in Eq. (16), they give the same IIA results once again. Thus the basic assum ption of this heuristic picture is the +' domains occur random ly, which seems like an accurate description at late tim es.

## IX . C O N C LU S IO N S

In this paper we have presented detailed analytical studies on a simple one dimensional kinetically constrained Ising model which was introduced in Ref. [5]. The kinetic constraints in this model are local and dynam ically generated. The e ect of these constraints was shown to slow down the dynam ics rather dram m atically. W e have shown that the average magnetization in this model decays extremely slowly with time in an inverse logarithm ic fashion to its nal saturation value. This kind of inverse logarithm ic law was observed in the behavior of the density of granularm aterial in experim ents on granular compaction [3] and was also seen in num erical simulations of various lattice based and 'tetris' like m odels [14]. There have been som e theoretical argum ents proposing various mechanisms responsible for this slow compaction [15,9]. These include the free volum e argument [15] and the argument based on an analogy with car parking models with the som ewhat ad hoc assum ption that the cars depark' from a lane at an in nitesim al rate [9]. In contrast, in this paper we have m apped our kinetic Ising model directly to a lattice model of granular compaction which incorporates the basic m in im alm icroscopic moves in the compaction process. The average m agnetization m (t) in the Ising m odel, via this m apping, gets related to the density of com paction (t) in the granular model as (t) = (1 m)=2. Hence, besides having nontrivial behavior and yet analytically solvable, our toy m odel of granular com paction correctly reproduces the inverse logarithm ic tim e dependence seen in the experim ents [3] and thereby proposes a new and entirely di erent m echanism for this slow com paction, quite di erent from the previousm odels such as the carparking m odels. It is also interesting to note a study of com paction in the Tetris m odel [16] shows that at late tim es the activity of the system, leading to com paction, occurs at boundaries between dom ains which can be identied in the system. The im age of com paction as a kinetically hindered coarsening process thus appears to be quite robust.

From a som ew hat broader perspective, our work addresses a general question: what is the e ect of kinetically generated disorders on the coarsening dynamics in domain growth problems? In the present work we have studied a speci c type of kinetic disorder, namely a dynam ically generated local magnetic eld. This eld acts locally on the topological defects responsible for the coarsening process (in this case simple domain walls). Our study suggests that such kinetic disorders, while slowing down the dynamics drastically, do not altogether inhibit the coarsening process as found in other constrained kinetic Ising models [2]. The domain growth problem s are rather common and occur in various physical system s [4]. Our work, therefore, opens up the possibility of studying the slowing down in coarsening dynam ics due to kinetic disorders in m any of these system s. For example, it would be interesting to study the e ect of dynam ically generated local elds in higher dim ensionional Ising models, in O (n) vector spin models and in liquid crystals, to mention a few. It is possible to have other types of local kinetic disorders than the one studied here and it would also be interesting to study their e ect in coarsening system s.

W e thank P.G rassberger for his earlier collaboration in this study, and also M.Barm a for useful discussions. W e acknowledge interesting exchanges with G.O dor, H. H inrichsen and G.Schutz.

- J.P. Bouchaud, L.F. Cugliandolo, J. Kurchan, and M. Mezard, in Spin Glasses and Random Field, edited by A.P.Young (World Scientic, Singapore, 1998).
- [2] G. H. Fredrickson, Ann. Rev. Phys. Chem. 39, 149 (1988);
  J. Jackle and S. Eisinger, Z. Phys. B 84, 115 (1991); JD.
  Shore, M. Holzer, and J.P. Sethna, Phys. Rev. B 46, 11376 (1992); J. Kisker, H. Reiger, and H. Schreckenberg, J. Phys. A 27, L853 (1994); F. Ritort, Phys. Rev. Lett. 75, 1190 (1995); M E J. Newman and C. Moore, Phys. Rev. E 60, 5068 (1999).
- [3] E R. Nowak, J.B. Knight, E. Ben-Naim, H.M. Jaeger, and S.R. Nagel, Phys. Rev. E 57, 1971 (1998)
- [4] A J. Bray, Adv. Phys. 43, 357 (1994).
- [5] SN.Majum dar, D.S. Dean and P.G rassberger, Phys. Rev.Lett. 86, 2301 (2001).
- [6] R.J.G lauber, J.M ath. Phys. 4, 294 (1963).
- [7] A J.Bray, J.Phys.A 22, L67 (1990); J.G.Am ar and F. Fam ily, Phys. Rev. A 41, 3258 (1990); S.N.Majum dar and D.A.Huse, Phys. Rev. E 52, 270 (1995).
- [8] P. Sollich and M R. Evans, Phys. Rev. Lett. 83, 3238 (1999).
- P.L.K rapivsky and E.Ben-Naim, J.Chem. Phys. 100
   6778, (1994); X.Jin, G.Tarjus and J.Talbot, J.Phys.
   A 27, L195 (1994); E.Ben-Naim, J.B.Knight and E.R.
   Nowak, Physica D 123, 380 (1998).
- [10] S.Kwon, J.Lee and H.Park, Phys. Rev. Lett. 85, 1682
   (2000); G.Odor and N.Menyhard, Phys. Rev. E 61, 6404 (2000); F.van W ijland, cond-m at/0010491.
- [11] P L.K rapivsky and E.Ben-Naim, Phys.Rev.E 56, 3788 (1997).
- [12] B.Derrida and R.Zeitak, Phys. Rev. E 54, 2513 (1996).
- [13] C.Sire and SN.M ajum dar, Phys.Rev.E 52,244 (1995).
- [14] E. Caglioti, V. Loreto, H.J.Herrm ann and M. Nicodemi, Phys.Rev.Lett.79, 1575 (1997); M. Nicodemi, J.Phys.I 17, 1535 (1997); M. Piccioni, M. Nicodemi and S.Galam, Phys.Rev.E 59, 3858 (1999).
- [15] J.B. Knight et al, Phys. Rev. E 51, 3957 (1995); T. Boutreux and P.G. de Gennes, Physica A 244, 1, (1997).
- [16] A. Baldassarri, S. Krishnam urthy, V. Loreto and S. Roux, Phys. Rev. E 87, 224301 (2001)

Figure Captions

Fig. 1. Picture of rolling o (left dom ain) and tapping up (right dom ain) in the granular interpretation of the spin model, the solid squares represent particles.

