

Comment on “Freezing by heating in a driven mesoscopic system”

Leo Radzihovsky and Noel A. Clark

Department of Physics, University of Colorado, Boulder, CO 80309

(Dated: March 22, 2022)

In a recent Letter Helbing et al.[1] found a fascinating fluctuation-induced ordering, which they dubbed “freezing by heating”, in a nonequilibrium model system of particles confined to a 2d narrow strip and driven to move in opposite directions: counter flowing chains of particles jam up and crystallize when sufficiently large noise is introduced into the system. In this Comment we point out that this kind of fluctuation-induced ordering is *not* an intrinsically *nonequilibrium* effect, and that it can be expected to occur in a variety of *equilibrium* systems, with reentrant order appearing as a function of temperature. We argue that this is a consequence of a familiar Helfrich-like interaction that had been used to understand a variety of phenomena ranging from the lamellar phase of fluctuating membranes to magnetization curves of a vortex liquid state of type II superconductors.

We demonstrate the mechanism behind reentrant melting with an equilibrium model of 2d colloids subjected to a 1d periodic potential, that received considerable attention recently due to the experimental discovery of *isothermal* melting that is reentrant as a function of strength of the periodic potential (produced by two interfering laser beams) [2], the so-called reentrant “laser-induced melting” (RLIM).[3]

Since at all T the orientational symmetry and translational symmetry along y are *explicitly* broken by the 1d periodic potential with troughs running along x and spacing $a = 2\pi/q$, the only remaining feature distinguishing a crystal from the modulated liquid is the spontaneous translational order along the troughs, characterized by quasi-long ranged u_x phonon correlations and a finite shear modulus μ_{xy} .

The mechanism behind RLIM can be understood as follows[4]. Just above melting temperature $T_m(V_q)$, the dominant effect of increasing a *weak* periodic potential V_q is to suppress thermal fluctuations, thereby leading to LIF[3]. In contrast, for *large* V_q , the suppression of the out-of-trough u_y fluctuations has the dominant effect of

reducing the inter-row interactions, thereby decoupling the particles in the neighboring rows, reducing the effective shear modulus $\mu_{xy}(T, V_q)$, and leading to RLIM.[2]

Since it is clearly the ratio T/V_q that determines the extent of the fluctuation-driven inter-row interaction, it follows that at large *fixed* V_q increasing T will increase the inter-row interaction and the modulus $\mu_{xy}(T, V_q)$. Because T_m is typically a monotonically increasing function of the shear modulus[4] thermal enhancement of shear modulus implies an equilibrium analog of the “freezing by heating” transition discovered by Helbing, et al.[1].

Above qualitative argument can be formalized by an explicit calculation. Although T dependence of both the bulk and the shear moduli maybe important and can be studied in full detail[4], in order to illustrate the effect, it is sufficient to focus on the effective shear modulus $\mu_{xy}(T, V_q)$. Clearly this modulus is proportional to the interaction potential between nearest particles in adjacent rows, typically of Debye-screened form $U_D \approx e^{-(a+u_y)/\xi}$ (though only its short-range nature is important), where ξ is the screening length. Since phonon fluctuations u_y transverse to the troughs are “optical” (i.e., “massive”, as opposed to the acoustic u_x phonon) we can average over fast u_y fluctuations obtaining an effective shear modulus $\mu_{xy}(T, V_q) \approx \mu_0 e^{\langle u_y^2 \rangle / 2\xi^2} \approx \mu_0 e^{T/V_q}$ that clearly increases with T . [5] We can then compute $\langle u_x^2 \rangle \approx T/\mu_{xy}(T, V_q) \approx T e^{-T/V_q}/\mu_0$ and use Lindemann criterion $\langle u_x^2 \rangle = d^2$ to estimate $T_m(V_q)$. This clearly gives two roots for T_m , with the lower corresponding to the conventional melting and the upper to the freezing by heating (see figure). At sufficiently high T , we expect above elastic calculation to become inaccurate and for the fluctuation enhancement of $\mu_{xy}(T, V_q)$ to saturate, leading to re-melting into a modulated liquid.

While extending above analysis to the nonequilibrium case studied by Helbing, et al.[1] remains an interesting and challenging problem, we believe that fluctuation-driven reentrant freezing mechanism described above is also at play in their system.

Support by the NSF MRSEC DMR-0213918 and the Packard Foundation (LR).




FIG. 1: Schematic phase diagram of colloidal system in the inset showing the mechanism for fluctuation driven freezing.

- [1] D. Helbing, *et al.*, *Phys. Rev. Lett.* **84**, 1240 (2000).
- [2] Q.-H. Wei *et al.*, *Phys. Rev. Lett.* **81**, 2606 (1998).
- [3] A. Chowdhury *et al.*, *Phys. Rev. Lett.* **55**, 833 (1985).
- [4] L. Radzihovsky, *et al.*, *Phys. Rev. E* **63**, 31503 (2001).
- [5] Qualitative increase of $\mu_{xy}(T)$ with T holds beyond this harmonic approximation.