

Comment on "Stability of Small Carbon-Nitride Heterofullerenes"

Guangyu Sun¹, Ruihua Xie², and Marc C. Nicklaus¹

¹Laboratory of Medicinal Chemistry, NCI-Frederick, Frederick, MD 21702

²National Institute of Standards and Technology, Gaithersburg, MD 20899-8423

(April 14, 2024)

In a recent Letter [1], Schultz et al. proposed the existence of heterofullerenes C_n-xN_x ($40 \leq n \leq 50$) based on the pattern of the peaks in the mass spectrum of carbon-nitride nanostructures produced by the arc-discharge technique. Using PM3 semiempirical calculations, they predicted the most stable structures of the N-substituted fullerenes, $C_{33}N_9$ and $C_{44}N_6$, to have C_3 symmetry. While some level of symmetry seems to maximize stability, the highest symmetry often sacrifices it, as in the case of fullerene C_{84} [2]. Using PM3 and density functional theory (DFT), we argue that a highly systematic investigation is needed to determine the most stable structures of both pristine and N-substituted fullerenes. We demonstrate this explicitly for two particular cases, C_{40} and $C_{44}N_6$, for which the authors of Ref. [1] only considered four and two isomers, respectively.

Mathematically, there are 40 possible isomers for C_{40} [3]. We optimized the geometries of the 37 isomers with non-vanishing HOMO-LUMO gaps using the PM3 Hamiltonian and the B3LYP functional with STO-3G, 3-21G and 6-31G* basis sets as implemented in Gaussian 98 [4]. The closed-shell formalism was used and the topological symmetry was maintained throughout. Isomers 1, 30 and 40 (note: the isomer index follows Ref. [3]) have zero HOMO-LUMO gaps at the Huckel level of theory. Geometric convergence could not be achieved for these cases at the PM3 level and no further calculation was performed.

The most stable isomer of C_{40} is predicted to be isomer 38 (D_2) by all levels of theory. A analysis of the heats of formation (not shown) indicated that isomer 38 is not among the four isomers studied in Ref. [1], although the authors did not identify the isomers. Isomers with low relative energies include isomer 39 (D_{5d} , 10.9 kcal/mol) and isomer 31 (C_3 , 16.1 kcal/mol), see Fig. 1. All other isomers have energies higher than 20 kcal/mol. These results are in agreement with earlier predictions [5,6]. The stabilities from different levels of theory agree with each other.

In Ref. [1], Schultz et al. showed an isomer with C_3 symmetry as the most stable structure for C_{50} substituted with six nitrogen atoms (Fig. 4b in Ref. [1]). Here we present a complete search of the N_6 -substituted isomers within the frame of $C_{50}270$ (D_3). Dividing the six

N atoms into two groups of three atoms and substituting different groups of carbon atoms in $C_{50}270$ gives 64 possible structures. We first performed calculations at PM3 and B3LYP/STO-3G levels, which gave similar relative energies. Three structures have relative energies lower than 16 kcal/mol, whereas all others have relative energies higher than 40 kcal/mol. Further calculations at the B3LYP/6-31G* level gave relative energies as 0.00 (D_3), 6.78 (C_3) and 9.63 (D_3) kcal/mol, confirming results from lower levels. The HOMO-LUMO gaps, 2.70 eV, 2.85 eV and 2.58 eV, respectively, are large by DFT standards, indicating their kinetic stability. Fig. 2 shows the most stable structure of $C_{50}270$ with N_6 -substitution having D_3 symmetry.

In summary, quantum chemical calculations at both DFT and PM3 levels indicate that either a complete survey or a more rational selection scheme is needed to determine the ground state structures of fullerenes.

-
- [1] D. Schultz, R. D. Roppa, F. A. Alvarez, and M. C. dos Santos, Phys. Rev. Lett. 90, 015501 (2003).
 - [2] G. Sun and M. Kertesz, J. Phys. Chem. A 105, 5212 (2001).
 - [3] P. W. Fowler and D. E. Manolopoulos, An Atlas of Fullerenes (Clarendon Press, Oxford, 1995).
 - [4] M. J. Frisch et al., Gaussian 98, Revision A11.3, Gaussian Inc., Pittsburgh, PA, 2002.
 - [5] B. L. Zhang et al., J. Chem. Phys. 97, 5007 (1992).
 - [6] E. A. Bertazzi et al., Phys. Chem. Chem. Phys. 1, 2913 (1999).

Caption of Figures

Fig. 1: Relative energy of isomers of fullerene C_{40} . The isomer index follows Ref. [3].

Fig. 2: The most stable structure of $C_{50}270$ with N_6 -substitution has D_3 symmetry.

Fig 1/PRL/Sun et al.




Fig 2/PRL/Sun et al.

