

An example of dissipative quantum system : finite differences for complex Ginibre ensemble

Maciej M. Duras

Institute of Physics, Cracow University of Technology, ulica Podchorążych 1, 30-084
Cracow, Poland

E-mail: m.duras@riad.usk.pk.edu.pl

"Quantum dynamics in terms of phase-space distributions"; May 22nd, 2000 to May 26th, 2000; Max Planck Institute for the Physics of Complex Systems; Dresden, Germany (2000).

AD 2000 May 16

1 Abstract

The Ginibre ensemble of complex random Hamiltonian matrices H is considered. Each quantum system described by H is a dissipative system and the eigenenergies Z_i of the Hamiltonian are complex-valued random variables. For generic N -dimensional Ginibre ensemble analytical formula for distribution of second difference 1Z_i of complex eigenenergies is presented. The distributions of real and imaginary parts of 1Z_i and also of its modulus and phase are provided for $N = 3$. The results are considered in view of Wigner and Dyson's electrostatic analogy. General law of homogenization of eigenenergies for different random matrix ensembles is formulated.

2 Introduction

Random Matrix Theory RMT studies quantum Hamiltonian operators H for which their matrix elements H_{ij} are random variables [1, 2, 3]. There were studied among others the following Gaussian Random Matrix ensembles GRME: orthogonal GOE, unitary GUE, symplectic GSE, as well as circular ensembles: orthogonal COE, unitary CUE, and symplectic CSE. The choice of ensemble is based on quantum symmetries ascribed to H . It was Eugene Wigner who firstly dealt with eigenenergy level repulsion phenomenon [1, 2, 3]. RMT is applicable now in many branches of physics: nuclear physics (slow neutron resonances, highly excited complex nuclei), condensed phase physics (nematic particles, random Ising model [spin glasses]), quantum chaos (quantum billiards, quantum dots), disordered mesoscopic systems (transport phenomena). Jean Ginibre considered another example of GRME dropping the assumption of hermiticity of Hamiltonians thus defining generic complex valued H [1, 2, 4, 5]. Hence, H belong to general linear Lie group $GL(N, \mathbb{C})$, where N is dimension and $\mathbb{C}$ is complex numbers field. Therefore, the eigenenergies Z_i of quantum system ascribed

to Ginibre ensemble are complex valued random variables. Jean Ginibre postulated the following joint probability density function of random vector of complex eigenvalues $Z_1; \dots; Z_N$ for $N \times N$ Hermitian matrices [1, 2, 4, 5]:

$$P(Z_1; \dots; Z_N) = \frac{1}{N!} \prod_{i < j}^N |z_i - z_j|^2 \exp\left(-\sum_{j=1}^N |z_j|^2\right); \quad (1)$$

where z_i are complex-valued sample points ($z_i \in \mathbb{C}$).

We emphasise here Wigner and Dyson's electrostatic analogy. A Coulomb gas of N unit charges moving on complex plane (Gauss's plane) $\mathbb{C}$ is considered. The vectors of positions of charges are z_i and potential energy of the system is:

$$U(Z_1; \dots; Z_N) = \sum_{i < j}^N \ln |z_i - z_j| + \frac{1}{2} \sum_{i=1}^N |z_i|^2; \quad (2)$$

If gas is in thermodynamic equilibrium at temperature $T = \frac{1}{2k_B}$ ($\beta = \frac{1}{k_B T} = 2$, k_B is Boltzmann's constant), then probability density function of vectors of positions is $P(Z_1; \dots; Z_N)$ Eq. (1). Therefore, complex eigenenergies Z_i of quantum system are analogous to vectors of positions of charges of Coulomb gas. Moreover, complex-valued spacings 1Z_i of complex eigenenergies of quantum system:

$${}^1Z_i = Z_{i+1} - Z_i; i = 1; \dots; (N-1); \quad (3)$$

are analogous to vectors of relative positions of electric charges. Finally, complex-valued second differences 2Z_i of complex eigenenergies:

$${}^2Z_i = Z_{i+2} - 2Z_{i+1} + Z_i; i = 1; \dots; (N-2); \quad (4)$$

are analogous to vectors of relative positions of vectors of relative positions of electric charges. The 2Z_i have their real parts $\text{Re } {}^2Z_i$, and imaginary parts $\text{Im } {}^2Z_i$, as well as radii (moduli) $|{}^2Z_i|$, and main arguments (angles) $\text{Arg } {}^2Z_i$. 2Z_i are extensions of real valued second differences:

$${}^2E_i = E_{i+2} - 2E_{i+1} + E_i; i = 1; \dots; (N-2); \quad (5)$$

of adjacent ordered increasingly real valued energies E_i defined for GOE, GUE, GSE, and Poisson ensemble PE (where Poisson ensemble is composed of uncorrelated randomly distributed eigenenergies) [6, 7, 8, 9].

3 Second Difference Distributions

Let us define following random variables for $N = 3$ dimensional Ginibre ensemble:

$$Y_1 = {}^2Z_1; A_1 = \text{Re } Y_1; B_1 = \text{Im } Y_1; \quad (6)$$

$$R_1 = \sqrt{Y_1}; \quad r_1 = \text{Arg} Y_1; \quad (7)$$

Their distributions for $N = 3$ dimensional Ginibre ensemble read [10]:

$$\begin{aligned} f_{Y_1}(Y_1) &= f_{(A_1, B_1)}(a_1; b_1) = \\ &= \frac{1}{576} [(a_1^2 + b_1^2)^2 + 24] \exp\left(-\frac{1}{6}(a_1^2 + b_1^2)\right); \end{aligned} \quad (8)$$

$$f_{A_1}(a_1) = \frac{1}{576} (a_1^4 + 6a_1^2 + 51) \exp\left(-\frac{1}{6}a_1^2\right); \quad (9)$$

$$f_{B_1}(b_1) = \frac{1}{576} (b_1^4 + 6b_1^2 + 51) \exp\left(-\frac{1}{6}b_1^2\right); \quad (10)$$

$$\begin{aligned} f_{R_1}(r_1) &= \\ &= \frac{1}{288} r_1 (r_1^4 + 24) \exp\left(-\frac{1}{6}r_1^2\right); \\ f_{r_1}(\varphi) &= \frac{1}{2\pi}; \quad \varphi \in [0; 2\pi]; \end{aligned} \quad (11)$$

For the generic N -dimensional Ginibre ensemble we study complex-valued random variable of second difference: $W_1 = Z_1^2$. Its distribution is $P_3(w_1)$ [10]:

$$\begin{aligned} P_3(w_1) &= \\ &= \sum_{j_1=0}^{N-1} \sum_{j_2=0}^{N-1} \sum_{j_3=0}^{N-1} \frac{1}{j_1! j_2! j_3!} I_{j_1 j_2 j_3}(w_1); \end{aligned} \quad (12)$$

$$\begin{aligned} I_{j_1 j_2 j_3}(w_1) &= \\ &= 2^{-2j_2} \frac{\varrho^{j_1 + j_2 + j_3}}{\varrho^{j_1-1} \varrho^{j_2-2} \varrho^{j_3-3}} F(w_1; -1; -2; -3) j_{i=0}; \end{aligned} \quad (13)$$

$$\begin{aligned} F(w_1; -1; -2; -3) &= \\ &= A(-1; -2; -3) \exp[B(-1; -2; -3) w_1^2]; \end{aligned} \quad (14)$$

$$\begin{aligned} A(-1; -2; -3) &= \\ &= \frac{(2\pi)^2}{(\Gamma(-1 + \frac{5}{4}) \Gamma(-2 + \frac{5}{4}) \Gamma(-3 + \frac{5}{4})) (\Gamma(-1))^2}; \end{aligned} \quad (15)$$

$$\begin{aligned} B(-1; -2; -3) &= \\ &= (\Gamma(-1) - 1) \frac{2\Gamma(-1) \Gamma(-2) \Gamma(-3) + \frac{1}{2}}{2\Gamma(-1) + \Gamma(-2) + \Gamma(-3) - \frac{9}{2}}; \end{aligned} \quad (16)$$

4 Conclusions

In order to compare our results for second differences for different ensembles we define dimensionless second differences:

$$C = \frac{^2E_1}{\langle S \rangle}; \quad (17)$$

$$X_1 = \frac{A_1}{\langle R_1 \rangle}; \quad (18)$$

where $\langle S \rangle$ are the mean values of spacings for GOE (3) ($\beta = 1$), for GUE (3) ($\beta = 2$), for GSE (3) ($\beta = 4$), for PE ($\beta = 0$) [6, 7, 8, 9], and $\langle R_1 \rangle$ is mean value of radius R_1 for $N = 3$ dimensional Ginibre ensemble [10].

We formulate homogenization law [6, 7, 8, 9, 10]: Eigenenergies for Gaussian ensembles, for Poisson ensemble, and for Ginibre ensemble tend to be homogeneously distributed. The second differences' distributions assume global maxima at origin for above ensembles. For Coulomb gas the vectors of relative positions of vectors of relative positions of charges statistically tend to be zero. It can be called stabilisation of structure of system of electric charges.

5 Acknowledgements

It is my pleasure to most deeply thank Professor Jakub Zakrzewski for formulating the problem.

References

- [1] Haake F 1990 Quantum Signatures of Chaos (Berlin Heidelberg New York: Springer-Verlag) Chapters 1, 3, 4, 8 pp 1-11, 33-77, 202-213
- [2] Guhr T, Muller-Groeling A and Weidenmüller H A 1998 Phys. Rept. 299 189-425
- [3] Mehta M L 1990a Random matrices (Boston: Academic Press) Chapters 1, 2, 9 pp 1-54, 182-193
- [4] Ginibre J 1965 J. Math. Phys. 6 440-449
- [5] Mehta M L 1990b Random matrices (Boston: Academic Press) Chapter 15 pp 294-310
- [6] Duras M M and Sokalski K 1996a Phys. Rev. E 54 3142-3148
- [7] Duras M M 1996b Finite difference and finite element distributions in statistical theory of energy levels in quantum systems (PhD thesis, Jagellonian University, Cracow, July 1996)

- [8] Duras M. M. and Sokalski K. 1999a Physica D 125 260-274
- [9] Duras M. M. 1999b Proceedings of the Sixth International Conference on Squeezed States and Uncertainty Relations, 24 May-29 May 1999, Naples, Italy (Greenbelt, Maryland: NASA) at press
- [10] Duras M. M. 2000 J. Opt. B: Quantum Semiclass. Opt. 2 1-5