

A numerical method for generation of quantum noise and solution of generalized c -number quantum Langevin equation

Dhruba Banerjee

Indian Association for the Cultivation of Science, Jadavpur, Kolkata 700 032, India

Bidhan Chandra Bag

Department of Chemistry, Visva Bharati, Shantiniketan 731 235, India

Suman Kumar Banik

*Indian Association for the Cultivation of Science, Jadavpur, Kolkata 700 032, India**

Deb Shankar Ray[†]

Indian Association for the Cultivation of Science, Jadavpur, Kolkata 700 032, India

(Dated: January 6, 2021)

Based on a coherent state representation of noise operator and an ensemble averaging procedure we have recently developed [Phys. Rev. E **65**, 021109 (2002); *ibid.* 051106 (2002)] a scheme for quantum Brownian motion to derive the equations for time evolution of *true* probability distribution functions in c -number phase space. We extend the treatment to develop a numerical method for generation of c -number noise with arbitrary correlation and strength at any temperature, along with the solution of the associated generalized quantum Langevin equation. The method is illustrated with the help of a calculation of quantum mean first passage time in a cubic potential to demonstrate quantum Kramers turnover and quantum Arrhenius plot.

PACS numbers: 05.40-a, 82.20.-w

I. INTRODUCTION

A system coupled to its environment is the standard paradigm for quantum theory of Brownian motion^{1,2,3,4,5,6,7,8,9,10,11,12}. Its overwhelming success in the treatment of various phenomena in quantum optics, transport processes in Josephson junction, coherence effects and macroscopic quantum tunneling in condensed matter physics, electron transfer in large molecules, thermal activation processes in chemical reactions is now well known and forms a large body of current literature. While the early development of quantum optics initiated in the sixties and seventies was based on density operator, semigroup, noise operator or master equation methods primarily within weak-coupling and Markov approximations, path integral approach to quantum Brownian motion attracted wide attention in the early eighties. Although this development had widened the scope of condensed matter and chemical physics significantly, so far as the large coupling between the system and the heat bath and large correlation times of the noise processes are concerned several problems still need to be addressed. *First*, a search for quantum analogue of Kramers' equation for a nonlinear system which describes quantum Brownian motion in phase space had remained elusive and at best resulted in equation of motion which contains higher (than second) derivatives of probability distribution functions^{2,13} whose positive definiteness is never guaranteed. As a result these quasi-probability distribution functions often become singular or negative in the full quantum domain. *Second*, although long correlation times and large coupling constants are treated non-perturbatively formally in an exact manner by path integrals, their analytic evaluation is generally completed within semi-classical schemes^{1,4,5,11}. This results in situations where the theory fails to retain its validity in the vacuum limit. *Third*, although the numerical techniques based on the path integral Monte Carlo are very successful in the treatment of equilibrium properties it is often difficult to implement in a dynamical scheme because of the well-known sign problem associated with varying phases in a quantum evolution over paths due to the oscillating nature of the real time propagator^{6,7,10,12}. *Fourth*, the treatment of non-Markovian dynamics within a quantum formulation as retardation effect in a memory functional is very difficult, if not impossible, to handle even numerically^{10,12}.

Keeping in view of these difficulties it is worthwhile to ask how to extend classical theory of Brownian motion to quantum domain for arbitrary friction and temperature. Based on a coherent state representation of the noise operator and an ensemble averaging procedure we have recently^{14,15,16,17} developed a scheme for quantum Brownian motion in terms of a c -number generalized Langevin equation. The approach allows us to use classical methods of non-Markovian dynamics to derive exact generalized quantum Kramers' equation and other relevant quantum analogues of classical diffusion, Fokker-Planck and Smoluchowski equations. The object of the present analysis is to extend the treatment (*i*) to develop a numerical method for generation of quantum noise as classical c -numbers (*ii*) and to

solve the associated generalized quantum Langevin equation. The numerical method valid for arbitrary temperature, damping strength and noise correlation, is exemplified by an application to calculation of quantum mean first passage time in a cubic potential. The approach is classical-looking in form, and is independent of path integral quantum Monte Carlo techniques and takes care of quantum effect order by order with simplicity and accuracy to a high degree.

The outlay of the paper is as follows: We start with a brief review of c -number representation of quantum noise and the associated quantum Langevin equation, as formulated recently by us, in Sec.II. This is followed by a numerical method for generation of quantum noise and solution of the Langevin equation in Sec.III. In Sec.IV we illustrate the procedure by calculating quantum mean first passage time in a cubic potential with an emphasis on turn-over features and Arrhenius plot in thermal activated processes assisted by tunneling.

II. A COHERENT STATE REPRESENTATION OF QUANTUM NOISE AND QUANTUM LANGEVIN EQUATION IN C-NUMBERS

We consider a particle of unit mass coupled to a medium comprised of a set of harmonic oscillators with frequency ω_j . This is described by the following Hamiltonian:

$$\hat{H} = \frac{\hat{p}^2}{2} + V(\hat{x}) + \sum_j \frac{\hat{p}_j^2}{2} + \frac{1}{2} \kappa_j (\hat{q}_j - \hat{x})^2 \quad (1)$$

Here \hat{x} and \hat{p} are co-ordinate and momentum operators of the particle and the set $\{\hat{q}_j, \hat{p}_j\}$ is the set of co-ordinate and momentum operators for the reservoir oscillators coupled linearly to the system through their coupling coefficients κ_j . The potential $V(\hat{x})$ is due to the external force field for the Brownian particle. The co-ordinate and momentum operators follow the usual commutation relations $[\hat{x}, \hat{p}] = i\hbar$ and $[\hat{q}_j, \hat{p}_j] = i\hbar\delta_{ij}$. Note that in writing down the Hamiltonian no rotating wave approximation has been used.

Eliminating the reservoir degrees of freedom in the usual way^{2,31,32} we obtain the operator Langevin equation for the particle,

$$\ddot{\hat{x}}(t) + \int_0^t dt' \gamma(t-t') \dot{\hat{x}}(t') + V'(\hat{x}) = \hat{F}(t) \quad , \quad (2)$$

where the noise operator $\hat{F}(t)$ and the memory kernel $\gamma(t)$ are given by

$$\hat{F}(t) = \sum_j \left[\{\hat{q}_j(0) - \hat{x}(0)\} \kappa_j \cos \omega_j t + \hat{p}_j(0) \kappa_j^{1/2} \sin \omega_j t \right] \quad (3)$$

and

$$\gamma(t) = \sum_j \kappa_j \cos \omega_j t \quad , \quad (4)$$

with $\kappa_j = \omega_j^2$ (masses have been assumed to be unity).

The Eq.(2) is an exact quantized operator Langevin equation² for which the noise properties of $\hat{F}(t)$ can be defined using a suitable initial canonical distribution of the bath co-ordinates and momenta. Our aim here is to replace it by an equivalent QGLE in c -numbers. To achieve this and to address the problem of quantum non-Markovian dynamics in terms of a *true probabilistic description* we proceed in two steps^{14,15,16,17}. We *first* carry out the *quantum mechanical average* of Eq.(2)

$$\langle \ddot{\hat{x}}(t) \rangle + \int_0^t dt' \gamma(t-t') \langle \dot{\hat{x}}(t') \rangle + \langle V'(\hat{x}) \rangle = \langle \hat{F}(t) \rangle \quad (5)$$

where the average $\langle \dots \rangle$ is taken over the initial product separable quantum states of the particle and the bath oscillators at $t = 0$, $|\phi\rangle\{|\alpha_1\rangle|\alpha_2\rangle \dots |\alpha_N\rangle\}$. Here $|\phi\rangle$ denotes any arbitrary initial state of the particle and $|\alpha_i\rangle$ corresponds to the initial coherent state of the i -th bath oscillator. $|\alpha_i\rangle$ is given by $|\alpha_i\rangle = \exp(-|\alpha_i|^2/2) \sum_{n_i=0}^{\infty} (\alpha_i^{n_i} / \sqrt{n_i!}) |n_i\rangle$, α_i being expressed in terms of the mean values of the co-ordinate and momentum of the i -th oscillator, $\langle \hat{q}_i(0) \rangle - \langle \hat{x}(0) \rangle = (\sqrt{\hbar}/2\omega_i)(\alpha_i + \alpha_i^*)$ and $\langle \hat{p}_i(0) \rangle = i\sqrt{\hbar\omega_i}/2(\alpha_i^* - \alpha_i)$, respectively. It is important to note that $\langle \hat{F}(t) \rangle$ of Eq.(5) is a classical-like noise term which, in general, is a non-zero number because of the quantum mechanical averaging over the

co-ordinate and momentum operators of the bath oscillators with respect to the initial coherent states and arbitrary initial state of the particle and is given by

$$\langle \hat{F}(t) \rangle = \sum_j \left[\{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \} \kappa_j \cos \omega_j t + \langle \hat{p}_j(0) \rangle \kappa_j^{1/2} \sin \omega_j t \right] . \quad (6)$$

It is convenient to rewrite the c -number equation (5) as follows;

$$\langle \ddot{\hat{x}}(t) \rangle + \int_0^t dt' \gamma(t-t') \langle \dot{\hat{x}}(t') \rangle + \langle V'(\hat{x}) \rangle = F(t) \quad (7)$$

where we let the quantum mechanical mean value $\langle \hat{F}(t) \rangle \equiv F(t)$. We now turn to the *second* averaging. To realize $F(t)$ as an effective c -number noise we now assume that the momenta $\langle \hat{p}_j(0) \rangle$ and the shifted co-ordinates $\{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \}$ of the bath oscillators are distributed according to a canonical distribution of Gaussian forms as

$$\mathcal{P}_j = \mathcal{N} \exp \left\{ \frac{-[\langle \hat{p}_j(0) \rangle^2 + \kappa_j \{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \}^2]}{2\hbar\omega_j (\bar{n}_j + \frac{1}{2})} \right\} \quad (8)$$

so that for any quantum mechanical mean value $O_j(\langle \hat{p}_j(0) \rangle, \{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \})$ the statistical average $\langle \dots \rangle_S$ is

$$\begin{aligned} \langle O_j \rangle_S &= \int O_j(\langle \hat{p}_j(0) \rangle, \{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \}) \\ &\quad \times \mathcal{P}_j(\langle \hat{p}_j(0) \rangle, \{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \}) \\ &\quad \times d\langle \hat{p}_j(0) \rangle d\{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \} . \end{aligned} \quad (9)$$

Here \bar{n}_j indicates the average thermal photon number of the j -th oscillator at temperature T and $\bar{n}_j = 1/[\exp(\hbar\omega_j/k_B T) - 1]$ and \mathcal{N} is the normalization constant.

The distribution (8) and the definition of statistical average (9) imply that $F(t)$ must satisfy

$$\langle F(t) \rangle_S = 0 \quad (10)$$

and

$$\langle F(t)F(t') \rangle_S = \frac{1}{2} \sum_j \kappa_j \hbar\omega_j \left(\coth \frac{\hbar\omega_j}{2k_B T} \right) \cos \omega_j(t-t') \quad (11)$$

That is c -number noise $F(t)$ is such that it is zero-centered and satisfies the standard fluctuation-dissipation relation (FDR) as known in the literature in terms of the quantum statistical average of the noise operators. We emphasize that the method of ensemble averaging over c -number noise $F(t)$ is distinctly different from traditional methods of quantum statistical averaging and also from the method of Glauber-Sudarshan P -distribution. We discuss this in Appendix A.

We now add the force term $V'(\langle \hat{x} \rangle)$ on both sides of Eq.(7) and rearrange it to obtain

$$\dot{X} = P \quad (12a)$$

$$\dot{P} = - \int_0^t dt' \gamma(t-t') P(t') - V'(X) + F(t) + Q(t) \quad (12b)$$

where we put $\langle \hat{x}(t) \rangle = X(t)$ for notational convenience and

$$Q(t) = V'(\langle \hat{x} \rangle) - \langle V'(\hat{x}) \rangle \quad (13)$$

represents the quantum correction due to system degrees of freedom. Since $Q(t)$ is a fluctuation term Eq.(12b) offers a simple interpretation. This implies that the QGLE is governed by a c -number quantum noise $F(t)$ originating from the heat bath characterized by the properties (10) and (11) and a quantum fluctuation term characteristic of

the non-linearity of the potential. To proceed further we need a recipe for the calculation of $Q(t)$. This has been discussed earlier in several contexts^{14,15,16,17,18,19}. For the present analysis we summarize it as follows.

Referring to the quantum nature of the system in the Heisenberg picture, one may write.

$$\hat{x}(t) = \langle \hat{x} \rangle + \delta \hat{x} \quad (14a)$$

$$\hat{p}(t) = \langle \hat{p} \rangle + \delta \hat{p} \quad (14b)$$

where $\langle \hat{x} \rangle$ and $\langle \hat{p} \rangle$ are the quantum-mechanical averages and $\delta \hat{x}$, $\delta \hat{p}$ are the operators. By construction $\langle \delta \hat{x} \rangle$ and $\langle \delta \hat{p} \rangle$ are zero and $[\delta \hat{x}, \delta \hat{p}] = i\hbar$. Using Eqs.(14a) and (14b) in $\langle V'(\hat{x}) \rangle$ and a Taylor series expansion around $\langle \hat{x} \rangle$ it is possible to express $Q(t)$ as

$$Q(t) = - \sum_{n \geq 2} \frac{1}{n!} V^{(n+1)}(X) \langle \delta \hat{x}^n(t) \rangle \quad (15)$$

Here $V^n(X)$ is the n -th derivative of the potential $V(X)$. To the second order $Q(t)$ is given by $Q(t) = -\frac{1}{2}V'''(X)\langle \delta \hat{x}^2 \rangle$ where $X(t)$ and $\langle \delta \hat{x}^2(t) \rangle$ can be obtained as explicit functions of time by solving the following set of coupled equations (16)-(18) along with Eqs.(12a) and (12b);

$$d\langle \delta \hat{x}^2 \rangle / dt = \langle \delta \hat{x} \delta \dot{\hat{p}} + \delta \dot{\hat{p}} \delta \hat{x} \rangle \quad (16)$$

$$d\langle \delta \hat{x} \delta \dot{\hat{p}} + \delta \dot{\hat{p}} \delta \hat{x} \rangle / dt = 2\langle \delta \dot{\hat{p}}^2 \rangle - 2V''(X)\langle \delta \hat{x}^2 \rangle \quad (17)$$

$$d\langle \delta \dot{\hat{p}}^2 \rangle / dt = -V''(X)\langle \delta \hat{x} \delta \dot{\hat{p}} + \delta \dot{\hat{p}} \delta \hat{x} \rangle \quad (18)$$

We now make some pertinent remarks on the above equations for quantum corrections and the related approximations and their successive improvements. The set of equations (16)-(18) for calculation of quantum correction term to the lowest order on based on⁸ the equations $\delta \dot{\hat{x}} = \delta \dot{\hat{p}}$ and $\delta \dot{\hat{p}} = -V''(X)\delta \hat{x}$, when the system-bath correlation is taken into account through X in $V''(X)$ which is obtained as a solution to Eqs.(12a) and (12b). The system-reservoir correlation is included at the non-Markovian level in the dynamics involving quantum correction terms. This implies that we have neglected the two time correlation of the system like $\langle \delta \hat{p}(t)\delta \hat{p}(t') \rangle$. The next improvement would be to include the contribution due to these terms at least in the Markovian level to the quantum correction equations so that the right hand sides of Eqs.(17) and (18) may be modified to add $-\Gamma\langle \delta \hat{x} \delta \dot{\hat{p}} + \delta \dot{\hat{p}} \delta \hat{x} \rangle$ and $-2\Gamma\langle \delta \dot{\hat{p}}^2 \rangle$, respectively. Γ is the Markovian limit of dissipation. The quantum correction dynamics is thus taken into consideration at the Markovian level through Γ term and at the non-Markovian level through $V''(X)$ term while the mean values follow non-Markovian dynamics. Being exponential in nature the contribution of the former terms are felt only at long times when the quantum fluctuations in the low order has less significant role to play with. Furthermore, it is now well established^{33,34,35,36} that curvature of the potential $V''(X)$ controls the initial growth of the quantum fluctuations. Since $X(t)$ is stochastic $V''(X)$ imparts stochasticity in $\delta \ddot{\hat{x}} = -V''(X)\delta \dot{\hat{x}}$, which makes the average dynamics of $\langle \delta \hat{x}^2 \rangle$ exponentially diverging³⁸. We thus emphasize that the $V''(X)$ term dominantly controls the evolution of quantum fluctuations to a good degree of accuracy. In the overdamped limit, however, it is easy to incorporate quantum correction to very high orders. To this end one may start from equations $\delta \dot{\hat{x}} = \delta \dot{\hat{p}}$ and $\delta \dot{\hat{p}} = -\Gamma\delta \dot{\hat{p}} - V''(X)\delta \hat{x}$. Neglecting the inertial term $\langle \delta \ddot{X} \rangle$ in this limit we may write $-\Gamma\delta \dot{\hat{x}} = V''(X)\delta \hat{x}$ to yield the solutions $\langle \delta \hat{x}^n(t) \rangle$ for arbitrary n in the expression for $Q(t)$. This has been discussed in greater detail in¹⁷ in the of quantum Smoluchowskii equation. For a more accurate estimate of quantum corrections one may use higher order terms in $Q(t)$ which can be calculated by solving higher order (say, twelve equations for fourth order) coupled equations as derived in Appendix B. Under very special circumstance, it has been possible to include quantum effects to all orders¹⁹.

The above method of c -number representation of quantum Langevin equation has been utilized recently by us to derive exact quantum Kramers' equation^{15,16} in terms of the probability distribution function for calculation of escape rate from a metastable well and to address several other issues. In what follows we present a method for generation of c -number noise for numerical simulation of quantum Langevin equation (12a) and (12b) in the next section.

III. A NUMERICAL METHOD FOR GENERATION OF QUANTUM NOISE

Eq.(11), the fluctuation-dissipation relation is the key element for generation of quantum noise. The left hand side of this equation suggests that $F(t)$ is a 'classical' random number and $\langle F(t)F(t') \rangle_S$ is a correlation function which is classical in form but quantum-mechanical in its content. That is $F(t)$ is a c -number Langevin force such that

Eq.(12b) (along with (12a)) represents a stochastic differential equation. Since one of the most commonly occurring noise processes is exponentially correlated colour noise, we develop the scheme first for this process. This is based on a simple idea that c -number quantum noise may be viewed as a *superposition* of several Ornstein-Uhlenbeck noise processes. It may be noted that Imamoğlu³⁷, a few years ago, has produced a scheme for approximating a class of reservoir spectral functions by a finite superposition of Lorentzian functions with positive coefficients corresponding to a set of fictitious harmonic oscillator modes to incorporate non-Markovian effects into the dynamics. The aim of the work³⁷ is to develop a non-Markovian dynamics in terms of a master equation corresponding to the system-reservoir interaction for the expanded model after the tracing operation. The method is an extension of the stochastic wave function method in quantum optics. The present superposition method is reminiscent of this approach in the time domain. We however emphasize two important points. Because of rotating wave and Born approximations the master equation is valid for weak system-reservoir coupling i.e., weak damping regime. The present method on the other hand uses a superposition of noises in the time domain and the quantum Langevin equation is valid for arbitrary coupling since no weak coupling approximation or rotating wave approximation has been used in the treatment. Secondly, while the stochastic wave function approach as well as the density operator equation have no classical analogue, the present scheme makes use of c -numbers to implement the classical numerical method quite freely to solve quantum Langevin equation.

A. SUPERPOSITION METHOD

We proceed in several steps as follows;

1. Step I

We first denote the correlation function $c(t - t')$ as

$$\langle F(t)F(t') \rangle_S = c(t - t') \quad (19)$$

and in the continuum limit $c(t - t')$ is given by

$$c(t - t') = \frac{1}{2} \int_0^\infty d\omega \kappa(\omega) \rho(\omega) \hbar \omega \coth\left(\frac{\hbar \omega}{2k_B T}\right) \cos \omega(t - t') \quad (20)$$

where we have introduced the density function $\rho(\omega)$ for the bath modes. In principle an a priori knowledge of $\kappa(\omega)\rho(\omega)$ is essential and primary factor for determining the evolution of stochastic dynamics governed by Eqs.(12a) and (12b). We assume a Lorentzian distribution of modes so that

$$\kappa(\omega)\rho(\omega) = \frac{2}{\pi} \frac{\Gamma}{1 + \omega^2 \tau_c^2} \quad (21)$$

where Γ and τ_c are the dissipation in the Markovian limit and correlation time, respectively. These two quantities and the temperature T are the three input parameters for our problem. Eq.(21) when used in (4) in the continuum limit yields an exponential memory kernel $\gamma(t) = (\Gamma/\tau_c) \exp(-|t|/\tau_c)$. For a given set of these parameters we first numerically evaluate the integral (20) as a function of the time $(t - t')$. The typical profile of such a correlation is shown in Fig.1(a) for three different temperatures (solid line) and for fixed $\Gamma = 1$ and $\tau_c = 1$. Similarly we present the numerically calculated correlation functions for different values of Γ for a fixed temperature $T = 0.0$ and correlation time $\tau_c = 1.0$; (Fig 1(b), solid line) and for different values of τ_c (for fixed temperature $T = 0.0$ and dissipation $\Gamma = 1.0$; (Fig 1(c), solid line).

2. Step II

In the next step we numerically *fit* the correlation function with a superposition of several exponentials as

$$c(t - t') = \sum_i \frac{D_i}{\tau_i} \exp\left(\frac{-|t - t'|}{\tau_i}\right) \quad i = 1, 2, 3... \quad (22)$$

The set D_i and τ_i , the constant parameters are thus known. In Fig.1 (a-c) we exhibit these fitted curves (dotted lines) against the corresponding solid curves calculated by numerical integration of Eq.(20). It may be checked that at high temperatures the required fitting curve is in general a single exponential, whereas at very low temperatures down to absolute zero biexponential fitting is an excellent description. At the intermediate temperatures one has to use even triexponential fitting, in general.

3. Step-III

The numerical agreement between the two sets of curves based on (20) and (22) in each of the Figs.1(a-c) suggests that one may generate a set of exponentially correlated colour noise variables η_i according to

$$\dot{\eta}_i = -\frac{\eta_i}{\tau_i} + \frac{1}{\tau_i}\xi_i(t) \quad (23)$$

where

$$\langle \xi_i(t) \rangle = 0 \quad (24)$$

$$\langle \xi_i(0)\xi_j(\tau) \rangle = 2D_i\delta_{ij}\delta(\tau) \quad (i = 1, 2, 3\dots) \quad (25)$$

in which $\xi_i(t)$ is a Gaussian white noise obeying (24) and (25); τ_i and D_i are being determined in Step-II from numerical fit. The quantum noise η_i is thus an Ornstein-Uhlenbeck process with properties

$$\langle \eta_i(t) \rangle = 0 \quad (26)$$

$$\langle \eta_i(t)\eta_j(t') \rangle = \delta_{ij}\frac{D_i}{\tau_i}\exp\left(\frac{-|t-t'|}{\tau_i}\right) \quad (27)$$

Clearly τ_i and D_i are the correlation time and the strength of colour noise variables η_i , respectively. It is also important to point out that they have nothing to do with true correlation time τ_c and strength Γ of the noise $F(t)$ as expressed in (21) and (20).

4. Step-IV

The c -number quantum noise $F(t)$ due to the heat bath is therefore given by

$$F(t) = \sum_{i=1,2\dots} \eta_i \quad (28)$$

Eq.(28) implies that $F(t)$ can be realized as a sum of several Ornstein-Uhlenbeck noises η_i satisfying $\langle F(t)F(t') \rangle_S = \sum_i \langle \eta_i(t)\eta_i(t') \rangle$.

Having obtained the scheme for generation of c -number quantum noise $F(t)$ we now proceed to solve the quantum Langevin equations (12a) and (12b) along with Eqs.(16)-(18). To this end the first step is the integration^{20,21} of the stochastic differential equation (23). We note that to generate Gaussian white noise $\xi_i(t)$ of (23) from two random numbers which are uniformly distributed on the unit interval, Box-Mueller algorithm may be used. To be specific we start the simulation with integration of (23) to generate an initial value of η_i as follows;

$$m = \text{random number} \quad (29)$$

$$n = \text{random number} \quad (30)$$

$$\eta_i = \left[-\frac{2D_i}{\tau_i} \ln(m) \right]^{1/2} \cos(2\pi n) \quad (31)$$

We then set $E_i = \exp(-\Delta t/\tau_i)$, where Δt is the integration step length for (23). Following the integral algorithm of Fox *et al*²⁰, we now generate the next step $\eta_i|_{t+\Delta t}$ from η_i according to the following lines;

$$a = \text{random number} \quad (32)$$

$$b = \text{random number} \quad (33)$$

$$h_i = \left[-\frac{2D_i}{\tau_i}(1 - E_i^2) \ln(a) \right]^{1/2} \cos(2\pi b) \quad (34)$$

and

$$\eta_i|_{t+\Delta t} = \eta_i E_i + h_i \quad (35)$$

After (35) the algorithm is repeated back to (32) and continued for a sufficient time evolution.

In the next step we couple the above-mentioned integral algorithm for solving (23) to standard fourth order Runge-Kutta (or other) algorithm for the solution of Eqs.(12a), (12b) and (16)-(18). For an accurate and fast algorithm η -integration (35) may be carried out at every $\Delta t/2$ step, while for Runge-Kutta integration of Eqs.(12a), (12b) and (16)-(18) step size Δt is a good choice. With Δt in the range $\sim 0.001 - 0.01$ the overall method gives excellent results for the solution of quantum Langevin equation.

B. SPECTRAL METHOD

Our method of superposition of Ornstein-Uhlenbeck noises for generation of quantum noise as discussed above primarily pertains to situations where the correlation function can be described by a superposition of several exponentials. Although in overwhelming cases of applications in condensed matter and chemical physics at low-temperature where in general non-Ohmic conditions prevail we encounter such situations, there are cases where the noise correlation exhibits power law behaviour or is Gaussian in nature. To extend the present scheme we now show how the spectral method can be implemented for generation of quantum noise. The method had been used earlier by several groups^{22,23,24,25} in the context of *classical* long and short range temporally correlated noises. We give here a brief outline for its extension to quantum noise.

The starting point of our analysis is the knowledge of quantum correlation function $c(t - t')$ as given in (20). Again this depends on $\kappa(\omega)\rho(\omega)$, which (for example, for short range Gaussian correlated noise may be taken as $2\Gamma \exp(-\omega^2\tau_c^2/4)$, Γ and τ_c being the strength and the correlation time, respectively) must be known a priori. Having known $c(t - t')$ as a function of time we first carry out its Fourier transform

$$\begin{aligned} \langle F(t)F(t') \rangle_S &= c(t - t') \\ C(\omega) &= \int \exp(-i\omega t)c(t)dt \end{aligned} \quad (36)$$

by discretizing the time in $N = 2^n$ intervals of size Δt (such that Δt is shorter than any other characteristic time of the system). The result is a string of discrete numbers $C(\omega_n)$. In discrete Fourier space the noise correlation is given by

$$\langle F(\omega_n)F(\omega'_m) \rangle_S = N\Delta t C(\omega_n)\delta_{m+n,0} \quad (37)$$

Therefore the noise in the Fourier domain can be constructed simply as

$$\begin{aligned} F(\omega_n) &= \sqrt{N\Delta t C(\omega_n)}\alpha(\omega_n) \\ \text{where } i &= 0, 1, 2, \dots, N; \quad F(\omega_0) = F(\omega_n); \quad \omega_n = \frac{2\pi n}{N\Delta t} \end{aligned} \quad (38)$$

Here α_n are the Gaussian complex random numbers with zero mean and anti-delta correlated noise, such that $\alpha_n = a_n + ib_n$; a_n and b_n (the real and imaginary parts of α_n) are Gaussian random numbers with zero mean value and variance $1/2$.

$$\langle a_n \rangle = \langle b_n \rangle = 0; \quad (39)$$

$$\langle a_n^2 \rangle = \langle b_n^2 \rangle = \frac{1}{2}; \quad \text{where } n \neq 0 \quad (40)$$

$$\text{and } a_0^2 = 1 \quad (41)$$

Box-Mueller algorithm may again be used to generate the set of random numbers a_n, b_n (and consequently α_n) according to the lines;

$$p_n = \text{random number} \quad (42)$$

$$q_n = \text{random number} \quad (43)$$

$$a_n = (-\ln p_n)^{1/2} \cos 2\pi q_n \quad (44)$$

and similarly for b_n . The discrete Fourier transform of the string $F(\omega_n)$ is then numerically calculated by the Fast Fourier Transform algorithm to generate the quantum random numbers $F(t_n)$. The recipe can be implemented for solving Eqs.(12a) and (12b) and (16)-(18) as discussed in the earlier subsection III-A.

IV. APPLICATION: QUANTUM MEAN FIRST PASSAGE TIME CALCULATIONS

We now present the numerical results on c -number quantum Langevin equation for a cubic potential. The mean first passage time whose inverse represents the quantum rate of crossing the barrier is computed for different temperatures, correlation times and damping constants. The notion of mean in mean first passage time calculation needs some clarification. We are concerned here with evolution of quantum mean values $q(= \langle \hat{q} \rangle)$ and $p(= \langle \hat{p} \rangle)$ which can be simultaneously determined unlike their operator counterparts \hat{q} and \hat{p} , respectively. The quantum fluctuation around the mean path are however taken into account by following the equations of quantum corrections. Since the system is coupled to heat bath it is also necessary to take statistical ‘mean’ over several thousands of realizations of trajectories of the quantum mean values. One of the major aims of this numerical simulation is to show that the method described in subsection III-A can be implemented very easily to describe both thermally activated processes and tunneling within an unified scheme.

To begin with we consider a potential of the form

$$V(x) = -\frac{1}{3}Ax^3 + Bx^2 \quad (45)$$

where A and B are the constants. The other input parameters for our calculations (Eqs.(20) and (21)) are temperature T , strength of noise correlation Γ and correlation time τ_c . For the potential $V(x)$, $Q(t)$ is given by a single quantum correction term $Q = A\langle \delta \hat{x}^2 \rangle$ according to Eq.(15). Theoretical consideration demands that the activation barrier ΔE must be much larger than the thermal noise, i.e, $\Delta E \gg k_B T$, where ΔE denotes the height of the potential barrier. In order to ensure the stability of the algorithm we have kept $\Delta t/\tau_c \ll 1$ where Δt is the integration step size. For a given parameter set we have computed the quantum first passage time for the quantum Brownian particle in the cubic potential starting from the bottom of the metastable well at $x = 0$ to reach the maximum at $x = 2B/A$ for a single realization of a stochastic path. The quantum mean first passage time (QMFPT) is calculated by averaging over the first passage times over 5000 trajectories. We have used $\hbar = k_B = 1$ for the entire treatment.

We now present the results for the quantum mean first passage times obtained by the simulation of the quantum damped cubic potential (45) driven by quantum colour noise [Eqs.(12a),(12b), (16)-(18) and (23)]. Our results are shown for the rate $1/\text{QMFPT}$ as a function of Γ for three different temperatures $T = 3, 1$ and 0.0 in Fig.2(a-c). We have kept the parameter $\Delta E = 12$ with $A = 0.5$ and τ_c very small. It is observed that at $T = 3$ with increasing Γ , $1/\text{QMFPT}$ undergoes a turnover from an inverse trend ($1/\text{MFPT} \propto 1/\Gamma$) to a linear behaviour ($1/\text{QMFPT} \propto \Gamma$) in accordance with Kramers’ classic result²⁶ subsequently supported by many others^{27,28,29,30,31,32}. The $1/\text{QMFPT}$ has a maximum in the intermediate range. As the temperature is lowered to $T = 1$ and $T = 0$ the turnover persists with a little change of shape near the maxima. The Fig.2c represents the typical vacuum field assisted tunneling since the thermal activated process is almost absent at $T \sim 0$.

In Fig.3 we plot the rate $1/\text{QMFPT}$ vs. τ_c for two different temperatures, with $A = 1.0$ and $\Delta E = 4.0$ and $\Gamma = 1.0$. It is observed that as the temperature is lowered to $T = 0$ the decrease of $1/\text{QMFPT}$ exhibits two distinct time scales.

In Fig.4(a-b) we exhibit Arrhenius plot, i.e, QMFPT^{-1} vs. $1/T$ for several values of Γ . We have fixed $\Delta E = 10.0$ with $A = 1.5$ and $\tau_c = 1.0$. The physical situation corresponding to non-Ohmic underdamped to overdamped regions has been explored. It is apparent that when the temperature is high $\ln \text{QMFPT}^{-1}$ vs. $1/T$ is typically linear corresponding to the century old Arrhenius classical result. However, when the temperature is lowered the quantum effects and T^2 dependence is observed. This result is in complete conformity with those reported earlier in connection with the theoretical as well as experimental studies in mixed oxide systems and gas kinetics^{1,15}. The variation of Arrhenius profiles with Γ is also indicative of the turnover phenomenon as discussed earlier.

V. CONCLUSION

In this paper we have proposed a simple numerical method for generation of quantum noise and for solution of the associated generalized quantum c -number Langevin equation. The method is based primarily on two aspects. First, it has been shown that it is possible to realize quantum noise (in a quantum Langevin dynamics) which satisfies quantum fluctuation-dissipation relation as c -numbers rather than as operators whose ordering and quantum statistical averaging procedure is markedly different from those followed in the present scheme. Second, the quantum correction terms which appear due to the non-linearity of the potential of the system, can be generated systematically order by order, so that the coupled set of equations for the stochastic dynamics of the system and the correction terms can be solved easily with sufficient degree of accuracy. The method of simulation is equipped to deal with arbitrary correlation time, temperature and damping strength. The mapping of quantum stochastic dynamics as a problem of classical simulation as presented here allows us to implement classical methods in quantum simulations. The method is much simpler to handle compared to path integral quantum Monte-Carlo as employed earlier in the problem of barrier crossing dynamics¹⁰. We have shown that our simulation results on quantum mean first passage time for a particle in a cubic metastable potential agree well with the results obtained earlier. We hope that with the prescription given, many problems where the time correlation does not follow a typical prescribed dynamics can be simplified within the framework of this simulation.

Acknowledgments

The authors are indebted to the Council of Scientific and Industrial Research (CSIR), Government of India, for financial support under Grant No. 01/(1740)/02/EMR-II.

APPENDIX A: FLUCTUATION-DISSIPATION RELATION

In this section we derive quantum fluctuation-dissipation relation in three different ways.

Method-I: Quantum-Statistical Averaging

To make the discussion self-contained we first establish that the quantum-statistical average value of a two-time correlation function of the random force operator $\hat{F}(t)$ using standard usage of quantum-statistical averaging $\langle \dots \rangle_{QS}$ can be written as

$$\frac{1}{2} \langle \hat{F}(t) \hat{F}(t') + \hat{F}(t') \hat{F}(t) \rangle_{QS} = \sum_j \frac{1}{2} \kappa_j \hbar \omega_j \left(\coth \frac{\hbar \omega_j}{2 k_B T} \right) \cos \omega_j (t - t') \quad (\text{A1})$$

where $\hat{F}(t)$ is defined by Eq.(3). Starting from a canonical distribution of bath oscillators at $t = 0$ with respect to the free oscillator Hamiltonian \hat{H}_0 , with a shift of origin $\hat{q}'_j(0) = \hat{q}_j(0) - \hat{x}(0)$, where

$$\hat{H}_0 = \sum_j \left(\frac{\hat{p}_j^2(0)}{2} + \frac{1}{2} \kappa_j \hat{q}'_j{}^2(0) \right), \quad (\text{A2})$$

we construct the expectation value of an arbitrary operator \hat{O} as

$$\langle \hat{O} \rangle_{QS} = \frac{\text{Tr } \hat{O} e^{-\hat{H}_0/k_B T}}{\text{Tr } e^{-\hat{H}_0/k_B T}}. \quad (\text{A3})$$

As usual Trace implies both quantum mechanical averaging with respect to number states of the oscillators $\prod_{n_1=0}^{\infty} \{|n_1\rangle\} \prod_{n_2=0}^{\infty} \{|n_2\rangle\} \dots \prod_{n_N=0}^{\infty} \{|n_N\rangle\}$ and the system state $|\phi\rangle$ followed by Boltzmann statistical averaging. Making use of the definition of the quantum statistical average Eq.(A3) we first derive

$$\langle \hat{q}'_j(0) \hat{q}'_k(0) \rangle_{QS} = \frac{\langle \hat{p}_j(0) \hat{p}_k(0) \rangle_{QS}}{\omega_j^2} = \delta_{jk} \frac{\hbar \coth(\omega_j \hbar / 2k_B T)}{2\omega_j} \quad (\text{A4})$$

and

$$\langle \hat{q}'_j(0) \hat{p}_k(0) \rangle_{QS} = -\langle \hat{p}_k(0) \hat{q}'_j(0) \rangle_{QS} = \frac{1}{2} i \hbar \delta_{jk} \quad (\text{A5})$$

Since the expectation value of an odd number of factors of $\hat{q}'_j(0)$ and $\hat{p}_j(0)$ vanishes and the expectation value of an even number of factors is the sum of products of the pairwise expectation values with the order of the factors preserved we note that Gaussian property of the distribution of $\hat{q}'_j(0)$ and $\hat{p}_j(0)$ is implied.

The relations (A4) and (A5) directly yield (A1). On the other hand quantum mechanical averaging with respect to the number states results in

$$\langle F(t) \rangle_{QS} = 0 \quad (\text{A6})$$

Method-II: Glauber-Sudarshan P -function

In this method we first calculate the quantum mechanical average of $1/2[\hat{F}(t)\hat{F}(t') + \hat{F}(t')\hat{F}(t)]$ with respect to the coherent states of the bath oscillators $|\alpha_1\rangle|\alpha_2\rangle\dots|\alpha_N\rangle$ and system state $|\phi\rangle$ at $t = 0$ where $|\alpha_i\rangle$ has been defined in the text. $|\alpha_i\rangle$ is the eigenstate of the annihilation operator \hat{a}_i such that $\hat{a}_i|\alpha_i\rangle = \alpha_i|\alpha_i\rangle$ where $\hat{q}'_i(0)$ and $\hat{p}_i(0)$ are expressed as usual as $\hat{q}'_i(0) = \sqrt{\hbar/2\omega_i}[\hat{a}_i^\dagger(0) + \hat{a}_i(0)]$ and $\hat{p}_i(0) = i\sqrt{\hbar\omega_i/2}[\hat{a}_i^\dagger(0) - \hat{a}_i(0)]$. We thus note that the coherent state averaging $\langle \dots \rangle$ leads to (we have discarded the cross terms like $\alpha_j \alpha_k^*$ since they will vanish after the statistical averaging that follows in the next step),

$$\begin{aligned} \frac{1}{2} \langle \hat{F}(t)\hat{F}(t') + \hat{F}(t')\hat{F}(t) \rangle &= \sum_j \left[\kappa_j \frac{\hbar}{2\omega_j} \left(\alpha_j^{*2} + 2\alpha_j^* \alpha_j + 1 + \alpha_j^2 \right) \cos \omega_j t \cos \omega_j t' \right. \\ &\quad - \frac{\kappa_j \hbar \omega_j}{2} \left(\alpha_j^{*2} - 2\alpha_j^* \alpha_j - 1 + \alpha_j^2 \right) \sin \omega_j t \sin \omega_j t' \\ &\quad \left. + \frac{i\hbar}{4} \kappa_j \omega_j \left(2\alpha_j^{*2} - 2\alpha_j^2 \right) \sin \omega_j (t - t') \right]. \end{aligned} \quad (\text{A7})$$

We now introduce Glauber-Sudarshan distribution known as P -function for the statistical averaging of the quantum mechanical expectation values,

$$\begin{aligned} P_{GS}(\alpha_j(0), \alpha_j^*(0)) &= N \exp \left(-\frac{|\alpha_j|^2}{\bar{n}_j} \right) \\ &= N \exp \left[-\left(\frac{\langle \hat{p}_j(0) \rangle^2 + \omega_j^2 \langle \hat{q}'_j(0) \rangle^2}{2\hbar\omega_j \bar{n}_j} \right) \right] \end{aligned} \quad (\text{A8})$$

where N is the normalization constant. Noting that with the distribution (A8) the normalized statistical averages $\langle \dots \rangle_{GS}$ are given by,

$$\langle \alpha_j^2 \rangle_{GS} = \langle \alpha_j^{*2} \rangle_{GS} = 0 \quad \text{and} \quad \langle \alpha_j \alpha_k^* \rangle_{GS} = \bar{n}_j \delta_{jk} \quad (\text{A9})$$

With relations (A9), the statistical average (with respect to the P -function) of the quantum mechanical mean value (A7) leads us to

$$\frac{1}{2} \langle \langle \hat{F}(t)\hat{F}(t') + \hat{F}(t')\hat{F}(t) \rangle \rangle_{GS} = \frac{1}{2} \sum_j \kappa_j \hbar \omega_j \left(\coth \frac{\hbar \omega_j}{2k_B T} \right) \cos \omega_j (t - t') \quad (\text{A10})$$

and

$$\langle \langle \hat{F}(t) \rangle \rangle_{GS} = 0 \quad (\text{A11})$$

Method-III: Our Distribution Function

In this method we bypass the operator ordering of $\hat{F}(t)$ s. Instead, we start from coherent state average of the noise operator $\hat{F}(t)$, as given by (6), i.e., the c -number noise

$$\langle \hat{F}(t) \rangle = \sum_j \left[\{ \langle \hat{q}_j(0) \rangle - \langle \hat{x}(0) \rangle \} \kappa_j \cos \omega_j t + \langle \hat{p}_j(0) \rangle \kappa_j^{1/2} \sin \omega_j t \right] . \quad (\text{A12})$$

and then calculate the product of two-time c -numbers $\langle \hat{F}(t) \rangle \langle \hat{F}(t') \rangle$. In our notation we have written

$$\langle \hat{F}(t) \rangle \langle \hat{F}(t') \rangle \equiv F(t)F(t') \quad (\text{A13})$$

Expressing $\langle \hat{q}'_j(0) \rangle$ and $\langle \hat{p}_j(0) \rangle$ in terms of α_j and α_j^* as stated earlier, we may write

$$\begin{aligned} F(t)F(t') &= \left\{ \sum_j \left(\kappa_j \sqrt{\frac{\hbar}{2\omega_j}} (\alpha_j^* + \alpha_j) \cos \omega_j t + i \sqrt{\frac{\hbar \omega_j}{2}} \omega_j (\alpha_j^* - \alpha_j) \sin \omega_j t \right) \right\} \\ &\times \left\{ \sum_k \left(\kappa_k \sqrt{\frac{\hbar}{2\omega_k}} (\alpha_k^* + \alpha_k) \cos \omega_k t' + i \sqrt{\frac{\hbar \omega_k}{2}} \omega_k (\alpha_k^* - \alpha_k) \sin \omega_k t' \right) \right\} . \end{aligned} \quad (\text{A14})$$

We now introduce our distribution function (8) for ensemble averaging $\langle \dots \rangle_S$ over the product $F(t)F(t')$

$$\begin{aligned} P_j(\alpha_j(0), \alpha_j^*(0)) &= N \exp \left(-\frac{|\alpha_j|^2}{\bar{n}_j + 1/2} \right) \\ &= N \exp \left[-\left(\frac{\langle \hat{p}_j(0) \rangle^2 + \omega_j^2 \langle \hat{q}'_j(0) \rangle^2}{2\hbar \omega_j (\bar{n}_j + 1/2)} \right) \right] \end{aligned} \quad (\text{A15})$$

Noting that

$$\langle \alpha_j^2 \rangle_S = \langle \alpha_j^{*2} \rangle_S = 0 \quad \text{and} \quad \langle \alpha_j \alpha_k^* \rangle_S = (\bar{n}_j + \frac{1}{2}) \delta_{jk}, \quad (\text{A16})$$

we obtain on statistical averaging over $F(t)F(t')$

$$\langle F(t)F(t') \rangle_S = \frac{1}{2} \sum_j \kappa_j \hbar \omega_j \left(\coth \frac{\hbar \omega_j}{2k_B T} \right) \cos \omega_j (t - t') \quad (\text{A17})$$

Several points are now in order. First, we note that the fluctuation-dissipation relation has been expressed in several different but equivalent forms in Eqs.(A1), (A10) and (A17) (or Eq.(8)). Because of the prescription of operator ordering, the left hand sides of Eqs.(A1) and (A10) do not permit us to identify a c -number noise term. Clearly

Eq.(A17) suits this purpose and quantum noise as a number $F(t)$ can be identified in (A17) or Eq.(8). The ensemble averaging procedure of method-III is therefore followed in this work. Second, our distribution (8) is distinctly different from the Glauber-Sudarshan distribution (A8) because of a vacuum term $1/2$ in the width of the distribution (8). Third, the origin of distinction of the methods II and III lies in the fact that in the latter case we need not carry out an operator ordering of noise as in the former case since we are essentially dealing with a correlation of numbers rather than operators. The uncertainty principle is taken care of through the vacuum term $1/2$ in the width of the distribution (8) rather than through operator ordering as in method-II. Fourth, the distribution (8) as well as the Glauber-Sudarshan distribution (A8) reduces to classical Maxwell-Boltzmann distribution in the limit $\hbar\omega_j \ll k_B T$. Being a quasi-classical distribution, (A8) becomes singular in the vacuum limit $\bar{n} \rightarrow 0$. The distribution (8) on the other hand remains a valid non-singular distribution of quantum mean values whose width is simply the natural width due to non-thermal vacuum fluctuations.

APPENDIX B: EVOLUTION EQUATIONS FOR HIGHER-ORDER QUANTUM CORRECTIONS

The equations for quantum corrections upto fourth order are listed below. Equations for the second order are:

$$\begin{aligned}\frac{d}{dt}\langle\delta X^2\rangle &= \langle\delta X\delta P + \delta P\delta X\rangle, \\ \frac{d}{dt}\langle\delta P^2\rangle &= -2\Gamma\langle\delta P^2\rangle - V''\langle\delta X\delta P + \delta P\delta X\rangle - V'''\langle\delta X\delta P\delta X\rangle, \\ \frac{d}{dt}\langle\delta X\delta P + \delta P\delta X\rangle &= -\Gamma\langle\delta X\delta P + \delta P\delta X\rangle 2\langle\delta P^2\rangle - 2V''\langle\delta X^2\rangle - V'''\langle\delta X^3\rangle,\end{aligned}\tag{B1}$$

Those for the third order are:

$$\begin{aligned}\frac{d}{dt}\langle\delta X^3\rangle &= 3\langle\delta X\delta P\delta X\rangle, \\ \frac{d}{dt}\langle\delta P^3\rangle &= -3\Gamma\langle\delta P^3\rangle - 3V''\langle\delta P\delta X\delta P\rangle + V'''\left(\frac{3}{2}\langle\delta X^2\rangle\langle\delta P^2\rangle - \frac{3}{2}\langle\delta P\delta X^2\delta P\rangle + \hbar^2\right), \\ \frac{d}{dt}\langle\delta X\delta P\delta X\rangle &= -\Gamma\langle\delta X\delta P\delta X\rangle + 2\langle\delta P\delta X\delta P\rangle - V''\langle\delta X^3\rangle - \frac{V'''}{2}\left(\langle\delta X^4\rangle - \langle\delta X^2\rangle^2\right), \\ \frac{d}{dt}\langle\delta P\delta X\delta P\rangle &= -2\Gamma\langle\delta P\delta X\delta P\rangle + \langle\delta P^3\rangle - 2V''\langle\delta X\delta P\delta X\rangle \\ &\quad + \frac{V'''}{2}\left(\langle\delta X^2\rangle\langle\delta X\delta P + \delta P\delta X\rangle - \langle\delta X^3\delta P + \delta P\delta X^3\rangle\right),\end{aligned}\tag{B2}$$

And lastly, the fourth order equations are:

$$\begin{aligned}\frac{d}{dt}\langle\delta X^4\rangle &= 2\langle\delta X^3\delta P + \delta P\delta X^3\rangle, \\ \frac{d}{dt}\langle\delta P^4\rangle &= -4\Gamma\langle\delta P^4\rangle - 2V''\langle\delta X\delta P^3 + \delta P^3\delta X\rangle + 2V'''\langle\delta X^2\rangle\langle\delta P^3\rangle, \\ \frac{d}{dt}\langle\delta X^3\delta P + \delta P\delta X^3\rangle &= -\Gamma\langle\delta X^3\delta P + \delta P\delta X^3\rangle - 2V''\langle\delta X^4\rangle - 3\hbar^2 + 6\langle\delta P\delta X^2\delta P\rangle \\ &\quad + V'''\langle\delta X^2\rangle\langle\delta X^3\rangle, \\ \frac{d}{dt}\langle\delta X\delta P^3 + \delta P^3\delta X\rangle &= -3\Gamma\langle\delta X\delta P^3 + \delta P^3\delta X\rangle + 2\langle\delta P^4\rangle + 3V''(\hbar^2 - 2\langle\delta P\delta X^2\delta P\rangle) \\ &\quad + 3V'''\langle\delta X^2\rangle\langle\delta P\delta X\delta P\rangle, \\ \frac{d}{dt}\langle\delta P\delta X^2\delta P\rangle &= -2\Gamma\langle\delta P\delta X^2\delta P\rangle - V''\langle\delta X^3\delta P + \delta P\delta X^3\rangle + \langle\delta P^3\delta X + \delta X\delta P^3\rangle \\ &\quad + V'''\langle\delta X^2\rangle\langle\delta X\delta P\delta X\rangle.\end{aligned}\tag{B3}$$

The derivatives of $V(X)$, i.e., V'' or V''' etc. in the above expressions are functions of X .

-
- * Present Address: Max-Planck-Institut für Physik komplexer Systeme, Nöthnitzer Str. 38, 01187 Dresden, Germany
- † Electronic address: pcdsr@mahendra.iacs.res.in
- ¹ U. Weiss, *Quantum Dissipative Systems*, (World Scientific, Singapore, 1999).
 - ² W.H. Louisell, *Quantum Statistical Properties of Radiation* (Wiley & Sons, NY, 1973).
 - ³ G. Gangopadhyay and D.S. Ray in *Advances in Multiphoton Processes and Spectroscopy*, vol. 8, edited by S.H. Lin, A.A. Villayes and F. Fujimura (World Scientific, Singapore, 1993).
 - ⁴ H. Grabert, P. Schramm, G.L. Ingold, Phys. Rep. **168**, 115 (1988).
 - ⁵ P.G. Wolynes, Phys. Rev. Lett. **47**, 968 (1981).
 - ⁶ R. Egger and C.H. Mak, Phys. Rev. B **50**, 15210 (1994).
 - ⁷ D.E. Makarov and N. Makri, Chem. Phys. Lett. **221**, 482 (1994).
 - ⁸ J. Dalibard, Y. Castin and K. Molmer Phys. Rev. Lett. **68**, 580 (1992).
 - ⁹ I. Percival, *Quantum State Diffusion* (Cambridge University Press, Cambridge, UK, 1998).
 - ¹⁰ M. Topaler and N. Makri, J. Chem. Phys. **101**, 7500 (1994).
 - ¹¹ J. Ankerhold, P. Pechukas and H. Grabert, Phys. Rev. Lett. **87**, 086802 (2001).
 - ¹² S.T. Stockburger and H. Grabert, Phys. Rev. Lett. **88**, 170407 (2002).
 - ¹³ J. Ray Chaudhuri, B.C. Bag and D.S. Ray J. Chem. Phys. **111**, 10852 (1999).
 - ¹⁴ S.K. Banik, B.C. Bag and D.S. Ray, Phys. Rev. E **65**, 051106 (2002).
 - ¹⁵ D. Banerjee, B.C. Bag, S.K. Banik and D.S. Ray, Phys. Rev. E **65**, 021109 (2002).
 - ¹⁶ D. Banerjee, S.K. Banik, B.C. Bag and D.S. Ray, Phys. Rev. E **66**, 051105 (2002).
 - ¹⁷ D. Banerjee, B.C. Bag, S.K. Banik and D.S. Ray, Physica A **318**, 6 (2003).
 - ¹⁸ B. Sundaram and P.W. Milonni, Phys. Rev. E **51**, 1971 (1995).
 - ¹⁹ A.K. Pattanayak and W.C. Schieve, Phys. Rev. E **50**, 3601 (1994).
 - ²⁰ R.F. Fox, I.R. Gatland, R. Roy and G. Vemuri, Phys. Rev. A **38**, 5938 (1988).
 - ²¹ C. Mahanta and T.G. Venkatesh, Phys. Rev. E **58**, 4141 (1998).
 - ²² H.A. Makse, S. Havlin, M. Schwartz and H.E. Stanley, Phys. Rev. E **53**, 5445 (1996).
 - ²³ A. Barabasi and H.E. Stanley, *Fractal Concepts in Surface Growth* (Cambridge University Press, NY, 1995).
 - ²⁴ A.H. Romero and J.M. Sancho, Phys. Rev. E **58**, 2833 (1998).
 - ²⁵ A.H. Romero and J.M. Sancho, J. Comp. Phys. **156**, 1 (1999).
 - ²⁶ H.A. Kramers, Physica (Amsterdam) **7**, 284 (1940).
 - ²⁷ R.F. Grote and J.T. Hynes, J. Chem. Phys. **73**, 2715 (1980).
 - ²⁸ P. Hänggi and F. Mojtabai, Phys. Rev. A **26**, 1168 (1982).
 - ²⁹ B. Carmeli and A. Nitzan, J. Chem. Phys. **79**, 393 (1983).
 - ³⁰ D. Kohen and D.J. Tannor, *Advances in Chemical Physics*, vol.111, edited by I. Prigogine and Stuart A. Rice, (John Wiley & Sons, Inc., 1999).
 - ³¹ J.K. Bhattacharjee, *Statistical Physics: Equilibrium and Non-Equilibrium Aspects* (Allied Publishers, New Delhi, 1997).
 - ³² K. Lindenberg and B.J. West, *The Nonequilibrium Statistical Mechanics of Open and Closed Systems* (VCH, New York, 1990).
 - ³³ L. Bonci, R. Roncaglia, B.J. West and P. Grigolini, Phys. Rev. Lett. **67**, 2593 (1991).
 - ³⁴ R.F. Fox and T.C. Elston, Phys. Rev. E **49**, 3683 (1994).
 - ³⁵ S. Chaudhuri, G. Gangopadhyay and D.S. Ray, Phys. Rev. E **54**, 2359 (1996).
 - ³⁶ B.C. Bag and D.S. Ray, J. Stat. Phys. **96**, 271 (1999).
 - ³⁷ A. Imamoglu, Phys. Rev. **50**, 3650 (1994); see also J. Dalibard, Y. Castin and K. Molmer, Phys. Rev. Lett. **68**, 580 (1992).
 - ³⁸ N.G. van Kampen, Phys. Rep. **24**, 171 (1976).




FIG. 1: Plot of correlation function $c(t)$ vs. t (units arbitrary) as given by Eq.(20) (solid line) and Eq.(22) (dotted line) for the set of parameter values mentioned in the text.




FIG. 2: Plot of inverse of quantum mean first passage time $\frac{1}{QMFPT}$ vs. dissipation parameter Γ (units arbitrary) for different temperatures: (a) $T = 3.0$ (b) $T = 1.0$ and (c) $T = 0.0$.




FIG. 3: Plot of inverse of quantum mean first passage time $\frac{1}{QMFPT}$ vs. correlation time τ_c (units arbitrary) for two different temperatures, $T = 0.0$ and $T = 1.0$, and for a fixed $\Gamma = 1.0$




FIG. 4: Plot of logarithm of inverse of quantum mean first passage time $\ln(\frac{1}{QMFPT})$ vs. $1/T$ (units arbitrary) for three different Γ values, $\Gamma = 2.0$, $\Gamma = 8.0$ and $\Gamma = 20.0$.