M etastable states, anom alous distributions and correlations in the HMF m odel

A lessandro P luchino, V ito Latora, Andrea Rapisarda

D ipartim ento di Fisica e Astronom ia, Universita di Catania, and INFN sezione di Catania, Via S.So a 64, I-95123 Catania, Italy

Revised version 01/04/2003 - Physica D in press

A bstract

We study the microscopic dynamics of the metastable Quasi-Stationary States (QSS) in the Hamiltonian Mean Field (HMF) model, a Hamiltonian system of N classical inertial spins with in nite-range interactions which shows a second order phase transition. In order to understand the origin of metastability, which appears in an energy region below the critical point, we consider two di erent classes of out-of-equilibrium initial conditions, both leading to QSS, and having respectively initial magnetization equal to one $(M \mid IC)$ and equal to zero $(M \mid IC)$. We com pare the corresponding -space, the resulting velocity pdfs and correlations, and the eventual aging features of the m icroscopic dynam ics. In both cases the m odel exhibits non-gaussian pdfs, though anom alous correlations are present only when the system is started with an initial magnetization equal to one. In the M 0 IC case the relaxation to equilibrium is almost exponential, while, for M 1 IC, when correlations and aging are found, the decay is a power-law and the overall behavior can be very well reproduced by a T sallis q-exponential function. These results contribute to clarify the overall scenario, which is more complex than previously expected and stress the importance of the dynamics in the relaxation process. The nonextensive statistical mechanics form alism proposed by T sallis seems to be valid, in the out-ofequilibrium phase, when correlations and strong long-term memory e ects em erge. This regime becomes stable if the N ! 1 limit is performed before the t! 1 lim it.

Key words: Ham iltonian dynam ics; Long-range interaction; Out-of-equilibrium statisticalmechanics PACS: 05.50.+q, 05.70Fh, 64.60Fr

Em ailaddresses: alessandro.pluchino@ct.infn.it (A lessandro P luchino), vito.latora@ct.infn.it (V ito Latora), andrea.rapisarda@ct.infn.it (A ndrea Rapisarda).

1 Introduction

There is by now a huge number of papers devoted to understanding the physical foundation and the possible applications of the nonextensive statistical mechanics formalism proposed by Constantino Tsallis in 1988 [1]. In particular, although the topic is still a matter of debate [2], the last few years have seen an important turning point for what concerns dissipative systems both lowdimensional (for example unimodal maps) β {5] and high-dimensional ones (for example turbulence, plasm a and nuclear physics, econophysics) [6{11]. Progress has been made also for conservative system s both for maps [12] and for high-dimensional ham iltonian systems with long-range interactions [13] 19], but open problems still remain and need a more detailed investigation. W ith this scenario in m ind, in this paper we present and discuss new num erical simulations relative to the Hamiltonian Mean Field (HMF) model, a system of N fully coupled classical spins which has been intensively studied in the last years. It is now adays well known [13,14,20 [26] that, for som e values of the initial energy, below the critical point, the system does not immediately relax to the Boltzm ann-Gibbs equilibrium, but remains trapped in anom abus metastable states and exhibits non-Gaussian velocity distributions that can be reproduced by the probability density functions (pdfs) predicted by the T sallis nonextensive therm odynam ics [1]. Such states have been nam ed Quasi-Stationary States (QSS) because they become stable in the therm odynamic limit if the N ! 1 limit is taken before the t ! 1 limit. In fact in such a case the force between spins tends to zero and the largest Lyapunov exponent vanish [13,14].

In this work we study two classes of out-of-equilibrium initial conditions that lead to QSS, and we discuss how the di erent initial conditions may a ect the dynam ics of the relaxation towards equilibrium. In particular, we concentrate our attention on the existence of correlations in the orientation angles and velocity of the spins, on the presence of anom abus velocity distributions, and on the clusters formations. The main result of this paper is that T sallis nonextensive therm odynam ics formalism seems to provide a general fram ework, to interpret the statistical properties of the dynam ically created QSS, for H am iltonian system swith long-range interactions [1], only when long-range correlations and fractal structures in phase space are present. Recently som e critical comments [27] have been raised on a previous study of two of us [13]. In this paper we provide new calculations and detailed discussions which reply in part to this criticism and clarify also some points advanced in another work [28]. A more detailed reply to the points raised in ref. [27] will be presented elsewhere [29].

The paper is organized as follows. In section II the details of the model are sum marized and the initial conditions used are discussed. Num erical results for

Q SS are discussed in section III, while velocity pdfs and dynam ical correlations are studied in a quantitative way in section IV . Conclusions are drawn in section V .

2 M odel and initial conditions

The HMF model, consists of N planar classical inertial spins interacting through an in nite-range potential [20]. The Ham iltonian can be written as:

$$H = K + V = \frac{x^{N}}{\frac{p_{i}^{2}}{2}} + \frac{1}{2N} \frac{x^{N}}{\frac{1}{2}} \left[1 - \cos(\frac{1}{2N})\right] :$$
(1)

In the latter $_{i}$ is the orientation angle of the ith spin and p_{i} the conjugate variable representing the rotational velocity, considering the mass of the rotators equal to one. The sum mation in V is extended to all couples of spins and is not restricted to rst neighbors. In order to make H form ally extensive, ie.V / N when N ! 1 [1,15], the coupling constant in the potential is divided by N. However, this often adopted prescription cures only a part of the problem since the energy remains non-additive, that is the system cannot be simply divided in two independent sub-system s. The magnetization M , i.e. the $m \text{ odulus of } M = \frac{1}{N} \prod_{i=1}^{P} m_i, w \text{ hith } m_i = [\cos(i); \sin(i)], \text{ is the order param}$ eter of the model. The latter shows a a second-order phase transition from a low-energy clustered phase with magnetization M 1, to a high-energy one, where the spins are hom ogeneously oriented on the unit circle and M 0.The m odel can be solved exactly in the canonical ensemble and the dependence of the energy density U = E = N on the tem perature T, is given by [20,21]

$$U = \frac{T}{2} + \frac{1}{2} \ 1 \ M^2 \quad : \tag{2}$$

The critical point is at energy density $U_c = \frac{3}{4}$ corresponding to a critical tem perature $T_c = \frac{1}{2}$ [20]. The dynam ics of HMF m odel can be investigated by starting the system with out-of-equilibrium initial conditions and integrating numerically the equations of motion [21]. In a special region of energy values $(\frac{1}{2} < U < U_c)$ the results of the simulations show, for a transient regime which depends on the system size, a disagreem ent with the canonical ensemble. In this region the dynam ics is characterized by Levy walks and anom alous di usion, while in correspondence the system shows a negative speci c heat [23]. Ensemble inequivalence and negative speci c heat have also been found in self-gravitating system s [31,32], nuclei and atom ic clusters [34{36}], though in the present m odel such anom alies emerge as dynam icalm etastable features [24,26] and vanish, if N is nite, for t ! 1. In this paper we focus on a

particular energy value belonging to the anom alous region, nam ely U = 0.69, and we study the time evolution of temperature, -space structures, dynam ical clusters in angle, velocity correlations and velocity distributions considering two out-of-equilibrium initial conditions in particular:

the rst one, used already in previous papers, consists of all orientation angles $_i = 0$ and velocity p_i uniform by distributed accordingly to the total energy available. In this case the potential energy is initially zero and we have the maximum of kinetic energy, while the total magnetization M is equal to 1;

in the second case considered, both angles and velocities are uniform ly distributed. This implies that the total magnetization M is initially zero, so that the potential energy is $\frac{1}{2}$, see eq.(2). Therefore in this case only a smaller fraction of total energy is left for the kinetic part.

For both initial conditions the totalm om entum is initially put to zero. In the following, for brevity, we will indicate these two initial conditions respectively by M 1 and M 0, thus referring to the initial value of the magnetization.

3 M etastable equilibrium and Quasi-Stationary States

W e discuss in this section the num erical results obtained starting from M 1 and M 0 initial conditions. D i erent sizes of the system are studied. The accuracy of the calculations is such that one gets an energy conservation $E = E = 10^{-5}$. A s in previous papers the fourth order Y oshida sim plectic algorithm [30] has been used for the integration. At variance with the previous publications the averages where taken only over the ensemble and not also along the time evolution. This point will be discussed in detail later.

In g.1 we plot, for system sizes N = 500 and N = 1000, the time evolution of twice the kinetic energy per particle, a quantity which is the tem perature in stationarity situations. D i erent regimes exist according to the initial conditions considered. M ore speci cally, when starting with M 1 initial condition (IC) four regimes are observable:

- (1) a fast decay from M = 1 to M = 0 in a time which is about 80–100. No sensible dependence on the size is observed;
- (2) a plateau region with M (N) < M $_{\rm eq}$ and T (N) < T $_{\rm eq}$ whose duration depends also on the size of the system as N [13];
- (3) a power-law relaxation towards the true equilibrium value $T_{eq} = 0.476$ [14];
- (4) an equilibrium regime with $M_{eq} = 0.307$.

In refs.[13,14] the kinetic energy per particle was actually computed considering not only an ensemble average, but also a time average. This was done to in prove statistics, though in principle time averages are justilled only when correlations do not exist. In g2 the old procedure and the new one are com pared. One observes that the two methods give the same plateau if a large num ber of events are considered in the ensemble average. The only di erence is in the decay from the initial condition to the QSS regime. One can easily check that also the scaling laws found in ref. [13] continue to be valid. In particular the lifetime of the plateau diverges with N and the value of the plateau tem perature T_{QSS} (N) converges towards the asymptotic tem perature $T_1 = 0.38 \text{ as N} ! 1$ following the same laws of refs.[13]. At variance with the criticism advanced in [27], it is evident that one can really talk of an approxim ate plateau for the average kinetic energy per particle, see g2. The log timescale used in g.1 and in previous papers [13] is useful to have an overall view of the tem poral behavior and does not change the m acroscopic quasistationary features. This will be clearer when analizing also the M 0 IC, for which the system is initially put in the state at M 0 and T 0:38 and there it remains for quite a long time which also in this case depends on the size studied. In fact when starting with M 0 IC only three regimes are observable:

- a plateau region with M
 and T
 38 whose length depends on the size as N
 This has been checked num erically but it is not shown here for lack of space;
- (2) a power-law relaxation towards the true equilibrium value, but faster than the previous M 1 case;
- (3) an equilibrium regime with $M_{eq} = 0.307$.

In general, for both \mathbb{IC} , the values obtained for the magnetization and the temperature in the plateau regime for N ! 1 are consistent with the caloric curve (2) extended below the critical point. Therefore this average procedure is a posteriori justi ed and consistent with a metastable macroscopic therm odynamic description, contrary to what claimed in ref.[27].

A lthough both initial conditions drive the system towards the same out-ofequilibrium macroscopic values for the magnetization and the temperature in the N ! 1 limit, for the QSS regime, the corresponding microscopic dynamics is very much di erent in the two cases. This fact can be seen when one plots the time evolution of the -space, as done in g.3 for a typical event and N = 10000. In red we represent the rotators having at time t = 0 the highest velocities. In this way it is possible to visualize the kind of mixing occurring for the initial conditions considered. It is im mediately evident that there are correlations and structures in the plateau regime only for M 1 IC, while for M 0 IC the distributions remain hom ogeneous with a very slow mixing of the initially fast rotators. This di erent behavior can be observed also in gure 4, where we plot the pdf in angle at various times. A typical realization for N = 10000 is plotted. One observes clusters formation in the plateau regime only for M1 while these are completely absent for M0 IC. However, at equilibrium, the pdfs are equivalent and the plots, for both M1 and M0, show only one rotating big cluster corresponding to M = 0.307 which is the equilibrium m agnetization (according to the canonical prediction eq.(2) with U = 0.69 and T = 0.476).

These num erical results reinforce our previous statem ent that although m acroscopically sim ilar the two m etastable plateaux have a very di erent m icroscopic dynam ics. Further evidence is provided if one studies the entropy and free energy as a function of time.

This is done in gs.5 and 6 respectively. To obtain the entropy we adopted a lattice over the -space m ade up of n = 10000 cells and we used the following form ula:

$$S(t) = \langle \prod_{k=1}^{X^{n}} f_{k}(t) \ln (f_{k}(t)) \rangle; \qquad (3)$$

where f_k is the relative frequency of points for the kth cell. The brackets < :::> denote ensemble averages. The curves plotted in the gures refer to an ensemble of 1000 events for N = 1000. This particular size was used in this case and in other calculations when a good average over the ensemble was essential in order to have a robust value, computer time being too long for greater sizes and such big ensemble averages. On the other hand no qualitative changes are anyhow observed for greater sizes. As intuitively expected from the temporal evolution of the -space, the entropy grow swith time towards equilibrium only for M 1 IC due to the high degree of order of this initial conguration, while it remains xed at the maximum value for the other, much more uniform, initial condition, M 0. Correspondingly, we compute the free energy by extrapolating to a dynamical situation the equilibrium therm odynam ic form ula

$$F(t) = U T(t)S(t)$$
: (4)

As shown in g.6 the free energy remains constant for M 0 in the QSS regime. This implies that for this IC the system is consistently in a metastable regime, which is not however the true one that is reached only at greater times, when the free energy of the system has a real global minimum . On the other hand, for the M 1 IC, the gure shows a dierent tem poralevolution before equilibration. A fter a short rapid transient decay, an almost at region, at a dierent free energy value, always emerges in the QSS regime, implying also in this case the presence of a metastable situation. The use of formula (4) is thus a posteriori justied since the temperature is almost constant in the plateau regime. It is also very useful in clarifying the scenario that en erges which is consistent with a quasi-equilibrium situation. This results reinforce the validity of our procedure to extract average therm odynam ic quantities from m icroscopic dynam ics and can be considered a further answer to the criticism raised in ref.[27]. On the other hand, the fact that there is no plateau for the tem poral evolution of the entropy for M 1 IC rejects the slow m icroscopic dynamics observed in the evolution of the -space structures. This is perfectly in line with the quantitative analysis perform ed by m eans of the correlation dim ension in ref.[13]. Thus one can summarize this situation by saying that only at a macrosopic level. the quasi-stationarity is valid, while m icroscopically the system never stops its slow temporal evolution. The evolution really stops only for N = 1, since in this case the magnetization in the QSS is exactly zero and so is the force which is proportional to M . In this sense the inversion of the time limit with the size limit to in nity stabilize the QSS regime. However, when considering nite N systems, the uctuations due to nite-size noise, of the order 1= N, induœ a force which is not exactly zero, and when they grow with time, the system is slow ly forced towards equilibration. As shown in ref.[13] the accuracy of our calculations is much smaller than this nite-size noise.

For the M 0 IC no entropy increase is observed. But also in this case some m icroscopic dynam ics exists. This is proved by the m ixing of fast particles in g.3. However no holes or structures are present in this case and this is the reason why the entropy tem poral evolution is almost at.

It is in portant to stress the fact that, for very long times, both the two initial conditions lead the system towards the equilibrium expected values.

W e would like to notice at this point that the slope of the entropy growth in g.5 does not give the K olm ogorov-Sinaientropy as dem onstrated num erically form aps [4,38]. In fact in our case we are considering the reduced -space and not the complete phase-space which has to be used. Unfortunately in our case the phase space has too m any dimensions and this fact prevents the direct investigation of the generalized q-entropy proposed by T sallis along the same lines of ref.[4] for m aps.

4 A nom alous velocity pdfs, velocity correlations and aging

In this section we study in a quantitative way the eventual presence of velocity correlations and memory e ects in the HMF dynamics. In Fig.7 we report the velocity pdfs for the plateau regime (at t = 1000) and at equilibrium (t = 1000000) and for the case N = 10000. For both IC, the initial velocity pdfs quickly acquire and maintain during the entire duration of the metastable state a non-G aussian shape. But while for M1 IC they can be tted with a

truncated generalized T sallis pdf [13], this is not possible for M 0 IC. In fact in this case the pdf shows a at alm ost rectangular shape, which clearly reveals the fact that structures and correlations, present in the other case, are here m issing. F inally, for very long times the system reaches always the equilibrium distribution and the agreem ent with the gaussian is alm ost perfect.

A quantitative way to estimate the velocity correlations is the calculation of the two-time autocorrelation function, de ned as

$$C (t; 0) = \frac{\langle P(t) P(0) \rangle \langle P(t) \rangle \langle P(t) \rangle \langle P(0) \rangle}{p(t) p(0)};$$
(5)

where $P = (p_1; p_2; ::: p_N)$ is the N-component velocity vector and the brackets < :::> indicate the average over the ensemble, while p(t) and p(0) are the standard deviations at time t and at time zero. In g.8 we plot the velocity autocorrelation functions for M 1 and M 0 IC in the QSS regime, obtained for the case N = 1000 and an average over 500 events. It is immediately evident that for M 0 the autocorrelation decays faster than for M 1 IC. This result clari es the controversial numerical results obtained in ref.[28] by using a slightly di erent, but equivalent, de nition for the autocorrelation function. In fact we do nd an alm ost exponential behavior only for the M 0 IC, while in ref. [28] the author claim ed to nd an exponential in both cases. In general we can reproduce the autocorrelation decay by means of the q-exponential function

$$e_{q}(x) = A \ 1 \ (1 \ q)^{\frac{X}{-1}}$$
 (6)

proposed by T sallis in his therm odynam ic nonextensive q-form alism [1], where A is the saturation value and the characteristic time. One gets the usual exponential for q = 1. In this way we can quantitatively discriminate between the two dimensional conditions. In fact we get for M 1 the value q = 1.55, revealing a power law scaling, while for M 0 we get a value of q much smaller, i.e. q = 1.12. This means that for M 0 the decay tends to be exponential (an exact exponential is obtained when q = 1). This almost exponential decay is also evident in the inset of the gure, where we show the same data in linear-log scale. An exponential t is also reported for comparison. Such num erical results may be considered a quantitative con mation of the dimension, in spite of the macroscopic presence of Q SS for both cases.

It is interesting to investigate also if the relaxation process observed in g.8 depends on the history of the system, i.e the possible existence of strong long-term m em ory e ects, the so-called aging phenom enon. This is typical of glassy system s but it has also been found in a great variety of com plex system s [39{ 41]. Recently it has been observed also in the HMF m odel by M ontem urro,

Tam arit and Anteneodo [18]. In the following we perform a similar analysis to study this e ect for both M1 and M0 IC, but considering only the N-component velocity vector de ned previously. In their case also angles were taken into account in the state vector entering the correlation function and only M1 IC were considered. The autocorrelation function to be considered is that one de ned before but calculated with respect to a set of four di erent waiting times t_w (that plays the role of age of the system)

$$C (t + t_w; t_w) = \frac{\langle P(t + t_w) P(t) \rangle \langle P(t + t_w) \rangle \langle P(t + t_w) \rangle}{p(t + t_w) p(t_w)} : (7)$$

In Fig.9 (a) we show the autocorrelation decay for M 1 IC, while in Fig.10 (a)that one for M 0 IC. If aging were not present, all the curves should overlap into one, because the function C (t+ t_w ; t_w) would depend only on the di erence between the two times t+ t_w and t_w , i.e. C (t+ t_w ; t_w) = C (t), and the decay would be exponential. Actually this is not what happens as shown in the gures.

For M 1 IC aging properties surely exist, i.e. it can be observed a characteristic two times dependence on t and t_w and the data can be scaled onto one curve, panel(b), considering as discussed in ref.[42]:

$$C (t + t_w; t_w) = f(\frac{t}{t_w});$$
 (8)

 $=\frac{1}{4}$. In such a way we obtain, for the tail a power law decay, i.e. with f (t=t,) $(t=t_{x})$ with scaling parameter = 1:54 that is here characteristic of a slow relaxation dynamics and long-term memory e ects. A g-exponential curve can reproduce very well this behavior with $q = \frac{+1}{-1} = 1:65$, A = 0:7 = 60. Notice that the scaling indices are di erent from those found and in ref.[18]. A sistem atic study of these scaling exponents would be desirable, but is beyond the scope of the present work and we postpone it to future investigations. As discussed in ref. [18] the origin of aging in the HMF model is not completely clear. The most likely scenario is that of a weak ergodicitybreaking. As originally proposed by Bouchaud for glassy systems in ref.[40], the latter occurs when the phase-space is a priori not broken into mutually inaccessible regions, as in the true ergodicity-breaking case, but the system can rem ain trapped for very long times in some regions of the complex potential landscape. In HMF the weak ergodicity-breaking can be related to the com plex dynam ics generated by the vanishing of the largest Lyapunov exponent and by a sort of dynam ical frustration due to the existence, in the QSS regime, of dierent small clusters (see q.4). These clusters compete in trapping more and more particles until only one of them remain when the system reach the standard equilibrium .

On the other hand, a di erent behavior is obtained for M O IC, as shown in g. 10. The same scaling law used for the M 1 IC case does not seem to apply, see panel (b), and the decay is now more rapid. In this case it is not so clear as before if aging does really exist, but again a quantitative di erence emerges for the two dynamics. A more detailed study in this respect is left for the future.

Before closing, it is important to notice that the aging observed is not restricted to the QSS region, but extends also to the following relaxation regime towards equilibration.

5 Conclusions

In this paper we have studied the microscopic dynamics towards relaxation in the HMF model, a Ham iltonian system of fully coupled inertial spins. Two di erent out-of-equilibrium initial conditions which lead to metastable longliving quasi-stationary states (QSS) have been studied. Our num erical results indicate that the dynam ics of the relaxation process is extremely rich and m ore complex than what previously thought.W e have found that for a certain class of initial conditions, i.e. M 1 IC, one can have metastable Q SS with non G aussian velocity pdfs which can be reproduced with truncated q-exponential curves and for which a nite size scaling seems to apply [13,14]. We have shown that in such a regime there are also strong velocity correlations in time which decay as a q-exponential. Moreover, con ming similar studies [18], in correspondence of this metastable states and in the following relaxation regime, aging in the velocity correlations has been found. Such a strong longterm memory e ect can be attributed to a glassy-like weak ergodicity-breaking due to a sort of dynam ical frustration which disappears at equilibrium. These new num erical facts, added to the anom alous di usion [21], the structures in angle and the weak mixing produced by a vanishing Lyapunov exponent [13,14], previously observed for this out-of-equilibrium regime, although do not constitute a rigorous proof, certainly provide with a strong indication that the nonextensive formalism proposed by T sallis could probably be applied also for Ham iltonian many-body systems, expecially when the in nite size limit is performed before in nite time limit. A rigorous link between the entropic index q and the dynam ical properties of nonextensive H am iltonian m any-body system s is still missing and must be found in order to con m de netely T sallis form alism. Such m etastable QSS provide an ideal benchm ark for testing the nonextensive theory, a project that we plan to continue in future works. We have also presented num erical evidence that when adopting a di erent class of out-of-equilibrium initial conditions, i.e. M 0 IC, one can have also m etastable QSS, but with no structures in the -space and a fast (almost exponential) decay of velocity correlations. In this case nonextensive statistics certainly

does not apply. Open questions, concerning aging for example, for this M 0 IC case still remain and will be studied in the future.

6 A cknow legm ents

We thank Celia Anteneodo and Constantino T sallis for stimulating discussions. One of us (A R.) would like to thank the Los A lam os N ational Laboratory for the nancial support.

References

- [1] C. Tsallis, J. Stat. Phys. 52, (1988) 479; for recent reviews see: C. Tsallis, Nonextensive Statistical Mechanics and Them odynamics, Lecture Notes in Physics, eds. S. Abe and Y. O kam oto, Springer, Berlin, (2001); Proceedings of NEXT2001, special issue of Physica A 305 (2002) eds. G. Kaniadakis, M. Lissia and A. Rapisarda; \Nonextensive Entropy- Interdisciplinary Applications", C. Tsallis and M. Gell-Mann eds., Oxford University Press (2003). See also http://tsallis.cat.dopfbr/biblio.htm for a regularly updated and complete bibliography.
- [2] A. Cho, Science 297 (2002) 1268; V. Latora, A. Rapisarda and A. Robledo, Science (2003) in press; S. Abe and A. K. Rajagopal, Science (2003) in press; A. Plastino, Science (2003) in press.
- [3] M L.Lyra and C.Tsallis, Phys.Rev.Lett. 80 (1998) 53; U.Timakli, C.Tsallis and M L.Lyra, Eur.Phys.J.B 10 (1999) 309; FABF.deMoura, U.Timakli and M L.Lyra Phys.Rev.E 62 (2000) 6361.
- [4] Latora, Baranger, Rapisarda and Tsallis, Phys. Lett. A 273 (2000) 97; U. Timakli, Phys. Rev. E 62 (2000) 7857 and Phys. Rev. E 66 (2002) 066212;
 U. Timakli U, C. Tsallis, M. L. Lyra, Phys. Rev. E 65 (2002) 036207; E P. Borges, C. Tsallis, G F. J. Ananos, P M C. de O liveira, Phys. Rev. Lett. 25 (2002) 254103.
- [5] F.Baldovin and A.Robledo Phys.Rev.E 66 (2002) 045104 (R) and Europhys. Lett. 60 (2002) 518.
- [6] C.Beck, G.S.Lew is and H.L.Swinney, Phys.Rev.E 63 (2001) 035303 (R). See also C.Beck, Physica A 295 (2001) 195, Phys.Lett.A 287 (2001) 240 and Phys.Rev.Lett. 87 (2001) 180201 and Physica A 305 (2002) 209; T.Arim itsu and N.Arim itsu, Physica A 305 (2002) 218.

- [7] E.G.D.Cohen, Physica A 305 (2002) 19.
- [8] C.Beck and E.G.D.Cohen Physica A 322 (2003) 267.
- B.M. Boghosian, Phys. Rev. E 53 (1996) 4754; M. Coraddu, M. Lissia, G. Mezzorani and P.Quarati, Physica A 305 (2002) 282
- [10] I.Bediaga, E M F.Curado, J.M iranda, Physica A 286 (2000) 156; G.W ilk and Z.W lodarczyk, Phys.Rev.Lett. 84 (2000) 2770; O.V.Utyuzh, G.W ilk G and Z.W lodarczyk J.ofPhys.G 26 (2000) L39; W M.Alberico, A.Lavagno and P. Quarati, Eur.Phys.J.C 12 (1999) 499.
- [11] L. Borland, Phys. Rev. Lett. 89 (2002) 098701
- [12] F. Baldovin C. T sallis and B. Schulze, Physica A 320 (2003) 184; F. Baldovin, E. Brigatti and C. T sallis [cond-m at/0302559]
- [13] V. Latora, A. Rapisarda and C. Tsallis, Phys. Rev. E 64 (2001) 056134 and Physica A 305 (2002) 129.
- [14] C. Tsallis, A. Rapisarda, V. Latora and F. Baldovin in "Dynamics and Thermodynamics of Systems with Long Range Interactions", T. Dauxois, S. Ru o, E. Arimondo, M. W ilkens Eds., Lecture Notes in Physics Vol. 602, Springer (2002) 140.
- [15] C.Anteneodo and C.Tsallis, Phys.Rev.Lett. 80 (1998) 5313; F.Tam arit and C.Anteneodo, Phys.Rev.Lett. 84 (2000) 208; A.Campa, A.Giansanti and D. Moroni, Phys.Rev.E 62 (2000) 303; Campa, A.Giansanti, D.Moroni and C. Tsallis, Phys.Lett.A 286 (2001) 251.
- [16] A. Campa, A. Giansanti and D. Moroni, Physica A 305 (2002) 137 and [condm at/0206173].
- [17] B J.C. Cabral and C. T sallis, Phys. Rev. E 66 (2002) 065101 (R).
- [18] M A. Montemurro, FA. Tamarit and C. Anteneodo, Phys. Rev. E 67 (2003) 031106.
- [19] F.D. Nobre, C. T sallis [cond-m at/0301492].
- [20] M. Antoni and S.Ru o, Phys. Rev. E 52 (1995) 2361. See also T. Dauxois, V. Latora, A. Rapisarda, S. Ru o and A. Torcini, in \Dynamics and Therm odynamics of System s with Long Range Interactions", T. Dauxois, S. Ru o, E. Arim ondo, M. W ilkens Eds., Lecture Notes in Physics Vol. 602, Springer (2002), p.458.
- [21] V.Latora, A.Rapisarda and S.Ru o, Phys. Rev. Lett. 80 (1998) 692; Physica
 D 280 (1999) 81 and Progr. Theor. Phys. Suppl. 139 (2000) 204.
- [22] V. Latora, A. Rapisarda and S. Ru o, Phys. Rev. Lett. 83 (1999) 2104 and Physica A 280 (2000) 81.
- [23] V. Latora and A. Rapisarda, Nucl. Phys. A 681 (2001) 331c.

- [24] M. Antoni, H. Hinrichsen and S. Ru o, Chaos Solitons and Fractals 13 (2002) 393.
- [25] M. Antoni, S. Ru o, A. Torcini Phys. Rev. E 66 (2002) 025103.
- [26] V. Latora and A. Rapisarda, Chaos Solitons and Fractals 13 (2002) 401.
- [27] D.H. Zanette and M.A. Montem urro, Phys. Rev. E 67 (2003) 031105.
- [28] Y.Y. Yam aguchi, [cond-m at/0209031].
- [29] V. Latora, A. Pluchino, A. Rapisarda and C. Tsallis, in preparation.
- [30] H.Yoshida, Phys. Lett. A 150 (1990) 262.
- [31] P. Hertel and W. Thirring, Ann. of Phys. 63 (1971) 520.
- [32] D. Lynden-Bell, Physica A 263 (1999) 293.
- [33] L.M ilanovic, H A Posch and W . Thirring, Phys. Rev. E 57 (1998) 2763.
- [34] D.H.E. Gross, M icrocanonical therm odynam ics: Phase transitions in Sm all systems, 66 Lectures Notes in Physics, W orld scientic, Singapore (2001) and refs. therein.
- [35] M. D'Agostino et al, Phys. Lett. B 473 (2000) 219.
- [36] M. Schmidt, R. Kusche, T. Hippler, J. Donges and W. Kronmueller, Phys. Rev. Lett. 86 (2001) 1191.
- [37] A. Torcini and M. Antoni, Phys. Rev. E 59 (1999) 2746.
- [38] V. Latora and M. Baranger, Phys. Rev. Lett. 82 (1999) 520.
- [39] P.G. Debenedetti and F.H. Stillinger, Nature 410 (2001) 259.
- [40] J.P. Bouchaud, L.F. Cugliandolo, J.Kurchan and M. Mezard Sping glasses and random elds, A.P. Young ed., World Scientic Singapore (1998).
- [41] L F. Cugliandolo, [cond-m at/0210312].
- [42] J.Kurchan, Phys. Rev. E 66 (2002) 017101.

Fig. 1. We show the tem poral behaviour of twice the average kinetic energy per particle for the two di erent initial conditions considered M 1 and M 0 and for the system sizes N = 500 (dashed lines) and N = 1000 (full lines). Ensemble averages over 1000 events were considered.

Fig.2. Com parison of twice the average kinetic perparticle (tem perature) considering two kind of averages: i) ensemble average (curves); ii) time and ensemble average (open circles). The latter has been used in previous papers to improve the statistics, while the former is the one adopted in the present paper. The curves plotted show that for ensemble averages over 1000 events the two procedures coincide. This comparison improves when considering larger sizes for which a smaller number of events is su cient in order to diminish the uctuations. See text for further details.

Fig. 3. Tem poral snapshots of the -space for the two initial conditions considered. We have structures in one case (M 1) which disappear after the plateau, while for the second initial condition (M 0) the phase space is always hom ogeneously lied. We put in red the rotators which have the highest velocities at time t = 0. This helps in order to understand the following dynam ical mixing. The plots refer to a single typical run for U = 0.69 and N = 10000.

Fig. 4. Pdf in angle vs time. This gure is a complement of the previous one and shows that, in the metastable QSS regime, we have the dynamical formation of small clusters in competition between each other for M 1 IC, but not for M 0 IC. However at equilibrium (t= 1000000) for both IC only one big rotating cluster remains, see text.

Fig. 5. T in e evolution of the entropy S for M 0 and M 1 initial conditions. See text for further details.

Fig. 6. Time evolution of the free energy F for M 0 and M 1 initial conditions. In correspondence of the metastable regime the free energy show a strong before going towards a minimum at equilibrium. See text for further details.

Fig. 7. Velocity pdfs in the QSS (t = 1000), top panel, and very close to equilibrium (t = 100000), bottom panel, for the two di erent initial conditions considered. The simulations refer to a size N = 10000. The equilibrium G aussian pdf is also plotted for com parison as full curve. In the QSS regime, one has a peaked pdf for M 1 IC and a at pdf for M 0 IC. In both cases strong deviations from the G aussian are evident. W hile the pdf for M 1 can be reproduced with a generalized and truncated T sallis pdf (see ref.[13]) this is not possible for M 0 IC.

Fig. 8. Velocity correlation functions, in the QSS regime, for the two intial conditions considered M 1 and M 0. Both of them can be reproduced by a q-exponential also shown as full (M 1 IC) and dashed (M 0 IC) curves. The corresponding param – eters are also reported. W e get q = 1.55 for M 1 IC and q = 1.12 for M 0 IC. Notice that in the latter case the value of q is close to 1 m eaning that the decay tends to be exponential. This is clearly seen in the inset where a linear-log scale was used and an exponential t is also reported for com parison. See text for further details.

Fig. 9. (a) W e plot for M 1 IC, N = 1000 and an ensemble average over 100 events the two-time autocorrelation function (7). Several waiting times are shown. (b) W e plot the curves in (a) by considering a scaling factor $(t_w)^{1=4}$. The nalpart of the tail is a power law. The scaled curves can be reproduced by a q-exponential with q = 1:65, A = 0:7 and = 60, full curve.

Fig. 10. (a) The same as g.9 but for M 0 IC. (b) The curves shown in (a) are scaled by the same factor $(t_w)^{1=4}$ used previously for M 1 IC. In this case, however the curves do not collapse into one and we get a faster decay.

This figure "f1.jpg" is available in "jpg" format from:

This figure "f2.jpg" is available in "jpg" format from:

This figure "f3p.jpg" is available in "jpg" format from:

This figure "f4.jpg" is available in "jpg" format from:

This figure "f5.jpg" is available in "jpg" format from:

This figure "f6.jpg" is available in "jpg" format from:

This figure "f7.jpg" is available in "jpg" format from:

This figure "f8.jpg" is available in "jpg" format from:

This figure "f9.jpg" is available in "jpg" format from:

This figure "f10.jpg" is available in "jpg" format from: