G round-state properties of nanographite system s with zigzag-shaped edges

Toshiya Hikihara and Xiao Hu

Computational Materials Science Center, National Institute for Materials Science, Tsukuba, Ibaragi 305-0047, Japan

Hsiu-Hau Lin^{1;2} and Chung-Yu Mou¹

¹ Department of Physics, National Tsing-Hua University, Hsinchu 300, Taiwan

² Physics D ivision, N ational Center for Theoretical Science, H sinchu 300, Taiwan

(Dated: April 14, 2024)

A -electron network in nanographite system s with zigzag edges exhibits strongly localized edge states, which are expected to have peculiar properties. We study elects of electron-electron interactions on ground-state properties of the system s by means of the weak-coupling renorm alization group and the density-matrix renorm alization-group method. It is shown that the ground state is a spin-singlet M ott insulator with nite charge and spin gaps. We also not that the edge states are robust against the electronic correlations, resulting in edge elective spins which can ip almost freely. The schematic picture for the low-energy physics of the system s is discussed.

PACS num bers: 71.10 H f, 73.22.-f, 75.75.+ a

I. IN TRODUCTION

Since the discoveries of fullerenes¹ and carbon nanotubes (CNTs),² the nature of \nanographites", graphite-based materials with nano-meter sizes, has attracted much attention from both fundamental science and applications. One of the most striking features of the materials is a wide variety of properties they show depending on their geometrical structure, i.e., size, shape, surface condition and so on. For example, CNTs can be eitherm etallic or semiconducting depending on the wrapping superlattice vector.³ Such a diversity of possible functions realized in nanographites makes them prom ising candidates for nano-scale devices.

Recently, it has been pointed out that a -electron system in nanographites with zigzag-shaped edges exhibits peculiar electronic states strongly localized around the edges, which are term ed \edge states".^{4,5,6} Such a state does not appear around arm chair-shaped edges. The nature of the edge states has been studied in the graphene sheet of nano-m eterw idth, nam ed the \nanographite ribbon" (NGR). Applying a tight-binding model for the NGR with zigzag edges, Fujita et al. have shown that the system has the electronic states localized on the edges, which penetrate from the edges into the bulk decaying exponentially.^{4,5} The edge states are believed to be responsible for a param agnetic behavior observed in activated carbon bers, which are taken to be an assembly of nanographite particles.^{7,8}

Since the edge states are characterized by almost at dispersions in a certain range of the momentum space, which result in a sharp peak in the density of state at the Ferm i energy, possible instabilities to various perturbations are in portant and have been investigated to date. C oncerning the Peierls instability due to electron-phonon interactions, it has been shown that no bond alternation occurs in the zigzag NGRs with realistic interaction strength.⁹ O n the other hand, e ects of electron-electron interactions on the edge states have been investigated by

a mean- eld approximation^{4,10} The studies have shown that an in nitesimal on-site interaction causes a spontaneous spin polarization at the zigzag edges. Similar results have been obtained for the zigzag CNTs by the density-functional theory.^{11,12} However, it is also known that the one-body approximation adopted in these approaches is not appropriate for one-dimensional (1D) quantum systems such as NGRs and CNTs. In fact, Lieb's theorem ¹³ prohibits the spontaneous spin polarization for the H ubbard m odel in the nanographite system s with zigzag edges. M ore detailed analyses with controlled approximations are therefore desirable for clarifying the low -energy properties of the nanographites.

In this paper, we study e ects of electron-electron interactions on the nanographite system swith zigzag edges using two powerful techniques: the weak-coupling renorm alization group (RG) and the density-m atrix renorm alization group (DMRG) method.14 These approaches allow us to treat the strong quantum uctuations in a controlled way for the 1D system s such as NGRs and CNTs. W ith the inclusion of electronic correlations, we show that the ground state of the zigzag NGR is a gapped spin-singlet and the arti cially broken symmetry of spin rotations in the mean-eld type analysis is restored by quantum uctuations. It is rather interesting that the edge states survive in the presence of electron-electron interactions and play a crucial role in the low-energy regime.

The rest of this paper is organized as follows. The model is presented in the next section. In Sec. III, we show results of the weak-coupling analysis. We rst discuss properties of the tight-binding model of the NGR system. Analytical eigen-wavefunctions and the corresponding energy spectrum are shown. And then, by keeping the low-lying edge states only and taking the electron-electron couplings into account, we derive the low-energy elective Ham iltonian in the continuum limit. It is shown that all coupling terms included in the Ham iltonian are relevant because of a large dynam ical exponent




FIG.1: Honeycom b lattice of carbon atom s in the zigzag NGRs and CNTs. The open circles and gray squares represent carbon sites of sublattice A and B, respectively.

of the dispersion of the edge states. The enhanced density of states at the Fermi energy generates nite charge and spin gaps, and consequently, the system exhibits a spin-singlet ground state without spontaneous symmetry breaking. The DMRG results are presented in Sec. IV. We nd that in the presence of the Hubbard interaction the ground state is a spin singlet with nite charge and spin gaps, and no spontaneous spin polarization appears. Besides, electrons around the zigzag edges correlate ferrom agnetically to compose e ective spins which are easily polarized by applying an external eld, while electrons in the bulk form a spin-singlet state which are hardly magnetized. The e ective spins are localized around the zigzag edges even for a rather strong coupling, showing that the localization character of the edge states persists robustly against the electron-electron interactions. The schem atic picture for the low energy physics of the system s is discussed. F inally, our results are sum m arized in Sec.V.

II. MODEL HAM ILTONIAN

In this paper, we consider a -electron system at halflling on a honeycom b lattice with zigzag edges as shown in Fig. 1. The e ect of electron-electron couplings is incorporated by introducing the on-site Hubbard interaction U. The Ham iltonian is

$$H = H_{0} + H_{U}; \qquad (1)$$

$$H_{0} = t(r;r^{0}) c^{V}(r)c(r^{0}) + hc;; (2)$$

$$H_{U} = U n_{*}(r)n_{\#}(r); \qquad (3)$$

where c (r) (c^{y} (r)) is the ferm ion annihilation (creation) operator at the site r = (x;y) and n (r) \dot{c} (r)c (r). The de nition of the site index is shown in Fig.1. The system size in the x- and y-directions are denoted by L_x and L_y , respectively. W ithin tight-binding approximation, the hopping amplitude $t(r; r^0)$ is assumed to be t between nearest-neighbor sites and otherwise 0. In the x-direction, the open boundary condition with L_x ! 1 is im posed for the zigzag NGR while the periodic boundary condition is im posed for the zigzag CNT. In the ydirection, the open boundary condition is im posed on the zigzag edges. W e set L_y to be even so that the system always has the rejection symmetry in the y-direction. All carbon sites at the edges are assumed to be term inated by hydrogen atom s.

W e note that the system is bipartite and the number of sites in each sublattice is equal, $N_A = N_B$. At the natural lling, with one -electron per site on average, the system is particle-hole symmetric. Thus, one can apply Lieb's theorem ¹³ to the system, which prohibits spontaneous spin polarizations in the ground state.

III. RENORMALIZATION GROUP

A. Band structure

To study the system in the weak-coupling lim it U t, it is natural to consider the case of U = 0 and analyze the band structure rst. We thus begin with diagonalizing the hopping H am iltonian H₀. Since the H am iltonian is translational invariant along the x-direction, we can perform the partial Fourier transform ation,¹⁵

$$c(k_{x};y) = \frac{1}{\frac{1}{L_{x}}} \exp[ik_{x}(x+)]c(x;y)$$
(4)

where = 1=2 for y = 0;1 (m od 4) and = 0 for y = 2;3 (m od 4). By the transform ation, the hopping H am iltonian H₀ is mapped into bond-alternating chains decoupled from each other,

where $t(y) = t_1$ 2tcos($k_x = 2$) if y is odd while $t(y) = t_2$ t if y is even. It is important to note that the e ective hopping t depends on the momentum k_x , and consequently, the motions in the x- and y-directions are entangled.

E ach of the bond-alternating chains is composed of two sublattices A (odd y) and B (even y). The hopping between these two sublattices lead to the coupled H arper equations,

where $y_A = 1;3;5;$:::; $L_y = 1$ are the lattice points of sublattice A. The open boundary condition in the ydirection requires $(L_y + 1) = 0$ and (0) = 0. From the coupled H arper equations, the wavefunctions of the eigenstates and the corresponding energy spectrum can be obtained in an analytical form. For the chains with $t_1 > t_2$ $(j_{x_x} j < 2 = 3)$, the wavefunction is given by

$$(k_{x}; y_{A}) \qquad \begin{array}{c} p_{y}(y_{A}) \\ p_{y}(y_{A}+1) \end{array} = \qquad \begin{array}{c} \sin \left[p_{y} y_{A} + '(p_{y}) \right] \\ \sin \left[p_{y}(y_{A}+1) \right] \end{array};$$

$$(8)$$

where the extra phase ', chosen to be in the range 0 ' $(p_y) < =2$, depends on the magnitude of momentum p_y ,

'
$$(p_y) = \tan^{-1} \frac{t_1}{t_1 + t_2} \tan p_y$$
: (9)

F in iteness of $L_{\rm y}$ causes the quantization of $p_{\rm y}$ which satis es the following constraint

$$(L_v + 1)p_v + '(p_v) = m$$
; (10)

where m is an integer. The energy spectrum for these states is given by

$$E (k_x; p_y) = \frac{q}{t_1^2 + t_2^2 + 2t_1t_2\cos(2p_y)} : (11)$$

Here we emphasize that the momentum in the ydirection, k_y , is no longer a good quantum number due to the open boundaries. However, the magnitude of the momentum p_y remains a good one and would be quite helpful for identifying dominating interactions later. Since the wavefunction Eq. (8) is extended into the bulk sites, we call the states \extended states".

For the chains with $t_1 < t_2$ ($jk_x j > 2 = 3$), peculiar localized states show up. One nds that there are only L_y 2 extended states in Eq. (8). The missing two states are linear combinations of localized states near the edges with complex momentum $k_y = =2$ i. The wavefunctions for these edge states are

$$(k_{x}; y_{A}) = e^{i y_{A} = 2} \qquad \sinh [(I_{y} + 1 y_{A})] \\ \sinh [(Y_{A} + 1)] \qquad : (12)$$

The imaginary part of the momentum ~~6~ 0 is the inverse of localization length of the edge states and satis es the constraint

$$(L_y + 1) = \tanh^1 \frac{t_1 + t_2}{t_1 \pm} \tanh :$$
 (13)

It is clear that the nonzero solution of $\$ only exists when $t_1 < t_2$. The energy spectrum for these edge states is given by

$$E(k_{x};) = \frac{q}{t_{1}^{2} + t_{2}^{2}} - 2t_{1}t_{2}\cosh(2): \quad (14)$$

We thereby obtain the whole band structure of the tight-binding Ham iltonian, Eqs. (11) and (14). The band




FIG.2: Band structure of the zigzag NGR with $\rm L_y$ = 12. The bold lines represent dispersions around the Ferm ipoint, k_x = .

structure for $L_y = 12$ is shown in Fig. 2, as an example. It is worth noticing that the energy spectrum of the edge states is always lower than those of the extended states. It means that the edge states would dominate the low-energy physics in the weak-coupling limit while the extended states in the bulk do not play a crucial role because allofthem are suppressed by a nite gap of order t= L_y .

B. Renorm alization group analysis

Now, we are ready to treat the interacting H am iltonian H $_{\rm U}$. A fter the partial Fourier transform ation, the interacting H am iltonian is written as,

$$H_{U}^{*} = \frac{U}{L_{x}} \frac{X}{f_{x_{1}gy=1}} (k_{x_{1}} k_{x_{2}} + k_{x_{3}} k_{x_{4}})$$
$$\frac{V}{f_{x_{1}gy=1}} (k_{x_{2}}; y)c_{\#}^{y} (k_{x_{3}}; y)c_{\#} (k_{x_{4}}; y): (15)$$

The delta function represents the momentum conservation in the x-direction. The Ham iltonian can be rewritten in terms of the eigenstates of H_0 by expanding the electron operator on the diagonal basis,

c
$$(k_x; y) = \sum_{p}^{X} (y)_{p} (k_x)$$
: (16)

where p_{p} is taken for all the extended states $p = p_{y}$ and, if ever, the edge states p =. The resultant H am iltonian is

$$H_{U}^{*} = \frac{U}{L_{x}} X X (k_{x_{1}} k_{x_{2}} + k_{x_{3}} k_{x_{4}})$$

$$X (k_{x_{1}} k_{x_{2}} + k_{x_{3}} k_{x_{4}})$$

$$y_{p_{1}}(y) y_{p_{2}}(y) y_{p_{3}}(y) y_{p_{4}}(y)$$

$$y_{p_{1}}^{y}(k_{x_{1}}) y_{p_{2}}(k_{x_{2}}) y_{p_{3}\#}(k_{x_{3}}) y_{p_{4}\#}(k_{x_{4}}):(17)$$

In general, pi are incom m ensurate and the dom inant contribution com es from the pairw ise equal pi vertices while other kinds of vertices only occur in a limited tiny phase space. This great reduction of dom inant interactions enables us to classify them into three categories. (i) The interactions involving only the edge states: Since the norm alized wavefunction of the edge states is order one near the edge and vanishingly small in the bulk, the wavefunction product of $p_i(y)$ in Eq. (17) after sum mation over y is order one. Thus the edge-edge interaction remains O (U). (ii) The interactions involving a pair of the extended states and a pair of the edge states: Due to the spreading of wavefunction in the extended state, the sum m ation of the wavefunction product in Eq. (17)is roughly j_{pv} (1) f 1=Ly. The edge-bulk interaction is therefore only O (U=L $_y$) and becomes smaller as the width of the NGR grows. (iii) The interactions involving only the extended states: From the sim ilar argum ent to the case of the edge-bulk interaction, the bulk-bulk interaction turns out to be 0 (U). The e ect of this interaction in the bulk system (L $_{\rm y}$! 1) has been studied in Refs. 15 and 16 by the weak-coupling analysis. It has turned out that the interaction opens a nite energy gap even if the noninteracting H am iltonian H_0 includes gapless chains with $k_x =$ 2 =3.

Let us move on to deriving the low-energy e ective H am iltonian in the continuum limit. In the weakcoupling limit, the e ective theory is dominated by the low-lying edge states near k_x because all other bands in the bulk are gapped. Thus, we may keep only the edge states and neglect all the gapped modes. That is to say, only the kinetic energy of the edge modes and the edgeedge interaction term must be considered in the limit. It is useful to construct the antibonding and bonding edge

(x) from the eld for the edge states de ned in Eq. (16),

_

$$(x) = \frac{dk_x}{2} e^{ik_x x} \qquad (k_x +); \qquad (18)$$

$$(k_x) = \frac{1}{\frac{p}{2}} [+ (k_x) (k_x)];$$
 (19)

where is a momentum cuto . Furthermore, we approximate the energy spectrum near k_x to the lowest non-vanishing order E (k) ' $c(k_x \ \tilde{f}, where the signs stand for the antibonding and bonding edge modes and the dynam ical exponent <math>z = L_y$ 1. The kinetic energy can then be expressed in terms of the elds (x) in the continuum limit

$$H_{0}^{2} = \frac{X}{dx} P_{p}^{Y}(x) (iQ_{x})^{z} (x); \quad (20)$$

where the constant c is set to unity for notation simplicity.

W hile it is rather straightforward to write down the kinetic term s in the form of Eq. (20), expressing the edgeedge interaction in term s of the eld operators is far from trivial. Our strategy to go from lattice to continuum lim it is to express the lattice operator for electrons in the edge states, e(x;y), in term softhe eld operator (k) and relate it to (x) by Fourier transform ation. To achieve the goal, we need to express e(x;y) in the eigenbasis of H'_0 ,

$$e(x;y) = \frac{Z}{2} \frac{dk_x}{2} e^{ik_x x} \sum_{p=1}^{X} p(y) p(k):$$
 (21)

Here a di culty com es from the entanglem ent of sum mation over k_x and P. Since depends on the longitudinal momentum k_x as in Eq. (13), the wavefunction $_{\rm P}$ (y) also has an implicit dependence on k_x . As a result, the lattice operatore (x;y) is not simply related to the continuous eld (k) by Fourier transform ation which requires the change of order between the sum mation over P and the integral over k_x . How ever, since the edge states with k_x ' are localized near the edges, it is reasonable to adapt the \sharp-edge" approximation which assumes that all the wavefunctions of the edge states take the simple form as for k_x = , i.e.,

$$(\mathbf{y}) = \frac{1}{2} (y_{;1} y_{;L_y}):$$
(22)

W ithin the sharp-edge approximation, the relation between the lattice and the eld operators becomes very simple,

$$e(x;1) = \frac{e^{1 x}}{2} [+ (x) + (x)];$$
 (23)

$$\varepsilon (x; L_y) = \frac{e^{1 x}}{p 2} [+ (x) (x)];$$
 (24)

$$e(x;y) = 0$$
 (2 y Ly 1): (25)

By substituting Eqs. (23) - (25) into the edge-edge interacting H am iltonian

$$H_{ee} = U \int_{r}^{X} e_{\pi}^{y} (x; y) e_{\pi} (x; y) e_{\#}^{y} (x; y) e_{\#} (x; y); \quad (26)$$

we obtain its continuum counterpart,

$$H'_{ee} = g_{c} (J_{+}^{2} + J^{2}) + g J_{+} J \qquad g J_{+} J$$

+ $\frac{g_{u}}{2} (I_{+} I^{y} + I I_{+}^{y}):$ (27)

Here the SU (2) invariant currents are de ned as

$$J_{P} = \frac{1}{2} \begin{array}{c} Y \\ P \end{array} \begin{array}{c} P \end{array} , \qquad (28)$$

$$J_{P} = \frac{1}{2} \frac{y}{P} P_{P} ; \qquad (29)$$

$$I_{P} = \frac{1}{2} \sum_{P}^{Y} P_{P} ; \qquad (30)$$

where and denotes Pauli matrices and an antisymmetric tensor, respectively. The bare values of these couplings generated by the on-site interaction U are $g_c = g = g = g_u = U$. We note that Eq. (27) is the most general form of the interacting H am iltonian which includes four-ferm ion interactions with preserving the U (1) SU (2) symmetry and the momentum conservation. In fact, one can use Eq. (27) to consider more complicated short-range interaction, which would generate a di erent set of bare couplings.

Finally, the low-energy eld theory of the zigzag NGR in the continuum limit is described by Eqs. (20) and (27). At the tree level, the scaling dimensions of the all four-ferm ion interactions are d(g) = z 1 by simple dimension counting. Since the dynamical exponent is greater than one for the NGR, all the couplings g_c ; g; g, and g_u are relevant under the RG transform ation. However, we note that relevance of these couplings does not necessarily warrant excitation gaps in the energy spectrum. To identify the ground state, it is necessary to apply another approach such as the bosonization technique. We will discuss it in the next subsection.

C. M ott insulating ground state

In this subsection, we consider what we can derive from the low-energy e ective H am iltonian obtained in the previous subsection. One clue is the fact that the e ective Ham iltonian has a form strikingly similar to the usual e ective theory for 1D system s except for the large dynam ical exponent z > 1. Hence, we discuss the analogy between the NGRs and a one-chain system to derive the ground-state properties of the NGR.On the other hand, the spectrum of the low-lying edge states in the NGR can be viewed as a two-chain system, whose bonding and antibonding bands touch the Ferm i energy at one Ferm i point as discussed below. Therefore, it is also useful to consider the NGR in the analogy with the two-chain system . W e will see that the both analyses give the same conclusion: the NGR has a spin-singlet ground state with nite charge and spin gaps.

Let us start with the stapproach, i.e., considering the NGR in the analogy to a one-chain system at halflling. In this analogy, the eld_+ () is regarded as a right- (left-) moving eld operator. Since the dispersion of the elds (x) is not linear, the rigorous m apping from the ferm ionic theory to the bosonic one is not possible. However, it is likely that the large dynam ical exponent z just serves to m ake all interactions relevant and we can still use the bosonization techniques to clarify physical properties of the ground states qualitatively. That is to say, we can treat the dynam ical exponent z = 1 + as aperturbation and still use the conventional bosonization rules to study the physical properties of the NGRs. Under the assumption, we follow conventional bosonization steps and introduce two pairs of bosonic elds

$$(x)' = e^{i[(x) + (x)]^{2}}; \qquad (31)$$

where (x) and (x) are respectively the phase and displacement uctuations, which obey the commutation

phaæ	(<u>2</u> ; <u>2</u>)	Cs	D
New	(;0)	0	0
CDW	(0;)	€ 0	0
D im er	(0;0)	0	€ 0

TABLE I: Com parison between di erent phases.

relation,

$$[(x); \circ (x^{0})] = 2 i[1 + sign(x + x)] \circ : (32)$$

Furtherm ore, we switch to the charge and spin basis dened by

$$(;;;) = \frac{1}{p-2}[(";") (";")]:$$
 (33)

The e ective interaction which is relevant to gap form ation in Eq. (27) can be written down in terms of these bosonic elds,

$$H_{ee}^{r} = g \cos(\frac{p}{2}) \quad g_{i} \cos(\frac{p}{2}): \quad (34)$$

The absence of their dual elds and in the interactions comes from the charge and spin conservations. For the on-site interaction, both g and g_u are positive and grow under RG transformation. Consequently, the corresponding bosonic elds are pinned at speci c values up to uctuations,

$$p = \frac{p}{2}$$
, $p = \frac{p}{2}$, 0: (35)

The signs of the couplings are of crucial in portance here. Expanding the cosine term around the pinned values, the system acquires nite gaps in both charge and spin sectors. Therefore, the ground state is a M ott insulator with both charge and spin gaps. The form ation of the charge gap is not surprising because the electron density is com m ensurate with the underlying lattice structure. The

nite spin gap indicates that the spin rotational sym metry is not broken spontaneously and the ground state is a spin singlet, as predicted by Lieb's theorem. The result di ers from the mean- eld theory predictions where the quantum uctuations are ignored. We believe that the strong uctuations in 1D systems would invalidate the mean- eld approach and restore the sym metry which is broken at the mean- eld level.

It is worth noting that for the usual weakly-coupled 1D system with linear energy spectrum, the bosonic elds never get pinned at the values listed in Eq. (35) because it is not stable in the RG ow. However, the new phase here is stable due to the large dynam ical exponent which corresponds to the van Hove singularity in the density of states. We summarize the pinned values for each phase in Table I. To di erentiate this new phase from the familiar charge density wave or the dimer phase, one needs to compute the order parameters, i.e., charge density modulation C_s and bond dimerization D de ned as

 C_{s} (1^x)ha^y(x)a (x)i

'
$$h_{+}^{y}$$
 i+ h_{-}^{y} i; (36)
D $\frac{(1)^{x}}{2}ha^{y}(x)a(x+1) + a^{y}(x+1)a(x)i$
' ih_{+}^{y} i+ ih_{-}^{y} i; (37)

where a (x) is the lattice operator in the one-chain system and related to the elds as a (x) ' $e^{ik_F x}$, (x) + $e^{ik_F x}$ (x) with $k_F = =2.0$ sing the relations

$$Y_{1}, V_{2}, V_{3}$$
 is $i = \frac{p}{2}e^{i} = \frac{p}{2};$ (38)

$$Y_{+\#}, ie^{i} = \sqrt{2}e^{+i} = \sqrt{2}i;$$
(39)

and substituting the pinned values of the bosonic edds, one obtain the expectation values of the order param eters. The results are listed in Table I. It is clear that there is no long-range order or symmetry breaking for the ground state of the zigzag NGRs.

Now we take another perspective and view the NGR as a system of two chains coupled via an interchain hopping. Figure 3 (a) shows the band structure of the two-chain system for the special point $t_{rung} = t_{leg} = 2$, where t_{rung} (t_{leg}) is an amplitude of the interchain (intrachain) hopping. As the hoppings are tuned at the special value, the chem ical potential cut through the bonding (antibonding) band at single Ferm i point $k_x = (k_x = 0)$ with vanishing Ferm i velocity v_{F} ! 0, rather than two pairs of Ferm ipoints in general. On the other hand, both the bonding and antibonding bands of the low -lying edge states in the NGRs have single Ferm i point $k_x = w$ ith $v_{\rm F}$! 0. Hence, in the low-energy limit, the peculiar band structure in the NGRs can be viewed as a two-chain system with shifting the momentum of the antibonding [See Fig. 2 and Fig. 3 (b)]. One eld $k_x ! k_x +$ m ight worry that the dynam ical exponent in the NGRs $z = L_y$ 1 may be di erent from that in the two-chain system z = 2. However, the dynamical exponent z in this limit only a ects how the Fermi velocity approaches zero and should not give rise to qualitative di erence.

In the weak-coupling lim it, the lattice electron operator a_i (x) in the i-th chain of the two-chain system can be described by two pairs of chiral elds

$$a_1 (x) = \frac{e^{i x}}{2} [+ (x) + e^{iQ x} (x)];$$
 (40)

$$a_2(x) = \frac{e^{1/x}}{\frac{p}{2}} [+(x) e^{iQ x} (x)];$$
 (41)

where Q = is the momentum di erence between Ferm i points of bonding and antibonding bands in the two-chain system. The relation of lattice operators to the eld operators for the two-chain system in Eqs. (40) and (41) are strikingly similar to those for the NGR in Eqs. (23) and (24), except the momentum shift Q. This enables us to translate the excitations and correlations in the twochain system directly into those for the NGR s.

For on-site repulsive interaction, the ground state of the two-chain system is the Mott insulator with nite (a)

FIG. 3: Schematic picture of the band structure of (a) the two-chain system with $t_{rung}=t_{leg}$! 2, and (b) the same system with them on entum shift Q = for the antibonding eld.The bold lines represent dispersions around Ferm ipoints.

charge and spin gaps. In addition, there exists neither broken symmetry nor local order parameter. This Mottinsulating phase resembles the peculiar phase we discussed above by making analogy to the one-chain case. Thus, both approaches support that the ground state of the NGRs is a spin singlet with nite charge and spin gaps.

The two-chain analogy also gives us inform ation about the spin-spin correlation. It is known that the two-chain system has magnon excitations with spin S = 1 and momentum P = (;). The momentum shift Q in the x direction shifts the low-lying magnon to P = (0;). This means that the spin-spin correlation functions in the NGRs are ferror agnetic within the same edge and antiferror agnetic between opposite edges. This result is consistent with the fact that the sites in the sam e edge belong to the same sublattice while those in opposite edges belong to di erent sublattices.

IV . NUMERICAL RESULTS

In this section, we present num erical results for the – electron system (1) at half–lling. Hereafter, we set t = 1. U sing the DM RG m ethod with in proved algorithm,^{14,17} we have calculated various energy gaps and correlation functions. The num ber of kept states is up to m = 1200 per block. Num erical errors due to the truncation of the DM RG calculation are estim ated from the di erence between the data with di erent m 's. W e discuss the results for the zigzag NGR s and the zigzag CNT s in the follow ing subsections.

A. Zigzag nanographite ribbons

In this subsection, we show results on the zigzag NGRs. We rst discuss the results for $L_y = 4$. In all the cases calculated, we have found that the ground state belongs to the subspace of M = 0, where M is the totalm agnetization in the system. In Fig. 4, we show the data of the




FIG. 4: L_x -dependence of the energy gaps in the NGRs for U = 1; (a) the charge gap _c, (b) the spin gap ⁽¹⁾_s. The solid and open circles represent the data for $L_y = 4$ and $L_y = 6$, respectively. The numerical errors due to the DMRG truncation are smaller than symbols. The gaps for $L_y = 4$ are extrapolated by tting the data to a polynomial form, $(L_x) = (1) + a = L_x + b = L_x^2$ where a and b are tting parameters. (c) The U-dependence of the extrapolated gaps for $L_y = 4$.

charge and spin gaps de ned as

$$c = \frac{1}{2} E_{0} \frac{N}{2} + 1; \frac{N}{2} + E_{0} \frac{N}{2} - 1; \frac{N}{2}$$

$$2E_{0} \frac{N}{2}; \frac{N}{2} ;$$

$$(42)$$

$${}_{s}^{(M)} = E_{0} \frac{N}{2} + M; \frac{N}{2} M E_{0} \frac{N}{2}; \frac{N}{2}; (43)$$

where $E_0 (N_*; N_*)$ is the lowest energy in the subspace of N * up-spin and N * down-spin electrons and N is the total number of sites. For $L_y = 4$, both the energy gaps are extrapolated to non-zero values at $L_x ! 1$. The U dependence of the extrapolated gaps suggests that both of the gaps start to open from in nitesim al U, i.e., $U_c =$ 0. We thus conclude that for arbitrary U > 0 the ground state of the zigzag NGRs is a spin singlet with nite charge and spin gaps, being consistent with the weakcoupling analysis.

To see how the spin polarizations are distributed in real space, we have calculated the local spin polarization

$$hS^{z}(r)i_{M} = \frac{1}{2}hn_{*}(r) n_{\#}(r)i_{M};$$
 (44)

where h_{M} denotes the expectation value in the low estenergy state in the subspace of M. We have found that in




FIG.5: Distribution of local spin polarization $hS^{z}(r)i_{M=1}$ in the NGRs with (a) $L_{y} = 4$ and (b) $L_{y} = 6$. The Hubbard coupling and the total magnetization are U = 1 and M = 1. Open and gray circles respectively represent the positive and negative values of the spin polarization while the areas of them are proportional to the magnitude.

the ground state the local spin polarization hS^z (r) $i_{M} = 0$ is zero at all sites within the num erical error, supporting that the ground state is spin singlet. There is no spontaneous spin polarization. Figure 5 (a) shows the distribution of the local spin polarization in the magnetized state of M = 1 for U = 1 and $L_y = 4$. As seen in the gure, the magnetization appears alm ost only at the edge sites. This suggests that the electrons in the edge states are polarized by an external eld more easily than those in the bulk. We note that the localization character of the magnetization seem s to persist even for a rather large coupling U = 4. In other words, the edge states survive robustly against the electron-electron interactions even if U becom es larger than the hopping am plitude t.

In Fig.6, we show the spin-spin correlation functions in the edge sites,

$$hS^{z}(x;1)S^{z}(x^{0};1)i_{M=0}$$
: (45)

To reduce boundary e ects, the correlations are calculated between the sites of the same distance from the center of the NGR. The correlations turn out to be always positive, suggesting a ferrom agnetic coupling between the sites in the same edge. We have also found that the spin correlation between the sites in opposite edges is always negative, i.e., antiferrom agnetic. These behaviors are consistent with the expectation from the two-chain analogy discussed above. The correlations decay exponentially for all U's re ecting the nite spin gap⁹, and are enhanced as U increases.

Finally, we discuss the dependence of low-energy properties of the zigzag NGRs on the width L_y . Since the low-energy physics of the NGRs is dominated by the edge states which are strongly localized around the zigzag edges, it is likely that the singlet-triplet spin gap $s^{(1)}$, which corresponds to the energy scale required to m agnetize the edge spins, com es from the antiferrom agnetic




FIG. 6: G round-state spin-spin correlation functions in the NGRs with $L_y = 4$ and $L_x = 20$. The circle, square, and triangle represent the data for U = 1, 2, and 4, respectively. The error bars represent the num erical error due to the DMRG truncation.

e ective coupling between the electrons in the edge states of opposite sides. Thus, it is natural to expect that the spin gap decreases as the width L_y increases. To see this, we have perform ed calculations for the NGR with $L_y = 6.^{20}$ As can be seen in Fig. 5 (b), the spin polarizations in the magnetized state of M = 1 are found to be strongly localized at the zigzag edges, suggesting that the edge states exist also for $L_y = 6$. Furtherm ore, we can clearly see in Fig. 4 that the spin gap for the same L_x decreases as the width of the ribbon L_y increases, as expected. Unfortunately, the system size calculated for $L_y = 6$ is not large enough to obtain the extrapolated gap $s^{(1)}$ ($L_x \ 1 \ 1$). Calculations for larger system s are desirable to clarify the situation for the NGR sw ith larger L_y .

B. Zigzag nanotubes with open edges

For the zigzag CNTs with open ends, we can apply basically the same argument as that for the zigzag NGRs. The dimension between the two systems is merely the fact that in the CNTs them omentum along the x-direction is quantized as $k_{\rm x}=2$ n=L $_{\rm x}$ due to the nite circum ference L $_{\rm x}$. To see what happens, we have done calculations for the zigzag CNT with L $_{\rm x}=2$. In this case, there is one edge state at each zigzag open end. A lthough the width of the CNT is quite narrow, we will show that characteristic magnetic properties expected for general zigzag CNTs already appear.

W e have found that the ground state always belongs to the subspace of M = 0 for all values of U > 0. In Fig. 7, we show the data of the charge gap $_{\rm c}$ and the spin gaps $_{\rm s}^{(1)}$ and $_{\rm s}^{(2)}$. It can be seen in the gure that all the gaps take nonzero values in all the case of U > 0 and L_y calculated. We therefore conclude that the ground state of the zigzag CNTs with nite L_y is a spin-singlet M ott




FIG.7: Charge gap $_{\rm c}$ and spin gaps $_{\rm s}^{(1)}$ and $_{\rm s}^{(2)}$ of the CNTs with $L_{\rm x} = 2$ for U = 1;2, and 4 as functions of the length of CNTs $L_{\rm y}$. The num erical errors due to the DMRG truncation are smaller than symbols. The dotted lines are guides for eye.

insulator, same as the NGRs. An interesting observation here is that the spin gap $^{(1)}_{s}$ decays exponentially as L_y increases while $^{(2)}_{s}$ converges to nite values at L_y ! 1. Hence, for large enough L_y , the magnetization M = 1 can be induced by applying an in nites in all eld whereas a nite eld is needed to magnetize the system to M = 2.

Next, we discuss how the spin polarizations are distributed in real space. Same as the NGRs, the local spin polarization in the ground state is zero at all sites. Figure 8 shows the distribution of the local spin polarization hS^z (r)i_M in magnetized states for U = 1. The data clearly show that in the state of M = 1 the induced magnetization are strongly localized at the ends of the zigzag CNT. This demonstrates the robustness of the localization character of the edge states against the coupling U. In the state of M = 2, on the other hand, the distribution of hS^z (r)i_M = 2 seems to be a superposition of the edge magnetization and magnetization excited in the bulk sites. We have found sim ilar distributions of hS^z (r)i_M for U = 2 and 4.

The results on the spin gaps and the local spin polarizations lead us to the conclusion that there are two di erent energy scales in magnetic excitations in zigzag CNTs: the energy to magnetize electrons in the edge states and the one to magnetize the bulk electrons. As the length of the CNT increases, the form er decays exponentially while the latter converges to a nite value, the spin gap of the bulk system . This suggests that one can regard the edge states and the bulk electrons as two alm ost independent objects; The edge states are essentially decoupled from the bulk singlet state in spite of the presence of U ' O (t). Very recently, it has been pointed out from a weak-coupling analysis that thicker CNTs including m ore than one edge states at each open ends also exhibit such a decoupling of the edge and bulk states.²¹ W e therefore expect that the conclusion above is valid




FIG.8: Distribution of local spin polarization in the CNTs with $L_x = 2$ for U = 1 and $L_y = 14$. The totalm agnetization is (a) M = 1 and (b) M = 2.0 pen and gray circles respectively represent the positive and negative values of the spin polarization while the areas of them are proportional to the m agnitude.

for general CNTs with zigzag open edges.

C . Picture of the low -energy states

From the above results, we can deduce a schematic picture to represent the low-energy physics of the zigzag NGRs and CNTs. In this picture, the -electron system consists of two parts: the electrons in the bulk form ing a spin-singlet state and those in the edge states at each zigzag edge. The electrons in the same edge are correlated ferrom agnetically with each other to compose a large e ective spin. The e ective spins interact via an effective antiferrom agnetic coupling across the bulk singlet state. Hence, the ground state of the system is a spin singlet in total. The e ective coupling becomes smaller as the distance between the e ective spins becom es larger, and nally, when the distance becom es large enough, the e ective spins can ip freely, giving a param agnetic response. The bulk electrons remain form ing a spin singlet unless a eld corresponding to the bulk spin gap is applied. We note that the picture above is basically consistent with the one discussed in Refs. 10 and 22.

V. DISCUSSION

In this paper, we have studied low -energy properties of nanographite system swith zigzag edges in the presence of on-site H ubbard interactions U using the weak-coupling RG and the DMRG method. We have analyzed the H ubbard model on the zigzag NGRs and the zigzag CNTs. We nd that in both systems the ground state for U > 0 is a spin-singlet M ott insulator with nite charge and spin gaps. It is also found that the localization property of the edge state persists even for a rather large value of U, resulting in the e ective spins localized around the zigzag edges.

Finally we wish to touch upon further extensions of this study. One issue to be studied is e ects of further hoppings and long-range electron-electron interactions. As for magnetism, these long-range terms violate the assumptions of Lieb's theorem and open the possibility of exotic magnetic states. However, it is also expected that the spin gap observed in the present study tends to stabilize the spin-singlet ground state against perturbations. Very recently, it has been pointed out by the density-functional theory¹² that zigzag CNTs with a

nite length can exhibit a high-spin ground state depending on the circum ference L_x , although the method may underestimate e ects of strong quantum uctuation in the nanographite system s. To clarify the situation, analyses with controlled approxim ations are needed. A nother issue is stacking e ects. Nanographite materials often take a stacking form, e.g., nanographite particles com posed by a few graphene sheets and multiwall CNTs. Hence, it is important for applications to study e ects of interlayer interactions. For nanographite particles, it has been shown that their magnetic properties change drastically depending on types of stacking geometry.²³ In multiwall CNTs, the di erence of radius and chirality between inner and outer layer is expected to a ect not only the edge states but also the bulk properties in a com plicated way. Further extensive studies are desirable to tack le the issue.

A cknow ledgm ents

W e are gratefulto K.W akabayashiand A.Tanaka for fruitful discussions. TH thanks S.O kada for sending a preprint before publication. TH and XH are supported by the M inistry of Education, Culture, Sports, Science and Technology, Japan, under the Priority G rant No. 14038240. HHL and CYM appreciate the nancial support from N ational Science Council in Taiwan through grant Nos. 91-2112-M 007-040 (HHL), 91-2120-M -007-001 (HHL) and 91-2112-M 007-049 (CYM).

¹ H.K roto et al., N ature 318, 162 (1985).

² S. Lijim a, Nature 354, 56 (1991).

³ For a review, see R. Saito, G. D resselhaus and M. S. D resselhaus, Physical Properties of Carbon N anotubes (Im pe-

rial College Press, 1998).

- ⁴ M.Fujita, K.W akabayashi, K.N akada and K.Kusakabe, J.Phys.Soc.Jpn.65, 1920 (1996).
- ⁵ K.Nakada, M.Fujita, G.D resselhaus and M.S.D resselhaus, Phys. Rev. B 54, 17954 (1996).
- ⁶ K.W akabayashi, M.Fujita, H.A jikiand M.Sigrist, Phys. Rev.B 59, 8271 (1999).
- ⁷ Y.Shibayam a, H.Sato, T.Enokiand M.Endo, Phys. Rev. Lett. 84, 1744 (2000).
- ⁸ Y.Shibayama, H.Sato, T.Enoki, X-X.Bi, M.S.D resselhaus and M.Endo, J.Phys. soc. Jpn. 69, 754 (2000).
- ⁹ M.Fujita, M.Igam iand K.Nakada, J.Phys.Soc.Jpn.66, 1864 (1997).
- ¹⁰ K.W akabayashi, M. Sigrist and M. Fujita, J. Phys. Soc. Jpn. 67, 2089 (1998).
- ¹¹ S. O kada and A. O shiyam a, Phys. Rev. Lett. 87, 146803 (2001).
- $^{\rm 12}\,$ S.O kada and A.O shiyam a, unpublished.
- ¹³ E.H.Lieb, Phys. Rev. Lett., 62, 1201 (1989).
- ¹⁴ S.R.W hite, Phys. Rev. Lett. 69, 2863 (1992); Phys. Rev.

- в 48,10345 (1993).
- ¹⁵ H.H.Lin, Phys. Rev. B 58, 4963 (1998).
- ¹⁶ L.Balents and M PA.Fisher, Phys.Rev.B 55, R11973 (1997).
- ¹⁷ S.R.W hite, Phys. Rev. Lett. 77, 3633 (1996).
- ¹⁸ T.Kennedy, J.Phys.Condens.Matter 2, 5737 (1990).
- ¹⁹ W e note that the suppression of the correlation functions from the linear behavior seen for large r L_x is due to the presence of the open boundaries and is not essential.
- For the NGR with L_y = 6, we rem ove the boundary sites (x;y) = (1;L_y) and (L_x;1), which are connected to other site by only one bond, to avoid undesirable boundary effects. The shape of the system calculated is shown in Fig. 5 (b).
- $^{21}\,$ S.Ryu and Y.Hatsugai, cond-m at/0211008.
- ²² K.Nakada, M.Igam iand M.Fujita, J.Phys.Soc.Jpn.67, 2388 (1998).
- ²³ K.Harigaya, Chem.Phys.Lett. 340, 123 (2001).