G row ing Surfaces with A nom alous D i usion -R esults for the Fractal K ardar-P arisi-Zhang E quation

Eytan Katzav

School of Physics and Astronom y, Raym ond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel

In this paper I study a model for a growing surface in the presence of anom alous di usion, also known as the Fractal K ardar-Parisi-Zhang equation (FKPZ). This equation includes a fractional Laplacian that accounts for the possibility that surface transport is caused by a hopping mechanism of a Levy ight. It is shown that for a speci c choice of parameters of the FKPZ equation, the equation can be solved exactly in one dimension, so that all the critical exponents, which describe the surface that grows under FKPZ, can be derived for that case. A flerwards, the Self-C onsistent E xpansion (SCE) is used to predict the critical exponents for the FKPZ model for any choice of the parameters and any spatial dimension. It is then veried that the results obtained using SCE recover the exact result in one dimension. At the end a simple picture for the behavior of the Fractal KPZ equation is suggested, and the upper critical dimension of this model is discussed.

The K ardar-Parisi-Zhang (KPZ) equation [1] for surface grow th under ballistic deposition was introduced as an extension of the Edwards-W ilkinson theory [2]. The interest in the KPZ equation exceeds far beyond the interest in evolving surfaces because of the following reasons: (a) The KPZ system is known to be equivalent to a number of very di erent physical system s. Exam ples are the directed polymer in a random medium, Schrodinger equation (in imaginary time) for a particle in the presence of a potential that is random in space and time and the important Burgers equation from hydrodynamics [2]. (b) The second reason, that is more important, to my mind, is that it serves as a relatively simple prototype of non-linear stochastic eld equations that are so common in condensed matter physics.

The equation for the height of the surface at the point r and time t, h (r;t), is given by

$$\frac{\partial h}{\partial t} = r^{2}h + \frac{1}{2}(rh)^{2} + (r;t); \qquad (1)$$

where autom atically the constant deposition rate is removed and (r;t) is a noise term such that

h
$$(x;t)i = 0$$

h $(x;t)$ $(x';t)i = 2D_0$ $(x x')$ $(t t');$ (2)

As can be seen in eq. (1) the basic relaxation mechanism in the KPZ equation is a Laplacian term that results from nearest neighbor hopping in the growing surface. During the last years, there has been a growing interest in other relaxational mechanisms [22]-[25], namely subdi usive di usion that seems to appear in the context of charge transport in am orphous sem iconductors [26, 27], NMR di usometry in disordered materials [28], and the dynamics of a bead in polymer networks [29]. The theoreticale ort to account for such phenomena led to the

formulation of the celebrated fractional Fokker-Planck equation (FFPE) [22]. This equation includes a fractional (Riem ann-Liouville) operator instead of the standard derivative of the Fokker-P lanck equation. Recently, M ann and W oyczynski [3] have suggested that in order to account for experim ental data, namely experim ents in which impurities were present on the growing surface [4], a modi cation of the KPZ equation has to be considered. They used the observation that the presence of an im purity can act as a strong trap for an adatom migrating at room tem perature, to conjecture that this process corresponds to Levy ights between trap sites. This conjecture then served as a justi cation for the introduction of a fractional Laplacian into the continuum equation of the growing surface as another relaxation mechanism . A ctually, the fractional Laplacian dom inates the standard Laplacian in the KPZ equation in the scaling regime (i.e. in the large scale lim it), so that the standard Laplacian can be ignored from the beginning. To summarize this exposition and to be more speci c, the equation they eventually suggest to describe the growing surface in the presence of self-sim ilar hopping surface di usion is the fractalKPZ (FKPZ) equation given by

$$\frac{\partial h}{\partial t} = (h) + \frac{1}{2} (r h)^2 + (r;t); \qquad (3)$$

where () $^{/2}$ (or in Fourier space (F) (q)

 $\dot{r}_{j}\dot{r}_{j}^{2}$ F (q)) is the fractional Laplacian, and in our context it is more convenient to choose = 2 with 0 < 2 (where the special case = 0 corresponds to the standard KPZ equation). In addition, (r;t) is a noise characterized by

h
$$(\mathbf{r};t)$$
 $(\mathbf{r}';t)i = 2D_0 \mathbf{j}\mathbf{r} \mathbf{r}^0 \mathbf{j}^2$ d $(t t^0):$ (4)

A ctually M ann and W oyczynski $[\beta]$ discussed the speci c case of white noise that corresponds to = 0 in the last equation, but since the m ore general case does not require speciale orts we discuss the FKPZ problem with spatially correlated noise. Furtherm ore, an exact solution

E lectronic address: eytak@ post.tau.ac.il

is possible only for a special case with correlated noise, so that the m ore general discussion is also interesting on that basis.

The proposed FKPZ equation generalizes the FFPE equation m entioned above in that it is a eld equation, rather than an equation for a single degree of freedom. Therefore, it is understood that such a generalization is essential in order to account for the dynam ics of a whole m edium experiencing anom alous di usion, and not just an arti cial problem. O byiously, the technical m athem atical di culties to be overcom e in this nonlinear case are form idable in com parison to those for the linear fractal kinetic equations.

However, in their paper M ann and W oyczynski [3] were not able to predict the critical exponents that describe the surface that grows under FKPZ.But before I m ake any new statements about this model let me brie y sum marize the various quantities of interest.

A very important quantity of interest is the roughness exponent that characterizes the surface in steady state. The roughness exponent is usually de ned using W (L;t) the roughness of the surface (that is de ned as the RM S of the height function h (r;t), in a system of linear size L). Then, in term s of W (L;t), is given by

$$W (L;t) / L :$$
 (5)

A nother in portant quantity of interest is the growth exponent that describes the short time behavior of the roughness W (L;t) (with at initial conditions)

$$W (L;t) / t:$$
 (6)

Finally, I introduce the dynam ic exponent z that describes the typical relaxation time scale of the system (i.e. the dependence of the equilibration time on the size of the system)

$$t_x / L^z$$
: (7)

It is well known [2] that these three exponents are not independent and that under very general considerations one should expect the following scaling relation

$$z = /$$
 (8)

(this relation is a direct consequence of the Fam ily-V icsek scaling relation [5].

In addition, for the K P Z equation there is another scaling relation, that comes from a sym metry of the equation under in nitesim altilting of the surface (this sym metry is just the fam ous G alilean invariance of the B urgers equation) [2]

$$+ z = 2$$
: (9)

It can easily be checked that this sym m etry holds in the case of FK P Z as well, because the fractal B urgers equation is evidently invariant under G alilean transform ation (see ref. [3], eq. (7.8)). Therefore, the last scaling relation is also relevant in the this discussion. Hence both

scaling relations, reduce the number of unknown exponents to just one (out of the three we started with). In some cases an extra scaling relation is possible (for example, in the case of the KPZ equation with long-range noise – see refs. [9, 32, 33]), so that the exponents can be obtained exactly by power-counting. Employing the term inology of the D ynam ic R enorm alization G roup (DRG) approach this can be explained by saying that certain terms in the dynam ic action do not renorm alize, and so an extra scaling relation arises (see [35]). This kind of solutions are naturally available also in the present problem. W henever the exponents are obtained due to such an extra condition it will be speci cally pointed out.

In this paper I show that for a speci c choice of parameters of the FKPZ equation (namely = 2). the equation can be solved exactly in one dimension, so that all the critical exponents can be derived easily for that case. A flerwards, in order to give a more complete picture (i.e. for any dimension d, and any spatial correlation index) I apply a method developed by Schwartz and Edwards [6]-[8] (also known as the Self-Consistent-Expansion (SCE) approach). This method has been previously applied successfully to the KPZ equation. The method gained much credit by being able to give a sensible prediction for the KPZ critical exponents in the strong coupling phase, where m any Renorm alization-G roup (RG) approaches failed, as well as D ynam ic R enorm alization G roup (D RG) [1, 9] (actually, it can be shown that the strong coupling regime is inaccessible by DRG even when it is used to all orders [30, 31]). It is then veri ed that the results obtained using SCE recover the exact result in one dimension.

As mentioned above, for the speci c case = 2 in one dimension an exact solution can be found for the FKPZ problem using the Fokker-Planck equation associated with its Langevin form (i.e. eq. (3)). This particular choice of parameters corresponds to a situation where the fractional exponent equals the exponent 2 that describes the decay of spatial correlations in the noise. Since kind this exact solutions is familiar in the KPZ community I will simply state the nal results given by = 1=2 and z = 3=2 (using the scaling relation (9)). Notice that these critical exponents for non zero 's and 's, as the classical KPZ case corresponds to = 2 = 0.

Now, the Self-Consistent Expansion (SCE) is applied in order to learn about the behavior of this system in more general contexts (other dimensions and cases where

62). SCE's starting point is the Fokker-P lanck form of FKPZ, from which it constructs a self-consistent expansion of the distribution of the eld concerned.

The expansion is formulated in terms of $_q$ and $!_q$, where $_q$ is the two-point function in momentum space, de ned by $_q = hh_q h_q i_s$, (the subscript S denotes steady state averaging), and $!_q$ is the characteristic frequency associated with h_q . It is expected that for sm all enough q, $_q$ and $!_q$ are power laws in q, $_{q} = Aq$ and $!_{q} = Bq^{z}$ (10)

where z is just the dynamic exponent, and the exponent is related to the roughness exponent by

$$=\frac{d}{2}$$
: (11)

The main idea is to write the Fokker-Planck equation $(P/(et = OP) in the form (P/(et = D_0 + O_1 + O_2)P), where O_0 is to be considered zero order in some parameter, O_1 is rst order and O_2 is second order. The evolution operator O_0 is chosen_ito have a simple form <math>O_0 = \frac{P}{q} \frac{e}{eh_q} D_q \frac{e}{eh_q} + !_q h_q$, where $D_q/!_q = q$. Note that at present q and $!_q$ are not known. I obtain next an equation for the two-point function. The expansion has the form $q = q + c_q$ (f $_pg; f!_pg$), because the low est order in the expansion already yields the unknow n q. In the same way an expansion for $!_q$ is also obtained in the form $!_q = !_q + d_q$ (f $_pg; f!_pg$). Now, the two-point function and the characteristic frequency are thus determ ined by the two coupled equations

$$c_q (f_p g; f!_p g) = 0$$
 and $d_q (f_p g; f!_p g) = 0$ (12)

These equations can be solved exactly in the asymptotic lim it to yield the required scaling exponents governing the steady state behavior and the time evolution. W orking to second order in the expansion, one gets the two coupled integral equations

$$D_{q} = \frac{X_{q} + 2 X_{m}}{Y_{m} + 2 Y_{q} + 1 + 1 + 1} \frac{M_{q} + M_{m} + 2 Y_{m}}{M_{q} + 1 + 1 + 1} = 2 \frac{X_{m}}{Y_{m}} \frac{M_{q} + M_{m} + M_{m} + 2 H_{m}}{M_{q} + 1 + 1 + 1} = 0; \qquad (13)$$

and

$$q \quad !_{q} \quad 2 \stackrel{X}{\underset{m}{}} M_{q'm} \frac{M_{mqm} + M_{m'q'}}{!_{r} + !_{m} + !_{q}} = 0; \quad (14)$$

where $D_q = D_0 q^2$, $q = q^2$ and $M_{qm} = \frac{p}{2}$, $m_{q;+m}$. In addition, in deriving eq. (14) I have used the Herring consistency equation [10]. In fact Herring's de nition of $!_q$ is one of many possibilities, each leading to a di erent consistency equation. But it can be shown, as previously done in [7], that this does not a ect the exponents (universality).

A detailed solution of equations (13) and (14) in the lim it of small q^0 s (i.e. large scales) in the line of refs. [7]-[8] yields a rich fam ily of solutions that I shall describe im m ediately.

First, there are two kinds of weak-coupling solutions - both with a dynamic exponent z = 2 (they are called weak-coupling because they are exactly the solutions obtained in the case of the FractalE dwards-W ilkinson equation, see [3]). Now, when the spatial correlations of the noise are relevant (i.e., when > 0) I obtain the solution = 2 + 2 -provided that d > 2 + 2. But if the spatial correlations of the noise are not relevant (i.e., when 0) I obtain the simpler solution = 2-provided that d > 2 - 3.

The second type of solutions is strong coupling solutions that obey the well known scaling relation $z = \frac{d+4}{2}$ obtained from eq. (14) (this scaling relation is just the above-mentioned scaling relation + z = 2 that is naturally obeyed by our analysis). The rst strong coupling solution is determined by the combination of the scaling relation and the transcendental equation F (;z) = 0, where F is given by

$$F(;z) = \begin{matrix} Z \\ d^{d}t \frac{t e t}{t^{z} + e t^{z} + 1} \\ h \\ ett + e t t t \end{matrix}$$

$$+ \begin{matrix} Z \\ d^{d}t \frac{t e t^{2}}{t^{z} + e t^{z} + 1} t e t ; (15) \end{matrix}$$

and ê is a unit vector in an arbitrary direction.

This solution is valid as long as the solutions of the last equations satisfy the following condition > max $\frac{d+4+4}{3}$; $\frac{d+4}{3}$; d+2.

It turns out that for d = 1 the equation F (; z ()) = 0 is exactly solvable, and yields = 2 and $z = \frac{3}{2}$ (it can be checked in m ediately by direct substitution). In this case the validity condition reads 2 > m ax $\frac{5+4}{3}$; 1+2or equivalently 0 < 1/2 and < 1/4. By using eq. (11) I translate the results into $= \frac{1}{2}$ and $z = \frac{3}{2}$ that are precisely the exact result presented above.

It should also be mentioned that for d 2 such an exact solution in closed form cannot be found, and one has to solve numerically the equation F (;z()) = 0. For convenience I denote the numerical value of this solution by $_0$ (d). For example, in two dimensions I obtain $_0$ (2) = 2:59.

The second strong coupling solution is obtained by power-counting, and it is relevant when > 0 (i.e. when the spatial correlations of the noise are relevant), d < 2+2 and F (;z()) < 0 (the last condition turns out to be equivalent to the condition $>_0$ (d) because from this value of and on the function F (;z()) is negative). This power counting solution can be written in closed form and is given by $z = \frac{d+4}{3}$ and $= \frac{d+4+4}{3} = z+2$.

The third strong coupling solution (that is in some sense the only "genuine" FKPZ solution, in the sense that it is the only solution that is dram atically in uenced by the fractional Laplacian, and at the same time is not a solution of the Fractal E dwards-W ilkinson equation) is also a obtained by power counting. M one speci cally, it is determined by the combination of the scaling relation $z = \frac{d+4}{2}$ and the extra relation d+ 4 2 z = 2. This solution can be written in closed form as = d+2 and z = 2. It turns out that this phase is relevant when d > m ax f2 3;2 3 + 2 g. In addition, it is needed that F (;z()) is positive. Therefore, this solution is possible only when $<_0$ (d).

The following table sum marizes all the possible phases found in this paper

Z		V alidity
2	2 + 2	> 0 and d > 2 3 + 2
2	2	0 and d > 2 3
2	d+ 2	0 < < 1 and
		d>maxf2 3;2 3+2g
d+ 4 2 3	$\frac{d+4+4}{3}$	> 0, d < 2 + 2 and
		$d + 4 + 4 > 3_{0}$ (d)
$\frac{d+4}{2}$	₀ (d)	$_{0}$ (d) > m ax $\frac{d+4+4}{3}$; $\frac{d+4}{3}$; d+2

TABLE I: A complete description of all the possible phases of the FKPZ problem, for any value of d; and . The rst two columns give the scaling exponents z and for a particular phase, and the third column states each phase's validity condition. Note that $_0$ (d) is the numerical solution of the transcendental equation F (;z) = 0 with the scaling relation $z = \frac{d+4}{2}$ – if such a solution exists.

In order to gain m ore insight on this system , it m ight be interesting to specialize to two extrem e cases: nam ely € 0 and = 0. The rst case = 0 and $\neq 0$ vs. (namely = 0 and \neq 0) corresponds to the local K P Z problem with long-range noise. This problem has been studied in the past using various methods - for example: DRG [9, 32, 33], Mode-Coupling [34] and SCE [8]. All m ethods agree on the basic picture that for a big enough noise exponent () one obtains a power-counting strongcoupling solution, given by $z = \frac{d+4}{3}$ (z is the dynam ic exponent). The controversy between the di erent methods is over the values of the scaling exponents for sm aller values of , and on the critical value $_0$ that separates between the two phases. Not surprisingly, the results given here agree with the previous SCE result presented in ref. [8].

The second case (namely $\neq 0$ and = 0) corresponds to the Fractal KPZ problem with white noise that is the original equation suggested by M ann and W oyczynski [3]. Specializing the general picture of phases presented in Table I above to the case = 0 yields new phase diagram s, namely a separate phase diagram for every dimension d. For example, in two dimensions there are three possible phases: I. The standard KPZ phase given by $_0(2) = 2:59$ that is valid for 0 0:295. II. The weak-coupling phase given by = 2that is valid for any > 0. III. The "third" strong coupling solution given by = d + 2 that is possible for 0 < 1. In the rst phase the dynam ic exponent is z = 1.705, while the other two phases share the same dynam ic exponent




FIG.1: The values of the scaling exponents (a) z and (b) as a function of the fractional parameter , for uncorrelated noise ($\,=\,$ 0) in two-dimensions.

The described two-dimensional (and white-noise) scenario is quite typical and appears in other dimensions as well – namely there are usually three possible phases that are possible for dimensions between them (as in the usual KPZ scenario – the phase transition is controlled by the strength of the dimensionless coupling constant). More precisely, this picture extends up to the upper critical dimension of the original KPZ problem $(d_{L_c}^{K_PZ})$ where the rst phase (phase I in Fig. 1) disappears (see the discussion below).

At this point, the full picture of possible solutions might seem too complex, so I want to suggest the follow ing sim ple interpretation for the behavior of the Fractal KPZ equation. If you remember, the starting point of this discussion was the introduction of the fractional Laplacian into the KPZ equation. This immediately implies that faster relaxations are now possible (faster in the sense that a sm aller dynam ical exponent is expected) when compared with the Edwards-W ilkinson equation. However, it is well known that already the KPZ nonlinearity introduces faster relaxations (at least for dim ensions lower than the upper critical dimension). Therefore, in the FKPZ system, the dynamics is controlled by the fastest "component": if the dynam ical exponent of the classical KPZ system is smaller than 2 then it dom inates, otherwise the new fractionaldynam icalexponent z = 2controls the dynam ics. However, at this point the picture gets a little bit m ore complicated (just like in the classical KPZ case), namely there are several possible phases with this new fractional dynamical exponent z = 2, and the transition between these phases is controlled by the strength of dimensionless coupling constant.

These conclusions have an important implication for the upper critical dimension of the FKPZ model (i.e. the dimension above which the dynamical exponent is the same as that of the linear theory). Namely, it is turns out that the FKPZ equation always has an upper critical dimension that is determined by the relation $_0 d_{uc}^{FKPZ} = d_{uc}^{FKPZ} + 2$ (or alternatively, using the roughness exponent – the upper critical dimension is the dimension where $_{KPZ} d_{uc}^{FKPZ} =)$. Notice, that this result does not dependent on the ongoing debate over the existence of the upper critical dimension for the

KPZ system (see [11]-[20]). A ctually, it merely requires that the roughness exponent of the classical KPZ system becomes arbitrarily small in higher dimensions - an assumption that is generally accepted (only Tu [21] had a dimension conjecture).

As one can see, the results obtained using the Self-Consistent Expansion (SCE) are quite general, and cover all possible values of the relevant parameters (and) as well as dimensions. It is also easily veried that these results recover the exact result obtained at the beginning of this paper for the case = 2. This situation, suggest that the SCE method is generally appropriate when dealing with non-linear continuum equations.

A cknow ledgement: I would like to thank M oshe Schwartz for useful discussions.

- [1] M. Kardar, G. Parisi and Y.-C. Zhang, Phys. Rev. Lett. 56, 889 (1986).
- [2] A.-L. Barabasi and H.E. Stanley, Fractal Concepts in Surface Growth (Cambridge Univ. Press, Cambridge, 1995).
- [3] JA. Mann Jr. and W A. W oyczynski, Physica A 291, 159 (2001).
- [4] G L.Kellog, Phys. Rev. Lett. 72, 1662 (1994).
- [5] F.Fam ily and T.Vicsek, J.Phys. A 18, L75 (1985).
- [6] M. Schwartz and S.F. Edwards, Europhys. Lett. 20, 301 (1992).
- [7] M. Schwartz and S.F. Edwards, Phys. Rev. E 57, 5730 (1998).
- [8] E. Katzav and M. Schwartz, Phys. Rev. E 60, 5677 (1999).
- [9] E. Medina, T. Hwa, M. Kardar and Y. C. Zhang, Phys. Rev. A 39, 3053 (1989).
- [10] J.R. Henring, Phys. Fluids 8, 2219 (1965); 9, 2106 (1966).
- [11] T. A la-N issila, Phys. Rev. Lett. 80, 887 (1998).
 J.M. K in , ibid. 80, 888 (1998).
 M. Lassig and H. K inzelbach, ibid. 80, 889 (1998).
 [21] E. Darlen en and M. Schwartz, Dhuging B. 224, 5
- [12] E. Perlsm an and M. Schwartz, Physica A 234, 523 (1996).
- [13] C.Castellano, M.M arsili and L.Pietronero, Phys. Rev. Lett. 80, 3527 (1998).
 - C.Castellano, A.Gabrielli, M.Marsili, M.A.Munoz and L.Pietronero, Phys. Rev. E 58, 5209 (1998).
- [14] T.Halpin-Healy, Phys. Rev. A 42, 711 (1990).
- [15] T. Blum and A. J. McKane, Phys. Rev. E 52, 4741 (1995).
- [16] J-P.Bouchaud and M.E.Cates, Phys. Rev. E 47, 1455 (1993).

erratum), Phys. Rev. E 48, 653 (1993).

- [17] E.Katzav and M.Schwartz, Physica A 309, 69 (2002).
- [18] J.Cook and B.Derrida, J.Phys. A 23, 1523 (1990).
- [19] F. Colaiori and M. A. Moore, Phys. Rev. Lett. 86, 3946 (2001).
- [20] E. Marinari, A. Pagnani, G. Parisi and Z. Racz, Phys. Rev. E 65, 26136 (2002).
- [21] Y.Tu, Phys. Rev. Lett. 73, 3109 (1994).
- [22] R.M etzler, E Barkai and J.K lafter, Phys. Rev. Lett. 82, 3563 (1999).
- [23] R. M etzler and J. K lafter, Chem. Phys. Lett. 321, 238 (2000).
- [24] R.Granek and J.K lafter, Europhys. Lett. 56, 15 (2001).
- [25] M. Sokolov, J.K lafter and A.B lum en, Phys. Rev. E 64, 21107 (2001).
- [26] H. Scher and E. Montroll, Phys. Rev. B 12, 2455 (1975).
- [27] Q.Gu, E.A.Schi, S.G rebner and R.Schwartz, Phys. Rev. Lett. 76, 3196 (1996).
- [28] A.Klemm, H.-P.Muller and R.Kimmich, Phys. Rev. E 55, 4413 (1997).
- [29] F.Amblard, A.C.Maggs, B.Yurke, A.N.Pargellis and S.Leibler, Phys. Rev. Lett. 77, 4470 (1996).
- [30] M. Lassig, Nucl. Phys. B 448, 559 (1995).
- [31] K J.W iese, J. Stat. Phys. 93, 143 (1998).
- [32] E. Frey, U.C. Tauber and H.K. Jansen, Europhys. Lett. 47, 14 (1999).
- [33] H K. Janssen, U C. Tauber and E. Frey, Europhys. J. B 9, 491 (1999).
- [34] A.Kr.Chattapohadhyay and J.K.Bhattacharjee, Europhys. Lett. 42, 119 (1998).
- [35] E.Frey and U.C. Tauber, Phys. Rev. E 50, 1024 (1994).