Dipole relaxation losses in DNA

M.Briman, N.P.Armitage, E.Helgren, and G.Gruner

Department of Physics and Astronomy, University of California, Los Angeles, CA 90095

(April 14, 2024)

The electrodynamic response of DNA in the millimeter wave range is investigated. By performing measurements under a wide range of humidity conditions and comparing the response of single strand DNA and double strand DNA, we show that the appreciable AC conductivity of DNA is not due to photon activated hopping between localized states, but instead due to dissipation from dipole motion in the surrounding water helix. Such a result, where the conductivity of DNA with the constrained motion of overdam ped dipoles, reconciles the vanishing DC conductivity of DNA with the considerable AC response.

PACS numbers: 87.14 Gg, 72.80 Le

The electrical conductivity of DNA has been a topic of much recent interest and controversy [1]. Measurements from dierent groups have reached a variety of conclusions about the nature of charge transport along the double helix. DNA has been reported to be m etallic [2], sem iconducting [3], insulating [4,5], and even a proxim ity e ect induced superconductor [6]. However, questions have been raised with regards to the role played by electrical contacts, length e ects, and the manner in which electrostatic dam age, residual salt concentrations, and other contam inations m ay have a ected these results [1]. M ore recent m easurem ents, where care was taken to both establish a direct chem ical bond between -DNA and Au electrodes and also control the excess ion concentration, have given com pelling evidence that the DC resistivity of the DNA double helix over long length scales (< 10 m) is very high indeed (> 10^{6} am) [7]. Such DC measurements contrast with recent contactless AC m easurem ents that have shown that there is appreciable conductivity at m icrowave and far-infrared frequencies [8,9] the magnitude of which approaches that of a welldoped sem iconductor [10].

P reviously, the AC conductivity in DNA was found to be well parameterized as a power-law in ! [8,9]. Such a dependence can be a general hallm ark of AC conductivity in disordered systems with photon assisted hopping between random localized states [11] and led to the reasonable interpretation that intrinsic disorder, counterion uctuations, and possibly other sources created a small number of electronic states on the base pair sequences in which charge conduction could occur. However, such a scenario would lead to thermally activated hopping conduction between localized states and is thus inconsistent with the very low DC conductivity [7]. A number of outstanding issues arise: A re there localized regions along the helix where a continuous conducting path is not present, but stillAC hopping between localized states over distances of a few base pairs can occur? A re there sensitive length dependencies in the DNA strands? Is

there a di erence between between the sam ples of various groups? Perhaps di erent charge conduction mechanism s play a role at nite frequency.

To the end of resolving som e of these m atters, we have perform ed AC conductivity experiments in the millimeter wave range under a wide range of hum idity conditions. We show that the appreciable AC conductivity of DNA in the microwave and far infrared regime should not be viewed as some sort of hopping between localized states and is instead likely due to dissipation in the dipole response of the water molecules in the surrounding hydration layer. It can be well described by a Debye-like relaxation of water molecules in the surrounding water helix. At low hum idities the response is well modelled by considering the rotation of single water molecules in the structuralwater layer. As the number of waterm olecules per base pair increases, dissipation due to the collective motion of water dipoles increases, until eventually the conductivity resembles that of bulk water. By measuring both single strand (ssDNA) and double strand DNA (dsDNA) over a wide range of hum idities we are able to show that, at least in principle, all the AC conductivity of DNA can be assigned to relaxation losses of water dipoles. This result reconciles the apparent com plete lack of DC conductivity with the appreciable AC response.

Double stranded DNA In swere obtained by vacuum drying of 7m M PBS solution containing 20 mg/ml sodium salt DNA extracted from calf thym us and salm on testes (Sigm a D 1501 and D 1626). The results were found to be independent of the use of calf or salm on DNA. Our choice for these concentrations deserves further explanation. It is wellknown that at a given tem perature double helical conformation of DNA can exist in solution only with a certain concentration of positive ions. Excess salt cannot be rem oved by vacuum drying, so large am ounts of residual salt in In s could introduce signi cant errors in conductivity, due to both the ionic conduction of the salt itself and its additional hydration during hum idity changes. M elting tem perature calculations [12,13] for

long native pieces of DNA with C-G content around 40% show that 2-10 m M concentration of sodium cations is enough to stabilize the double helix at room tem perature. Films were prepared with di ering salt amounts and it was found that as long as the excess salt m ass fraction is kept between 2-5% the nalresults were not signi cantly a ected. In order to improve the DNA/salt mass ratio we used a high concentration of DNA, but 20 mg/ml appears to be the lim it. Higher concentrations makes it di cult for DNA bers to dissolve and the solution becomes too viscous, which prevents producing the at uniform Im swhich are of param ount im portance for the quasi-optical resonant technique. Single stranded DNA In swere prepared from the same original solution as the double stranded ones, with prelim inary heating up to 95 C for 30 m inutes and fast cooling down to 4 C . In both dsDNA and ssDNA cases the conform ational state was checked by uorescent m icroscope m easurements. The dry Im swere 20 to 30 m icrons thick and were made on top of 1mm thick sapphire windows. Immediately after solution deposition onto the sapphire substrates the air inside the viscous solution was expelled by vacuum centrifuging at 500g, otherwise the evaporation process causes the form ation of air bubbles that destroy the Im uniform ity.

The AC conductivity was measured in the millimeter spectral range. Backward wave oscillators (BW 0) in a quasi-optical setup (100 G hz - 1 T H z) were em ployed as coherent sources in a transmission con guration. This di cult to access frequency range is particularly relevant as it corresponds to the approxim at eexpected time frame for relaxation processes in room tem perature liquids (1-10 ps). Importantly, it is also below the energy range where one expects to have appreciable structural excitations. The technique and analysis are well established [14]. We utilize the fact that for plane waves incident norm ally on a slab of material, transmission resonances occur when the slab is an integer number of half wavelengths. Thus, using a 1 mm sapphire disc as a substrate, resonances occurred approxim ately every 50 G H z. Having analyzed the transmission through the sapphire alone prior to m ounting the sam ple, the optical properties of the substrate were well characterized. Thus using a two-layer transmission model, each resonance can be analyzed to extract the optical properties of the DNA In , allowing for a 1.5 cm^{-1} resolution of the spectra.

Sam ples were measured at room tem perature at several xed hum idity levels which were maintained by putting them in a herm etically sealed environment with a saturated salt solution [15]. The change in thickness and mass of the DNA Im s at di erent hum idities were tracked by separate measurements within a controlled environment for each sample in a glove box. The total number of water molecules per nucleotide A can be correlated to the relative hum idity x (x=0-1) through the so-called B ranauer-Emmett-Teller (BET) equation [16]

$$A = \frac{BCx}{(1 x)(1 x + Cx)};$$
 (1)

The constant B is the maximum number of water m olecules in the rst layer sites. A coording to the statistical form ulation of the BET equation by Hill [17], m obile water molecules within the double helix can be characterized as 2 types. The rst are ones within the initial hydration layer, which are directly attached to DNA and have a characteristic binding energy 1. W ater m olecules of the second and all other layers can be approxim ated as having a binding energy $\ _{\rm L}$. To a good approximation this $_{\rm L}$ can be taken to be that of bulk water. These param eters enter into the BET equation through the expression for C which equals $D e^{(\frac{1}{kT})}$ where D is related to the partition function of water. A lso we should note that there is, in actuality, a structural 0-th layer of water molecules, containing 2.5-3 water molecules per nucleotide that cannot be rem oved from the helix under typical conditions [18].




FIG.1. Absorbtion of water molecules per nucleotide as a function of hum idity. The data represented by the open circles is taken from Falk et. al.

That it is reasonable that the mobile water layers of DNA can be modelled by distinguishing 2 di erent sets of water param eters was rst established by Falk et al.'s [15] use of the BET equation to describe the hydration of sodium and lithium DNA salts from calf thym us and salm on testes. They found good agreem ent between experimental data and theory with constants B = 2.2 and C = 20. We perform ed a sim ilar hydration study of our dsDNA and ssDNA lm s; as shown in Fig. 1 the hydration of our lm s are perfectly consistent with Falk's result. Note that there is no appreciable di erence in the hydration between dsDNA and ssDNA.

In Fig. 2 we present data for the extracted $_1$ (!) of

both dsDNA and ssDNA thin lms. One can see that in both cases, the conductivity is an increasing function of frequency. Since the conductivity is also an increasing function of hum idity, one may wish to try to separate the relative contributions of charge motion along the DNA backbone from that of the surrounding water molecules.




FIG.2. Frequency dependence of the conductivity of calf thym usDNA at di erent relative hum idity levels. (a) Double stranded DNA (b) Single stranded DNA (c) A com parison of conductivity between single and double stranded DNA.

First, one can consider that there should be two main e ects of hydration in our dsDNA lms. There is the hydration itself, where water molecules are added in layers to the double helix; this is well described by BET equation [16]. Additionally, the conform ational state of dsDNA also changes as a function of hum idity. For exam – ple, sodium salt calf thym usDNA is in a B-like disordered form at hum idities from 0-40%, above which it transfers to the A form, and nally to a well ordered B-form at hum idities higher than 80% [19,20]. Additional water molecules certainly contribute to the increase in conductivity, but at high hum idities there is the possibility that som e of the conduction might be due to an increase in electron transfer along the dsDNA helix in the ordered B form . However since such an e ect would be much reduced in disordered and denaturalized ssDNA lm s and since Fig. 2 shows that to within the experimental uncertainty the conductivity of dsDNA and ssDNA in the millimeter wave range is identical, it is most natural to suggest that water is the major contribution to the AC conductivity. From this comparison of dsDNA and ss-DNA, we nd no evidence for charge conduction along the DNA backbone.

In Fig. 3 we plot the the conductivity 1 of the DNA In s normalized by the expected volume fraction of waterm olecules including both the hydration layers plus the structural water. A lthough this normalization reduces the spread in the thin Im conductivity at the lowest frequencies it does not reduce it to zero, showing that if the largest contribution to the conductivity comes from water, the character of its contribution changes as a function of hum idity.

The complex dielectric constant of bulk water has been shown to be well described by a biexponential D ebye relaxation model [21{23], where the rst relaxation process [21], characterized by a time scale $_{\rm D}$ = 8:5 ps, corresponds to the collective motion of tetrahedral water clusters, and the second from faster single molecular rotations [24] with a time scale $_{\rm F}$ = 170 fs. For bulk water, the contribution of each relaxation process is determined by the static dielectric constant $_{\rm S}$ (T) = 87:91e $^{0:0046T\ f^{\rm C}}$, $_{\rm L}$ = 52, and the dielectric constant at high frequencies $_{\rm L}$ = 3:3.

$$b(w) = \frac{1}{1 + i!} + \frac{s}{1 + i!} + \frac{1}{1 + i!} + \frac{1}{1 + i!}$$
(2)

W hen applying Eq. 2 to the dipole relaxation losses of DNA, one expects that the relative contributions of the two frequency dependent terms will change as increasing hum idity increases the average e ective coordinate number. For instance, at 0% hum idity it is reasonable to assume that the rst term which is due to the collective motion of water clusters, cannot play a role as the structural water is not tetrahedrally coordinated. For high hydration levels, where multiple water layers exist around the dipole helix, the relaxation losses of the water layer may approach those of bulk water. We can com pare the above equation using the independently known values [21] for $_{\rm D}$, $_{\rm S}$, $_{\rm F}$ and $_{\rm 1}$ to the experimental data norm alized to the expected volum e fraction of the water from the independently determ ined water uptake curves shown in Fig. 1. In Fig. 3, along with the experimental data at two representative hum idity levels, two theoreticalcurves for 0% and 100% hum idity are plotted. W ith the only two assumptions being that at 0% hum idity, the sole relaxational losses com e from singly coordinated water molecules in the structural water layer and that it is only at higher hum idity levels where the collective losses can gradually play a greater role, the theoretical curves provide a very good t to the data over alm ost all of the m easured frequency range. At low hum idity the data is well m atched by the theory incorporating only single m olecule rotations. At high hum idity, the data begins to approach the behavior of 'free' water. For these two lim its the theoretical curves have no free param eters.




FIG.3. Conductivity of dsDNA and ssDNA lms normalized by the volume fraction of all water molecules (structural plus hydration layer). For clarity, only 0% and 84% hum idities are shown. The solid line represents the conductivity of pure water as modelled by the biexponentialD ebye model using the parameters of R onne et al. The dashed line shows just the contribution from single water molecule relaxation.

The only appreciable discrepancy between theory and experiment is the high frequency data at low hum idity, where the biexponentialD ebyem odelunderestim ates the conductivity. This may be due to a number of reasons. At very low relative hum idities it is possible for the ionic phosphate groups on the DNA backbone to form stable dihydrates which may give their own contribution to relaxation losses through their additional degree of freedom [15]. A hematively, it may also be that at higher frequencies for low hydration sam ples, the weak restoring force from charge-dipole interaction in the structural water layer begins to become more signi cant and our biexponentialD ebye m odel becomes less applicable.

In conclusion, we have found that the considerable AC conductivity of DNA can be largely ascribed to relaxational bases of the surrounding water dipoles. The conductivity of ssD NA and dsD NA was found to be identical to within the experim ental error, indicating that there is essentially no charge conduction along the DNA backbone itself. The conclusion that the observed conductivity derives from the water layer is supported by the fact that, over m uch of the range, it can be well described by a biexponentialD ebye m odel, where the only free param – eter is the relative contributions of single water m olecule and tetrahedral water cluster relaxation m odes. G enerally speaking, because m any large biom olecules have surrounding water layers, a result such as ours shows that one must be aware of the possibility of such relaxation losses when investigating the electrodynam ic response of such system s.

W e would like to thank K.G reskoviak for help with sample preparation. The research at UCLA was supported by the National Science Foundation grant DMR-0077251.

- [1] C.Dekker and M.Ratner, Physics W orld 14, 29 (2001).
- [2] H. W. Fink and C. Schonenberger, Nature 398, 407 (1999).
- [3] D. Porath et al, Nature 403, 635 (2000).
- [4] E.Braun et al, Nature 391, 775 (1998).
- [5] P.J. de Pablo et al, Phys. Rev. Lett. 85, 4992 (2000).
- [6] A.Y.Kasum ov et al, Science 291, 280 (2001).
- [7] Zhang et al. Phys. Rev. Lett. 89, 198102 (2002).
- [B] P. Tran, B. A lavi and G. G runer, Phys. Rev. Lett. 85, 1564 (2000).
- [9] E.Helgren et al. cond-m at/0111299.
- [10] E. Helgren, N P. A mm itage, and G. G runer, Phys. Rev. Lett. 89, 246601 (2002).
- [11] A.L.E fros and B.I.Shklovskii, J.Physics C 8, L49 (1975).
- [12] E.Breslauer et al, Proc.Nat.Acad.Sci.83, 3746 (1986).
- [13] N. Sugim oto et al, Nucl. A cids Res. 24, 4501 (1996).
- [14] A. Schwartz et al., Rev. Sci. Instrum . 66, 2943 (1995).
- [15] M. Falk, K. Hartman, R. Lord, J. Am. Chem. Soc. 84, 3843 (1962).
- [16] S.Braunauer, P.Emmett, E.Teller, J.Am. Chem. Soc. 80, 309 (1938).
- [17] T.L.Hill, J.Chem. Phys. 14, 263 (1946).
- [18] N. Tao and S. Lindsay, Biopolym ers 28, 1019 (1989).
- [19] V.Y.Maleev et al., Bio zika 38, 768, (1993).
- [20] S.M. Lindsay et al. Biopolymers, 27, 1015, (1988).
- [21] C.Ronne et al., J.Chem. Phys. 107, 5319 (1997)
- [22] J.K indt, C.Schm uttenm aer, J.Phys.Chem .100, 10373 (1996).
- [23] J. Barthel, R. Buchner, Pure and Appl. Chem. 63, 1473 (1991).
- [24] N.Agmon, J.Phys.Chem. 100, 1072 (1996).