

Duality in process of noncommutative deformation and topological nature of integral Cherepanov-Rice

Trinh V.K

Department of physics of polymers and crystals, MGU

E-mail : khoa@polly.phys.msu.ru

Summary: In this papers it is showed, that for the noncommutative deformation simultaneously exist also loading deformation H , and unloading deformation H^v . The real deformation is a combination of these types deformations. The criterion of destruction J reflects topological character of environment, i.e. defines properties of symmetry of environment at destruction. It is possible to tell, that during destruction the energy is released not continuously, and discretely. This situation is reflected through topological number Q or number of unloading, connected to him.

1. As is known, after removal of loading an elastic body always comes back in an initial status. The given definition of elasticity is a little simplified. If the pressure(voltage) exceeds a limit of elasticity, the dependence between loading and deformation ceases to be one-digit and depends on the order of the application of loading [1]. For simplicity, it is supposed, that the equation of a curve is

$$\sigma = \Phi(e)$$

fair at any program of the application of load, when a stress monotonously grows. Nevertheless for real materials the business is more complicated. Let we have finished loading up to a point $A = (\sigma', e')$ of areas of plasticity. After that is made unloading. Then the stress σ decreases up to zero. In this process it is reflected not only plastic behaviour of material, but also elastic. More careful experiments have shown, that the law of unloading is not described precisely by direct line. Replacing it by its closest straight line, we find, that its inclination not corresponds(meets)

in accuracy to the initial module of elasticity. At polymeric materials, and also at composite materials, for example, fibreglasses, the law of unloading differs from the law Hooke very essentially. Apparently, at the application of load of material in it the defects are formed. Naturally there is a question: Whether exists duality of statuses of loading and unloading? And in general, whether exists duality during plastic deformations? Let's consider this problem in given clause.

2. As is known, under action of loading material passes in plastic status. All data of experiments show, that in this status a the stress and the deformation is strongly in fluctuation. More that, diagram stress - deformation $\sigma = f(e)$ is similar with the diagram pressure - volume $P = f(V)$ in a liquid [2]. On the other hand , the local phase transition in the end of a crack is observed [3-5]. Thus, status, close to destruction it is possible to consider(count) as a status of transition or critical, in which spontaneously arises a structure [6]. Density of distribution of probability of transition to plasticity is the solution of the equation Fokker - Panck. However, using potential character of process of deformation, instead of the equation Fokker - Planck we can use the equation Schrodinger. It means, that it is possible to consider(count) process of plastic deformation by process with supersymmetry. Here tensor of deformation is considered in the role of parameter of order. In work [1] A.A. Iliushin has presented on discussion the concept of a trajectory of deformation in space E_5 . Actually, the space E_5 is a layer in stratified space of deformation [6]. It means, that a field of deformation we can consider(count) as a gauge field, in which generalized trajectory of deformations Iliushin is the section of a stratification of deformation. At identification of the generalized trajectory of deformation with knots we have shown topological character of process of deformation connected with invariant Witten. Thus, density of distributions of probability of transition to plasticity is received[6] :

$$Z = \int \exp[-kS_{cs}(e)]\mathcal{D}e,$$

Here k — it is factor elasticity, S_{cs} — action Chern - Simons

$$S_{cs} = \int \Gamma_{cs},$$

where Γ_{cs} - the form Chern - Simons for variety of deformation.

3. The spontaneous infringement of symmetry has allowed us to receive laminar structure in process of thermomechanical deformation [6]. Here from this mechanism we can describe two processes of loading and the unloadings with their topological character, which will be considered by dual processes. Processes of deformation in the end of a line dislocation or the cracks are those examples of unloading, which we would like to consider. These processes of clearing of energy of the unloadings were used in model of seismology. Nevertheless it is necessary to notice, that the nature of process of plastic deformation - it is noncommutative nature. Thus, the process of deformation is given $P = \{E_{ij}(x, t)\}$. It is noncommutative space of deformation. The deformation wave is the noncommutative wave. Besides it is possible to consider a scalar field describing actions of defect on deformation [7]. For the complete description of the distributions of a deformation wave it is necessary to us to use the space being tensor product $C(R_4) \otimes M_n$, and $C(R_4)$ - algebra of smooth functions determined on usual space - time, and M_n - algebra $n \times n$ of matrixes. Now our task is a construction of noncommutative bundles of deformations above $C(R_4) \otimes M_n$. This model of deformation was constructed in [6]. So, we admit(allow) connection as $\omega = A_\mu dx^\mu + (B_k - iE_k \otimes 1)\theta^k$, where A_μ is function in V , receiving value in $M_n(C) \times M_r(C)$, $\mu \in \{0, 1.., s\}$, B_k - function above V , having the value in $M_n(C) \times M_r(C)$. Thus, action will be expressed in the form:

$$S = - \int_{R^{s+1}} \frac{1}{4n} Tr(F_{\mu\nu} F^{\mu\nu}) + \frac{1}{2n} Tr[(\nabla_\lambda \phi_k)(\nabla^\lambda \phi^k)] \\ + \frac{1}{4n} \Sigma Tr[[\phi_k, \phi_l] - \mathbf{m} \Sigma C_{klm} \phi_m]^2$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$, $\phi_k = \mathbf{m} B_k$, $\nabla_\lambda \phi_k = \partial_\lambda \phi_k + [A_\lambda, \phi_k]$

” Magnetic monopole ”, arising during evolution gauge fields, actually is the deformation of the unloading. Really, from action S we shall receive the equation of movement

$$\delta S = 0$$

This is nonlinear equation, his(its) solution with final energy is important for research of the topological character of system. It is wonderful, that, not solving this equation, it is possible to receive some information on its properties. For this purpose from Lagrangian it is simply necessary to find expression for retentive energy of system, which for static decisions, with the account that $A_0^a = 0$, looks like

$$W = - \int_{R^s} d^s x \frac{1}{4n} Tr(F_{\mu\nu} F^{\mu\nu}) + \frac{1}{2n} Tr[(\nabla_\lambda \phi_k)(\nabla^\lambda \phi^k)] \\ + \frac{1}{4n} \Sigma Tr[[\phi_k, \phi_l] - \mathbf{m} \Sigma C_{klm} \phi_m]^2$$

This integral looks like integral Cherepanov - Rice without kinetic energy. Energy achieves minimum and addresses in zero at

$$A_0(x) = 0, \quad \phi_i(x) = \frac{1}{2} \mathbf{m} E_i, \quad \nabla_\lambda \phi^i = 0$$

Last equation turns to the equation $\partial_i \phi^a = 0$. However, in this case there are two gauge orbit. Let Ω_0 - vacuum appropriate to an orbit ($A_\mu = 0$, $\phi_k = 0$), and Ω_1 - vacuum appropriate to orbit ($A_\mu = 0$, $\phi_k = i \mathbf{m} E_k$). Thus, this condition is carried out for each orbit. In a case, when it is required for finiteness of energy $W < \infty$, then there is a following condition: because of that the condition of finiteness W

consists in enough fast aspiration of fields to some configuration with $W = 0$ on spatial infinity, it is visible, that the condition for ϕ at $r \equiv |x| \rightarrow \infty$ will look like $r^{3/2} \nabla_i \phi \rightarrow 0$, $\phi \rightarrow F$. For getting character of the field A we shall present ∇_i in spherical coordinates $\{r, \theta, \phi\}$, θ - the component is determined by expression $(\nabla \phi^a)_\theta = \frac{1}{r} \frac{\partial \phi^a}{\partial \theta} + g C^{abc} A_\theta^b \phi^c$. For performance of the condition given above, in the field A_θ^b to be satisfied condition $A_\theta^b = \frac{1}{r} A(\mathbf{n})$, where $A(\mathbf{n})$ depends only on a spatial corner: $\sum n_i^2 = 1$. Other components have similar character. Thus, there are available boundary conditions in the infinity

$$A_\mu^a = \frac{1}{r} A(\mathbf{n}), \quad \phi^a = \frac{1}{2} \mathbf{m} E_a, \quad r^{3/2} \nabla_i \phi = 0$$

So, shall receive topological index, which was similar expression in commutative case

$$Q = \int d^3x K_0$$

and topological current looks like

$$K_\mu = \frac{1}{8\pi} \text{Tr}(\partial^\mu \hat{\phi}^a \partial^\nu \hat{\phi}^b) \partial^\sigma \hat{\phi}^c$$

Where $\hat{\phi}_a = \phi^a / |\phi|$, $|\phi| = [\text{Tr}(\phi\phi)]^{1/2}$. The charge of unloading looks like $m = \int \frac{K_0}{g} d^3x = Q/g$. In a case $n, r = 1$ we shall receive usual monopole t Hoof- Polyakov.

4. Let deformation environment has symmetry G . For simplicity let's assume, that G - it is semisimple group. In process of deformations the spontaneous infringement of symmetry $G \rightarrow H \subset G$ will appear. Our purpose - to investigate structure of a component $A(\theta)$. In commutative case in [8] approved, that a component $A(\theta)$ of monopole is full determined through the Higgs field ϕ . In ours noncommutative case on the basis of the equation of movement we also receive the same statement. From the requirement of finiteness of energy in a stationary

status or from integral such as Cherepanov - Rice we receive topological invariant. Shall copy this result in the equivalent form, which does not diminish a generality of the problem:

$$\exp(4\pi i \epsilon A(\theta)) = 1$$

where ϵ - a constant of connection of defect with environment. Let $\mathcal{L}(H)$ - Lie algebra of group H , and $T_1, T_2 \dots T_r$ its generators. Thus, under the theory of Lie group and Lie algebra [10] it is possible to find gauge maps $S \in H$, so that

$$\epsilon A(\theta) = S \sum_{i=1}^r \beta_i T_i S^{-1}$$

The system $\beta_1, \beta_2 \dots \beta_r$ - is named as weight of statuses of unloading. Using this concept, it is possible to copy condition of "quantization" as

$$\exp(4\pi i \sum_{i=1}^r \beta_i T_i) = 1$$

If H is group for loading, it is necessary to search for group H^v for unloading. Thus, our purpose is to deform system of weight of group H^v , satisfying to the condition above mentioned. For this purpose we use method in [9]. Let for group H there is a system of roots $\alpha \in \Phi(H)$. Then system of roots ω of universal enveloping group \tilde{H} will look like

$$\Lambda(\tilde{H}) = \{\omega ; 2\omega \cdot \frac{\alpha}{\alpha^2} \in Z ; \alpha \in \Phi(H)\}$$

and $\Phi(H)$ - system of roots of group H . First stage given in [9] is to deform system β so that the condition was satisfied :

$$\widetilde{\exp}(4\pi i \beta \cdot T) \in Z(\tilde{H})$$

where the centre - $Z(\tilde{H})$. Here there is a system of dual roots $\Phi^v(H) = \{\alpha^v = N^{-1} \frac{\alpha}{\alpha^2} ; \alpha \in \Phi(H)\}$. For the elementary case, when H - simple group, N will be number as $N^2 = (\sum \frac{1}{\alpha^2}) / \sum \alpha^2 =$. System of weight

$\Lambda(\tilde{H}^v)$ has the form $\Lambda(\tilde{H}^v) = \{\omega ; 2\omega N\alpha \in Z ; \alpha \in \Phi(H)\}$. The greatest system of weight β , satisfying condition $2\alpha.\beta \in Z$, $\alpha \in \Phi(H)$, is written as $\beta = N\omega$, $\omega \in \Lambda(\tilde{H}^v)$. The following step is to look for system carrying out a condition $\exp(4\pi i \sum \beta_i T_i) = 1$ or condition $2\beta.\omega \in Z$, $\omega \in \Lambda(H)$. The general form of system β , carrying out the described above condition, looks like $N^{-1}\beta \in \Lambda(H^v)$. System β determines all characters of group H^v

5. So, process of destruction of environment the following. A crack (defect) is given in environment with group of symmetry G . Under action of loading system of environments with a crack passes in a status of destruction (status of phase transition). Thus there is a spontaneous infringement symmetry $G \rightarrow H$. More precisely, it is the process $G \rightarrow H \otimes H^v$. It means, that simultaneously exist also loading H , and unloading H^v . The real deformation is a combination of these types deformations. The presence of topological objects creates a difference between loading and unloading. From the requirement of finiteness of the energy of deformation and under influence of spontaneous infringement of symmetry for stationary status we shall receive, that integral of energy is invariant. Simultaneously the expression under integral is potential of deformation. Thus, this integral looks like integral Cherepanov - Rice J without kinetic parts of energy. In the classical mechanics of destruction integral Cherepanov - Rice is considered as criterion of destruction. Actually, on our model, the criterion of destruction J reflects topological character of environment, i.e. defines properties of symmetry of environment at destruction. It is possible to tell, that during destruction the energy is released not continuously, and discretely. This situation is reflected through topological number Q or number of unloading, connected to him

- [1] A.A. Iiushin, Mechanics of continuous environments, MGU 1994; V.D. Klusnikov, Physical and mathematical bases of durability and Plasticity, MGY 1994; L.I. Sedov, Mechanics of continuous environments, M. Science, 1973;
- [2] M.Ausloos, Solid Stat. Comm., 1986, vol.59, No.6, p.401-404.
- [3] V.G. Gargin, Superfirm materials, 1982, No.2, c.17-20;
- [4] V.A. Pesin, N.N. Tkachenko, L.I. Feldchuk 1979, JFCh. Volume 53, No.2, p.2794.
- [5] Li-Shing Li, Pabst R.J., Material Sci., 1980, vol.15, No 10, p.2861
- [6] Trinh V.K., Report of RAN, 2000., volume 372, No 4, c.473; The report of RAN, 2001., volume 378, No 3, c.343; analysis and its The application, Thesis of the reports of VZMS-2000 conference; mathematical modeling in natural and humanitarian sciences, Voronezh, January 20-27, 2000.; cond-mat/9907290; the bulletin MGU: physics and astrophysics, 2002, no. 1, p.49.
- [7] H.Kleinert, Phys. Lett., Vol 89A, No 6;
- [8] Radgiaraman P., Soliton and instanton in quantum The theories of a field, M. The world: 1985;
- [9] Godard P., Nuyts J., Olive D., Nucear Physics, B125 (1977) 1-28.
- [10] Serr Z.P., Lie Group and Lie algebra, M.: Mir, 1969;