O rbital entanglem ent and violation of B ell inequalities in m esoscopic conductors

P. Sam uelsson, E.V. Sukhorukov and M. Buttiker

Departem ent de Physique Theorique, Universite de Geneve, CH-1211 Geneve 4, Switzerland.

We propose a spin-independent scheme to generate and detect two-particle entanglement in a mesoscopic normal-superconductor system. A superconductor, weakly coupled to the normal conductor, generates an orbitally entangled state by injecting pairs of electrons into di erent leads of the normal conductor. The entanglement is detected via violation of a Bell inequality, form ulated in terms of zero-frequency current cross-correlators. It is shown that the Bell inequality can be violated for arbitrary strong dephasing in the normal conductor.

PACS num bers: 74.50.+ r, 73.50.T d, 73.23 Ad

Entanglem ent is one of the most intriguing features predicted by quantum theory [1]. It leads to correlation between distant particles, which can not be described by any local, realistic theory [2]. This nonlocal property of entanglem ent has been demonstrated convincingly in optics [3], where entangled pairs of photons have been studied over several decades. A part from the fundam ental aspects, there is a grow ing interest in using the properties of entangled particles for applied purposes, such as quantum cryptography [4] and quantum computation [5].

Recently, much interest has been shown for entanglement of electrons in solid state devices. A controlled generation and manipulation of electronic entanglement is of importance for a large scale implementation of quantum information and computation schemes. Electrons are however, in contrast to photons, massive and electrically charged particles, which raises new fundamental questions and new experimental challenges. Existing suggestions are based on creating [6, 7], manipulating and detecting [8, 9, 10] spin-entangled pairs of electrons. This requires experimental control of individual spins via spin

Iters or locally directed m agnetic elds on a m esoscopic scale. Here we propose a spin-independent scheme for creating and detecting orbital entanglement in a mesoscopic norm al-superconductor system.

We show that a superconductor, weakly coupled to a norm al conductor (see Fig. 1), creates an orbitally entangled state by emitting a coherent superposition of pairs of electrons into di erent leads of the norm al conductor. The zero-frequency correlation between currents owing into di erent norm al reservoirs is shown to be equivalent to a pair coincidence measurement: only correlations between the electrons from the same entangled pair contribute. As a consequence, a standard Bell Inequality (BI) can be directly form ulated in terms of the zero-frequency current correlators. We nd that a violation of the BI, dem onstrating the entanglem ent of the pair state, can be obtained for arbitrary dephasing in the norm al conductor.

We rst consider a simplied version of the system (see Fig. 1), a more detailed discussion is given below.

A single [11] superconductor (S) is weakly coupled to a norm alconductor, a ballistic two-dimensional electron gas, via two tunnel barriers 1 and 2 with transparency

1. The norm al conductor consists of four arm s, 1A;1B;2A and 2B, with equal lengths L. The arm s 1A and 2A (1B and 2B) are crossed in a controllable beam splitter A (B), parameterized via the angle $_{A}$ ($_{B}$), and then connected to norm al reservoirs + A and A (+ B and B). The beam splitters [12] are assumed to support




FIG.1: The system : A single superconductor (S) is connected to four norm alarm s via two tunnel barriers 1 and 2 (thick black lines). The arm s are joined pairwise in beam splitters A and B and end in norm al reservoirs + and .

only one propagating m ode. The states j + j; i and j ; i for electrons going out into the norm al reservoirs and the states jl; i and j2; i of the electrons em itted from superconductor are related via a scattering m atrix:

$$j$$
; i \cos \sin j^2 ; i ; = A;B:(1)
j; i \sin \cos j^1 ; i ; = A;B:(1)

The angles $_{\rm A}$ and $_{\rm B}$ can be tuned between 0 and =2 by tuning the beam splitter from \open", when the electrons are transmitted through from 1(2) to (+), to \closed", when the electrons are completely relected, from 1(2) to + (). We consider the low temperature limit, kT eV. A negative voltage eV is applied to all the norm alreservoirs and the superconductor is grounded. The voltage eV is smaller than the superconducting gap , so no single particle transport takes place. It is further assumed that the size of the system is smaller than the phase breaking length.

We ist present a simple and transparent explanation of how the entanglement is generated and detected, a rigorous derivation follows below. The superconductor emits pairs of particles into the norm alarms. Since the superconductor is a single object, coherent on a macroscopic scale, the state of an emitted pair is a linear superposition of states corresponding to a pair emitted through barrier 1 and 2. The emitted pair can either split, with one electron going to each beam splitter A and B, or both electrons can go to the same beam splitter. How ever, the latter process does not contribute [13] to the quantity of interest, the current cross-correlations, to leading order in pair emission (Andreev re ection) probability, proportional to ². The relevant part of the state of the emitted pair can thus be written as

$$ji=j_{12}i$$

a product of a state j $_{12}i$, orbitally entangled with respect to em ission across barrier 1 and 2 (called 12-space below), and a state j $_{AB}i$, describing one electron going towards A and one towards B, containing all additional information, such as energy and spin dependence. The beam splitters A and B, parameterized via the angles $_{A}$ and $_{B}$, rotate the state j $_{12}i$ [see Eq. (1)].

The entanglam ent is detected by violation of a BI.W e recall that in the original form ulation [2], a source em itting spin-1=2 singlets was considered. The BI, as form ulated in Ref. [14],

$$S = \frac{1}{E} \begin{pmatrix} a ; b \end{pmatrix} = E \begin{pmatrix} a ; b \\ a ; b \end{pmatrix} + E \begin{pmatrix} 0 \\ a ; b \end{pmatrix} + E \begin{pmatrix} 0 \\ a ; b \\ B \end{pmatrix} = 2; \quad (3)$$

is expressed in terms of spin correlation functions [15]

$$E(A; B) = P_{++} P_{+} P_{+} + P_{-}$$
 (4)

Here P ($_{A}$; $_{B}$) are the joint probabilities to observe one particle in detector A with a spin = [+() denoting up (down)] along the $_{A}$ -direction, and the other in detector B with a spin = along the $_{B}$ -direction. The joint probabilities are given by

P
$$(_{A}; _{B}) = 1 + \cos[2(_{A} _{B})]$$
: (5)

Inserting the values for P from Eq. (5) into Eq. (4), we get E ($_{A}$; $_{B}$) = cos[2($_{A}$ $_{B}$)]. We then directly see that for e.g. angles $_{A}$ = =8; $_{B}$ = =4; $_{A}^{0}$ = 3 =8 and $_{B}^{0}$ = =2, the BI in Eq. (3) is maximally violated, i.e. we get S = 2 2.

In our orbital setup (see Fig 1), it is clear from Eq. (2) that the 12-space plays the role of a pseudo-spin space and the norm al reservoirs act as detectors. We can thus, in direct analog to Ref. [2], form ulate a BI in term s of any observable which is directly proportional to the corresponding joint probability P ($_{A}$; $_{B}$) for our state j $_{12}$ i [here ; = denote the reservoirs, see Fig.

1]. We nd below that the zero-frequency current crosscorrelator is given by

S 2 dth
$$\hat{f}_{A}$$
 (t) \hat{f}_{B} (0) $i = P_{0}P$ (_A; _B); (6)

i.e. directly proportional to the joint probability distribution. Here $P_0 = e^{3}V^{-2}=2h$, and $\hat{f}(t) = \hat{f}(t)$ if i is the uctuating part of the current $\hat{f}(t)$ in reservoir . This leads to the important result that the Bell inequality in Eq. (3) can be directly formulated in terms of the zero-frequency current correlators in Eq. (6). We note [15] that when inserting S directly into the correlation functions Eq. (4) (i.e. substituting P with S), we must divide by the sum of all correlators $S_{++} + S_{+} + S_{-+} + S_{--} = P_0$, which just elim inates P_0 .

The simple result in Eq. (6) can be understood by considering the properties of the time-dependent correlator h \hat{f}_A (t) \hat{f}_B (0)i. It is nite only for times t $< \ _c$, where $_c = h=eV$ is the correlation time of the emitted pair (see Fig. 2). In the tunneling limit under consideration, 1, the correlation time is much smaller than the average time between the arrival of two pairs e=I h=eV². As a result, only the two electrons within a pair are correlated with each other, while electrons in di erent pairs are completely uncorrelated. Thus, the zero frequency current correlator in Eq. (6) is just a coincidence counting measurem ent running over a long time, collecting statistics over a large num ber of pairs [16].

For a rigorous derivation of the above result, we rst discuss the role of the superconductor as an em itter of pairs of orbitally entangled electrons. Since the system is phase coherent, we can work within the scattering approach to norm al-superconducting systems [17, 18]. The starting point is the many-body state of the norm al reservoirs, describing injection of hole quasiparticles at energies from 0 to eV (suppressing spin notation),

$$j_{in} i = \begin{cases} Y \\ y_{1A} (E) \\ y_{2A} (E) \\ y_{1B} (E) \\ y_{2B} (E) \\$$

where the ground state jDi is the vacuum for quasiparticles in the norm alreservoir. The operator $\frac{y}{1A}$ (E) creates a hole plane wave at energy E (counted from the superconducting chemical potential $_{\rm S}$) in lead 1A, going out from the norm al reservoirs towards the superconductor, and sim ilarly for the other operators. The commutation relations for the -operators are standard ferm ionic.

To obtain the state of the quasiparticles going out from the superconductor, j_{out}i, we note that the operators creating and destroying quasiparticles going out from the superconductor (towards the norm al reservoirs) are related [17, 19] to the operators of the incoming quasiparticles via a scattering matrix. The scattering am plitude for a hole injected in arm 1A, to be back-rejected as an electron in arm 2A, is denoted $r_{2A,1A}^{eh}$ and similarly for the other scattering am plitudes. In the tunnel lim it under consideration, the am plitude to backscatter as the same type of quasiparticle is r_{j,j^0}^{hh} . 1. The am plitude for Andreev re ection, r_{j,j^0}^{eh} . is given by i = 4 (independent on energy). Processes where a hole incident on barrier 1(2) is backscattered as an electron at barrier 2(1), i.e. when a pair in the superconductor breaks up, are exponentially suppressed with the distance between the two em ission points [6] and can be neglected in the present setup.

The tunneling lim it 1 m akes it relevant to change the perspective from a quasiparticle picture to an allelectron picture. An Andreev re ection, occuring with a sm all probability ², can be considered as a perturbation with respect to the ground state jDi in the normal reservoirs (a lled Ferm i sea of electrons at energies E < eV). It creates an excitation consisting of a pair of electrons (see Fig.2). Form ally, performing a B ogoliubov transform ation (i.e. $_j$ (E) = C_j^v (E), with C_j^i (E) being a standard electron creation operator), the state of the quasiparticles going out from the superconductor becomes [20] to rst order in

$$j_{out}i = jDi + j\tilde{i} + ji;$$
 (8)

where the states

$$j\hat{\ }i = \frac{i}{4} \int_{0}^{Z eV} dE \left\{ \begin{array}{c} X & X \\ X & X \\ 0 & j=1;2 = A;B \end{array} \right\} c_{j}^{Y} (E) c_{j} (E)^{5} j0;$$

$$ji = \frac{i}{4} \int_{eV}^{Z eV} h \\ dE c_{1A}^{Y} (E) c_{1B}^{Y} (E) \\ eV & i \\ + c_{2A}^{Y} (E) c_{2B}^{Y} (E) j0; \qquad (9)$$

describe orbitally entangled electron \wavepacket" pairs, i.e. a superposition of pairs at di erent energies [21]. In a rst quantized notation, the state j i is just the state in Eq. (2). We emphasize that the change from a quasiparticle to an all-electron picture, providing a clear picture of the entanglem ent, does not alter the physics.

The detection of the entanglement is done via the zero frequency current cross-correlators S in Eq. (6). To obtain S , we insert the second quantization expression for the current operator [19], $\hat{1}$ (t) = (e=h) dE dE ⁰exp[i(E E^{0})t=h]c^y (E^{0})c (E), into Eq. (6) and average with respect to the state ji in Eqs. (8) and (9). The beam splitters are taken into account by relating the c_i -operators in arm s 1A and 2A (1B and 2B) to the c -operators in the reservoirs + A A (+ B and B) via the scattering matrix in and Eq. (1). The average current, equal in all arm s , ishI i $I = (e^2 = 2h)^2 V$, independent of the beam splitter transparency.

The ground state jDi in Eq. (8) does not contribute to the correlator. Moreover, calculating the current correlator we nd that the state j~i, describing two electrons




FIG. 2: Left, upper: In the lled stream of incoming holes (empty circles) from the norm al reservoir, occasionally a hole is back-re ected as an electron (solid circles). The \m issing hole" (i.e an electron) and the Andreev re ected electron constitute the pair em itted from the superconductor. Left, lower: The correlation time $_{c} = h=eV$ (width of the wave packet) and the average time between em ission of two subse- $_{\rm c}$ of the current correlator. The quent pairs $e=I = h=eV^{2}$ $_{\rm c}$ between the em issions of the smalltime di erence h= two electrons in the pair is shown as a split of the wavepacket. Right: The transmission probabilities T_A (dashed), T_A^0 (dotted) and T_B^0 (solid) as a function of T_B [$T = \cos^2()$, giving optim alviolation of the Bellinequalities for dephasing param eters = 1 and = 0:3.

em itted into the sam e norm al lead, only contributes [13] to the cross correlator at order ⁴ and can thus be neglected. From Eq. (6) we then nd that S ($_{\rm A}$; $_{\rm B}$) = P_0P ($_{\rm A}$; $_{\rm B}$), where P is given in Eq. (5), just as announced. We note that it is the structure of j i in 12-space, i.e. j $_{12}$ i, that determ ines the angle dependence of P , all properties of j $_{\rm AB}$ i just gives rise to the constant P_0 . Moreover, the calculations show that the time dependent correlator h \hat{f}_1 (t) \hat{f}_2 (0) i vanishes as ($_{\rm C}$ =t)² for t $_{\rm C}$. As is pointed out above, S ($_{\rm A}$; $_{\rm B}$) can thus be inserted in Eq. (4) and be used to violate the BI in Eq. (3).

Until now, ideal conditions have been considered. One possible source of disturbances is dephasing. In our system, dephasing can quite generally be expressed in terms of a density matrix = [jllihl1j+ j2ih22j+ (jllih22j+ jllih22j)=2, where the o -diagonal elements, giving rise to the entanglement, are suppressed by a phenomenological dephasing parameter 0 1. The correlators E ($_A$; $_B$) in Eq. (4) then take the form

 $E (A_{A}; B_{B}) = \cos(2 A_{A})\cos(2 B_{B}) + \sin(2 A_{A})\sin(2 B_{B});$ (10)

By adjusting the four angles $_{A}$; $_{A}^{0}$; $_{B}$ and $_{B}^{0}$ we nd the maxim alBell parameter in Eq. (3) is

$$S = 2^{p} \frac{1}{1 + 2^{2}}$$
 (11)

which violates the BI for any > 0. The optimal violation angles, all in the rst quadrant, are tan (2 $_{\rm A}$) =

 $\cot(s)$, $\tan(2 \frac{0}{A}) = \tan(s)$ and $\tan(B$ $\frac{0}{B}) = \operatorname{sign}[\cos(2 \frac{1}{A})][(\tan^2(s) + \frac{1}{2})=(\frac{1}{2}\tan^2(s) + \frac{1}{2})]$ 1) $l^{1=2}$, where $s = B + B^{0}$ can be chosen at will. The corresponding transmission probabilities $T = \cos^{2}(0)$ are shown for = 1 and = 0.3 in Fig. 2.

The BI can thus in principle be violated for any am ount of dephasing. However it might be di cult to produce beam splitters which can reach all transmission probabilities between 0 and 1. This is not a serious problem in the absence of dephasing, = 1, a violation can be obtained for a large, order of unity, fraction of the \transm ission probability space". However, in the lim it of strong dephasing, 1, the set of transm ission probabilities for optim al violation contain transm issions close to both 0 and 1, see Fig. 2. Expecting unity transm ission to be most complicated to reach experim entally, we note that by instead choosing transm ission probabilities $T_A = T_B = 0$, $T_B^0 = 1=2$ and T_A^0 , the inequality in Eq. (3) becomes $2jl + T_{A}^{0}j$ 2. This gives a violation, although not maxim al, for all 1.

A part from dephasing there are several other e ects such as additional scattering phases, in purity scattering or asym m etric tunnelbarriers, which m ight alter the possibility to violate the BI. It turns out that all these e ects can be taken into account by replacing $! \, ^0 \cos(0)$ in Eq. (11), with the important conclusion that none of these e ects will destroy the possibility to violate the BI.

The phase factor $_0$ is the sum of possible scattering phases from the beam splitters [the scattering am plitudes in Eq. (1) are taken real], phases $k_{\rm F}$ L due to a difference in length, L, between the norm alarm s (see Fig. 1), scattering phases from weak in purities and a possible phase di erence between the superconductor at the two tunnel contacts 1 and 2. As a consequence e.g. the superconducting phase (in a bop geom etry) can be modulated to com pensate for the other phases.

The factor ⁰ plays the same role as dephasing. One possible contribution to ⁰ is energy dependent phases which oscillate rapidly on a scale of eV, suppressing the entangled part of the current correlator. For di erent lengths of the norm al arms, there is always a phase

E L=hv_F. This phase can however be neglected for L hv_F =eV, which for eV is fulled for L smaller than the superconducting coherence length hv_F = . Another possibility is that, due to asymmetries of the tunnel barriers $_{1} \in _{2}$, the amplitude for the process where the pair is emitted to jlli is di erent from the process where it is emitted to jlli bis di erent from the process where it is emitted to jlli $_{2}^{1}$. This gives rise to a state, in 12-space, $(_{1}$ jlli + $_{2}$ jlli = $\frac{1}{2} + \frac{2}{2}$. In this case [22] 0 = 2 $_{1}$ $_{2}$ = $(_{1}^{2} + \frac{2}{2})$. Thus, it is in principle possible to violate BI for arbitrary asymmetry.

We nally point out that the constraint on single mode beam splitter can not easily be relaxed. A multimode beam splitter will, in the suggested setup, probably have a di erent transparency dependence for the di erent modes. This decreases the space of accessible angles and will eventually make the violation of the BI impossible.

In conclusion, we have investigated a spin-independent

scheme to generate and detect two-particle orbital entanglement in a mesoscopic normal-superconductor system. The cross-correlator between the currents in the two leads depends in a nonlocal way on transparencies of beam splitters in the two leads. These nonlocal correlations arises due to the entanglement of the injected pair. For appropriate choices of transparencies, the correlators give a violation of a BI for arbitrary strong dephasing.

W e acknow ledge discussions with Fabio Taddeiand Valerio Scarani. This work was supported by the Swiss National Science Foundation and the program for Materials with Novel E lectronic Properties.

- [1] A. Einstein, B. Podolsky and N. Rosen, Phys. Rev. 47, 777 (1935); D. Bohm and Y. Aharonov, Phys. Rev. 108, 1070 (1957).
- [2] J.S.Bell, Physics 1, 195 (1964); Rev. M od. Phys. 38, 447 (1966).
- [3] For recent experim ents, see e.g. W . T ittel et al., Phys. Rev.Lett 81, 3563 (1998); G.W eihs et al., ibid 81 5039, (1998).
- [4] N.G isin et al, Rev.M od.Phys.74,145 (2002).
- [5] A. Steane, Rep. Prog. Phys. 61, 117 (1998).
- [6] P.Recher, E.V. Sukhorukov, and D. Loss, Phys. Rev. B 63, 165314 (2001).
- [7] P.Recher and D.Loss, Phys. Rev. B 65 165327 (2002);
 C.Bena et al, Phys. Rev. Lett 89, 037901 (2002).
- [8] G.Burkard, D.Loss and E.V.Sukhorukov, Phys. Rev. B
 61,16303 (2000); F.Taddeiand R.Fazio, ibid 65,134522 (2002); J.C.Egues, G.Burkard, and D.Loss, Phys. Rev. Lett. 89, 176401 (2002).
- [9] G B. Lesovik, T. M artin and G. B latter, Eur. Phys. J. B 24, 287 (2001).
- [10] N M. Chtchelkatchev et al, Phys. Rev. B 66, 161320 (2002).
- [11] The two parts of the superconductor can be contacted e.g. via a loop.
- [12] R.Liu et al., Nature 391, 263 (1998); W D.O liver et al., Science 284, 299 (1999).
- [13] P. Sam uelsson and M. Buttiker, Phys. Rev. Lett. 89, 046601 (2002); Phys. Rev. B 66, 201306 (2002).
- [14] J.F. Clauser et al., Phys. Rev. Lett. 23, 880 (1969).
- [15] A. Aspect, P. G rangier, and G. Roger, Phys. Rev. Lett. 49, 91 (1982).
- [16] In contrast to R ef. [10] there is no constraint on the m easurem ent tim e in our scheme.
- [17] S.D atta, P.F.Bagwell, and M.P.Anantram, Phys.Low-Dim.Struct.3,1 (1996); C.W. J.Beenakker, Rev.Mod. Phys, 69, 731 (1997).
- [18] The same result can be obtained using the tunneling Hamiltonian formalism in ref. [6].
- [19] M. Buttiker, Phys. Rev. B. 46, 12485 (1992); Ya. M. Blanter, and M. Buttiker, Phys. Rep. 336, 1 (2000).
- [20] For a related approach in optics, see e.g. Z.Y. Ou et al., Phys. Rev. A. 41, 566 (1990).
- [21] In contrast to R efs. [8], in our setup there is no additional constraint on the energy spectrum of the entangled electrons.
- [22] N.Gisin, Phys.Lett. A 154, 201 (1991); A.Abouraddy et.al, Phys.Rev.A 64, 050101 (2001).