

# Comment on 'Effect of on-site Coulomb repulsion on superconductivity in the boson-fermion model' (Phys. Rev. B66, 134512 (2002))

A.S. Alexandrov

Department of Physics, Loughborough University, Loughborough LE11 3TU, United Kingdom

The two-dimensional (2D) boson-fermion model (BFM) of high-temperature superconductors, numerically studied by Domanski (Phys. Rev. B66, 134512 (2002)) is not a superconductor. The critical temperature of the model is zero for any symmetry of the order parameter. The opposite conclusion advocated by Domanski stems from a mean-field approximation (MFA) neglecting the boson self-energy which is qualitatively erroneous for any-dimensional BFM.

PACS numbers: PACS: 71.20.-z, 74.20.Mn, 74.20.Rp, 74.25.Dw

Recently Domanski [1] and some other authors (see references in [1]) claimed that 2D BFM with hybridised fermions and immobile hard-core bosons is capable to reproduce the phase diagram of cuprates. The model is defined by the Hamiltonian,

$$H = \sum_{\mathbf{k}, \sigma=\uparrow, \downarrow} \xi_{\mathbf{k}} c_{\mathbf{k}, \sigma}^{\dagger} c_{\mathbf{k}, \sigma} + E_0 \sum_{\mathbf{q}} b_{\mathbf{q}}^{\dagger} b_{\mathbf{q}} + gN^{-1/2} \sum_{\mathbf{q}, \mathbf{k}} (\phi_{\mathbf{k}} b_{\mathbf{q}}^{\dagger} c_{-\mathbf{k}+\mathbf{q}/2, \uparrow} c_{\mathbf{k}+\mathbf{q}/2, \downarrow} + H.c.), \quad (1)$$

where  $\xi_{\mathbf{k}} = -2t(\cos k_x + \cos k_y) - \mu$  is the 2D energy spectrum of fermions,  $E_0 \equiv \Delta_B - 2\mu$  is the bare boson energy with respect to the chemical potential  $\mu$ ,  $g$  is the magnitude of the anisotropic hybridisation interaction,  $\phi_{\mathbf{k}} = \phi_{-\mathbf{k}}$  is the anisotropy factor, and  $N$  is the number of cells. Ref. [1] argued that 'superconductivity is induced in this model from the anisotropic charge exchange interaction ( $g\phi_{\mathbf{k}}$ ) between the conduction-band fermions and the immobile hard-core bosons', and 'the on-site Coulomb repulsion  $U$  competes with this pairing' reducing the critical temperature  $T_c$  less than by 25%. The author of Ref. [1] neglected our study of BFM [2], which revealed a devastating effect of the boson self-energy on  $T_c$ . Here I show that because of this effect  $T_c = 0K$  in the model, Eq.(1), even in the absence of the Coulomb repulsion,  $U = 0$ , and the mean-field approximation of Ref. [1] is meaningless for any-dimensional BFM.

Using MFA Ref. [1] decouples bosons and fermions in Eq.(1) replacing boson operators by  $c$ -numbers for  $\mathbf{q} = 0$ . Then  $T_c$  is numerically calculated using the linearised BCS equation for the fermionic order-parameter  $\Delta_{\mathbf{k}}$ ,

$$\Delta_{\mathbf{k}} = \frac{\tilde{g}^2 \phi_{\mathbf{k}}}{E_0 N} \sum_{\mathbf{k}'} \phi_{\mathbf{k}'} \frac{\Delta_{\mathbf{k}'} \tanh(\xi_{\mathbf{k}'} / 2k_B T)}{2\xi_{\mathbf{k}'}} \quad (2)$$

where  $E_0$  is determined by the atomic density of bosons ( $n^B$ ) as (Eq.(9) in Ref. [1])

$$\tanh \frac{E_0}{2k_B T} = 1 - 2n^B, \quad (3)$$

with  $T = T_c$  and  $\tilde{g}^2 = g^2(1 - 2n^B)$ . While Eq.(2) is perfectly correct, Eq.(3) is incorrect because the boson

self-energy  $\Sigma_b(\mathbf{q}, \Omega_n)$  due to the exchange interaction is missing. At first sight [1] the self-energy is small if  $g$  is small in comparison to the kinetic energy of fermions. However  $\Sigma_b(0, 0)$  diverges logarithmically at zero temperature [2], no matter how weak the interaction is. Therefore it should be kept in the density sum-rule, Eq.(3). Introducing the boson Green's function

$$D(\mathbf{q}, \Omega_n) = \frac{1 - 2n^B}{i\Omega_n - E_0 - \Sigma_b(\mathbf{q}, \Omega_n)} \quad (4)$$

one must replace incorrect Eq.(3) by

$$-\frac{k_B T}{N} \sum_{\mathbf{q}, n} e^{i\Omega_n \tau} D(\mathbf{q}, \Omega_n) = n^B, \quad (5)$$

where  $\tau = +0$ , and  $\Omega_n = 2\pi k_B T n$  ( $n = 0, \pm 1, \pm 2, \dots$ ). The divergent (cooperon) contribution to  $\Sigma_b(\mathbf{q}, \Omega_n)$  is given by [2]

$$\Sigma_b(\mathbf{q}, \Omega_n) = -\frac{\tilde{g}^2}{2N} \sum_{\mathbf{k}} \phi_{\mathbf{k}}^2 \times \frac{\tanh[\xi_{\mathbf{k}-\mathbf{q}/2} / (2k_B T)] + \tanh[\xi_{\mathbf{k}+\mathbf{q}/2} / (2k_B T)]}{\xi_{\mathbf{k}-\mathbf{q}/2} + \xi_{\mathbf{k}+\mathbf{q}/2} - i\Omega_n}, \quad (6)$$

so that one obtains

$$\Sigma_b(\mathbf{q}, 0) = \Sigma_b(0, 0) + \frac{q^2}{2M^*} + \mathcal{O}(q^4) \quad (7)$$

for small  $\mathbf{q}$  with any anisotropy factor compatible with the point-group symmetry of the cuprates. Here  $M^*$  is the boson mass, calculated analytically in Ref.[2] with the isotropic exchange interaction and parabolic fermion band dispersion (see also Ref.[3]), and  $\hbar = 1$ . The BCS-like equation (2) has a nontrivial solution for  $\Delta_{\mathbf{k}}$ , if

$$E_0 = -\Sigma_b(0, 0). \quad (8)$$

Substituting Eq.(7) and Eq.(8) into the sum-rule, Eq.(5) one obtains a logarithmically divergent integral with respect to  $\mathbf{q}$ , and

$$T_c = \frac{const}{\int_0^{\infty} dq/q} = 0. \quad (9)$$

Using a ‘bubble’ approximation for the self-energy Ref.[2] proved that the Cooper pairing of fermions in BFM is impossible without the Bose-Einstein condensation (BEC) of real bosons. The bubble approximation is actually exact because of the logarithmic divergence of the Cooperon diagram, as was also confirmed in Ref.[4]. Hence, the devastating result, Eq.(9) is a direct consequence of the well-known theorem, which states that BEC is impossible in 2D.

One may erroneously believe that MFA results [1] can be still applied in three-dimensions, where BEC is possible. However, increasing dimensionality does not make MFA a meaningful approximation for the boson-fermion model. This approximation leads to a naive conclusion that a BCS-like superconducting state occurs below the critical temperature  $T_c \simeq \mu \exp(-E_0/z_c)$  via fermion pairs being *virtually* excited into *unoccupied* bosonic states [5, 6]. Here  $z_c = \tilde{g}^2 N(0)$  and  $N(0)$  is the density of states (DOS) in the fermionic band near the Fermi level  $\mu$ . However, the Cooper pairing of fermions is impossible via virtual unoccupied bosonic states. It occurs only simultaneously with the Bose-Einstein condensation of real bosons in the exact theory of 3D BFM [2].

The origin of the simultaneous condensation of the fermionic and bosonic fields in 3D BFM lies in the softening of the boson mode at  $T = T_c$  caused by its hybridization with fermions. Indeed, Eq.(8) does not depend on the dimensionality, so that the analytical continuation of Eq.(4) to real frequencies  $\omega$  yields the partial boson DOS as  $\rho(\omega) = (1 - 2n_B)\delta(\omega)$  at  $T = T_c$  for  $\mathbf{q} = 0$  in any-dimensional BFM for any coupling with fermions.

Taking into account the boson damping and dispersion shows that the boson spectrum significantly changes for all momenta. Continuing the self-energy, Eq.(6) to real frequencies yields the damping (i.e. the imaginary part of the self-energy) as [2]

$$\gamma(\mathbf{q}, \omega) = \frac{\pi z_c}{4q\xi} \ln \left[ \frac{\cosh(q\xi + \omega/(4k_B T_c))}{\cosh(-q\xi + \omega/(4k_B T_c))} \right], \quad (10)$$

where  $\xi = v_F/(4k_B T_c)$  is a coherence length. The damping is significant when  $q\xi \ll 1$ . In this region  $\gamma(\mathbf{q}, \omega) = \omega\pi z_c/(8k_B T_c)$  is comparable or even larger than the boson energy  $\omega$ . Hence bosons look like overdamped diffusive modes, rather than quasiparticles in the long-wave limit [2, 3], contrary to the erroneous conclusion of Ref.[7], that there is ‘the onset of coherent free-particle-like motion of the bosons’ in this limit. Only outside the long-wave region, the damping becomes small. Indeed, using Eq.(10) one obtains  $\gamma(\mathbf{q}, \omega) = \omega\pi z_c/(2qv_F) \ll \omega$ , so that bosons at  $q \gg 1/\xi$  are well defined quasiparticles with a logarithmic dispersion,  $\omega(q) = z_c \ln(q\xi)$  [2]. As a result the boson dispersion is distributed over the whole energy interval from zero up to  $E_0$ , but not a delta-function at  $E_0$  even in the weak-coupling limit.

The main mathematical problem with MFA stems from the density sum rule, Eq.(5) which determines the chemical potential of the system and consequently the bare boson energy  $E_0(T)$  as a function of temperature. In

the framework of MFA one takes the bare boson energy in Eq.(2) as a temperature independent parameter,  $E_0 = z_c \ln(\mu/T_c)$ , or determines it from the conservation of the total number of particles, Eq.(5) neglecting the boson self-energy  $\Sigma_b(\mathbf{q}, \Omega)$  [1, 5, 6]). Then Eq.(2) looks like the conventional mean-field BCS equation, or the Ginzburg-Landau equation (near the transition) with a negative coefficient  $\alpha \propto T - T_c$  at  $T < T_c$  in the linear term with respect to  $\Delta(T)$ . Hence, one concludes that the phase transition is almost the conventional BCS-like transition, at least at  $E_0 \gg T_c$  [5, 6], and, using the Gor’kov expansion in powers of  $\Delta$ , finds a finite upper critical field  $H_{c2}(T)$  [8]. These findings are mathematically and physically wrong. Indeed, the term of the sum in Eq.(5) with  $\Omega_n = 0$  is given by the integral

$$T \int \frac{d\mathbf{q}}{2\pi^3} \frac{1}{E_0 + \Sigma_b(\mathbf{q}, 0)}. \quad (11)$$

The integral converges, if and only if  $E_0 \geq -\Sigma_b(0, 0)$ . In fact,  $E_0 + \Sigma_b(0, 0)$  is strictly zero in the Bose-condensed state, because  $\mu_b = -[E_0 + \Sigma_b(0, 0)]$  corresponds to the boson chemical potential relative to the lower edge of the boson energy spectrum. More generally,  $\mu_b = 0$  corresponds to the appearance of the Goldstone-Bogoliubov mode due to a broken symmetry below  $T_c$ . This exact result makes the BSC equation (2) simply an identity [2] with  $\alpha \equiv 0$  at any temperature below  $T_c$ . On the other hand, MFA violates the density sum-rule, predicting the wrong negative  $\alpha(T)$  below  $T_c$ .

Since  $\alpha(T) = 0$ , the Levanyuk-Ginzburg parameter [9] is infinite,  $Gi = \infty$ . It means that the phase transition is never a BCS-like second-order phase transition even at large  $E_0$  and small  $g$ . In fact, the transition is driven by the Bose-Einstein condensation of *real* bosons with  $\mathbf{q} = 0$ , which occur due to the complete softening of their spectrum at  $T_c$  in 3D BFM. Remarkably, the conventional upper critical field, determined as the field, where a non-trivial solution of the linearised Gor’kov equation occurs, is zero in BFM,  $H_{c2}(T) = 0$ , because  $\alpha(T) = 0$  below  $T_c$ . It is not a finite  $H_{c2}(T)$  found in Ref. [8] using MFA. Even at temperatures well below  $T_c$  the condensed state is fundamentally different from the BCS-like MFA ground state, because of the *pairing* of bosons [10]. The pair-boson condensate significantly modifies the thermodynamic properties of the condensed BFM compared with the MFA predictions.

This qualitative failure of MFA might be rather unexpected, if one believes that bosons in Eq.(1) play the same role as phonons in the BCS superconductor. This is not the case for two reasons. The first one is the density sum-rule, Eq.(5), for bosons which is not applied to phonons. The second being that the boson self-energy is given by the divergent (at  $T = 0$ ) Cooperon diagram, while the self-energy of phonons is finite at small coupling.

I have to conclude that the numerical work by Doman-ski [1] does not make any sense in any dimension. There is nothing in 2D BFM to compete with because the model

is not a superconductor even without the Coulomb repulsion. The MFA results [1, 8] do not make any sense in three dimensions either, because the divergent self-energy has been neglected in calculating  $T_c$  and  $H_{c2}(T)$ .

The common wisdom that at weak coupling the boson-fermion model is adequately described by the BCS theory, is negated by our results.

- 
- [1] T. Domański, Phys. Rev. **B66**, 134512 (2002).
  - [2] A. S. Alexandrov, J. Phys.: Condens. Matter **8**, 6923 (1996); Physica **C274**, 237 (1997); *ibid* **316**, 239 (1999).
  - [3] I. Grosu, C. Blaga, and M. Crisan, J. Supercond. **13**, 459 (2000).
  - [4] R. Friedberg, H. C. Ren and O. Tchernyshyov, J. Phys.: Condens. Matter **10**, 3089 (1998); A. S. Alexandrov, J. Phys.: Condens. Matter **10**, 3093 (1998).
  - [5] R. Friedberg and T.D. Lee, Phys. Rev. **B40**, 6745 (1989); R. Friedberg, T.D. Lee, and H.C. Ren, Phys.Rev. **B42**, 4122 (1990).
  - [6] R. Micnas, J. Ranninger, and S. Robaszkiewicz, Rev. Mod. Phys. **62**, 113 (1990); J. Ranninger and J.M. Robin, Physica **C235**, 279 (1995).
  - [7] J. Ranninger, J.M. Robin, and M. Eschrig, Phys. Rev. Lett. **74**, 4027 (1995).
  - [8] T. Domański, M. M. Maška and M. Mierzejewski, Phys. Rev. B **67**, 134507 (2003).
  - [9] L.D. Landau and E.M. Lifshitz, *Statistical Physics, 3rd Edition, Part 1* (eds. E.M. Lifshitz and L.P. Pitaevskii, Pergamon Press, Oxford- New York (1980)), page 476.
  - [10] A.S. Alexandrov, cond-mat/0402123, unpublished.