M ixing e ects for the structural relaxation in binary hard-sphere liquids

G.Fo,W.Gotze, F.Sciortino,P.Tartaglia, and Th.Voigtm ann Dipartimento di Fisica and INFM Center for Statistical Mechanics and Complexity,

Universita di Roma \La Sapienza", Piazzale Aldo Moro 2, 00185 Roma, Italy

(Dated: April 14, 2024)

We report extensive molecular-dynam ics simulation results for binary mixtures of hard spheres for di erent size disparities and di erent mixing percentages, for packing fractions up to 0:605 and over a characteristic time interval spanning up to ve orders in magnitude. We explore the changes in the evolution of glassy dynam ics due to mixing and discover two opposite scenarios: for large size disparity, increasing the mixing percentage of sm all particles leads to a speed-up of long-time dynam ics, while sm all disparity leads to a slowing down. These results agree with predictions based on the mode coupling theory for ideal glass transitions.

PACS num bers: 64.70 Pf, 82.70 Dd

This paper deals with structural relaxation in liquids, i.e. the dynam ical phenom ena which are precursors of the glass transition. These processes are characterized by rather complex dependencies of correlation functions on time and parameters like the density and the tem – perature. The introduction of several new experimental techniques and great progress in molecular dynamics sim – ulation studies has unfolded a wealth of facts about the evolution of structural relaxation upon cooling or com – pressing glass-form ing liquids [1, 2].

To establish an understanding of structural relaxation and the glass transition, one should focus on system s as simple as possible. A tom ic one-component system s with conventional interaction potentials cannot be used since they crystallize before structural relaxation is fully developed. Beginning with the work of H ansen and collaborators [3], binary m ixtures of simple particles were used form olecular dynam ics studies of the glass transition. A binary Lennard-Jones system has been used extensively in recent years to analyze structural relaxation [4].

In previous work, m ixing was introduced merely as a means of suppressing crystallization [5]. In the following we analyze system atically the e ect of mixing on structural relaxation to identify the in uence of composition changes and variation of the particle size disparity on the glassy dynamics. We focus on binary hard-sphere mixtures (HSM), i.e. particles interacting via an hard-core potential, which we study via extensive molecular dynamics simulations.

A speci c m otivation for our studies com es from the light-scattering data by W illiams and van M egen for a H SM of colloidal particles [6]. The system was prepared to approximate a binary H SM for the ratio = 0.60 of the particle diam eters. Three m ixing e ects have been reported. If the percentage of the smaller particles increases from 10% to 20% of the relative packing fraction, then (i) the time scale for the naldecay of the density correlators decreases; (ii) the plateau value for interm ediate times increases; (iii) the initial part of the structural relaxation slows down. The rst e ects m eans that m ix-

ing has stabilized the liquid as if the smaller particles provide som e lubri cation. This e ect has som e analog in the plasticization observed in dense polymeric liquids due to mixing with polymers of shorter lengths. However, the e ects (ii) and (iii) indicate a stiening of the dynam ics upon mixing. These e ects have never been reported for conventional system s, and one might wonder whether these are structural relaxation phenomena rather than colloid-speci c features caused by, e.g., hydrodynam ic interactions or polydispersity. Since our simulation studies are done im plementing a New tonian microscopic dynam ics, this question will be answered by our results.

A nother m otivation of our work is provided by recent predictions for binary HSM based on mode-coupling theory (MCT) [7] calculations. This theory, which allows rst principle evaluations of density- uctuation correlators within a regime where structural relaxation dom inates the dynam ics, explains the light-scattering data for slightly polydisperse hard-sphere colloidal suspensions [8]. Extending MCT to binary HSM, the three mentioned mixing e ects have been identied as structural relaxation properties [9]. The light scattering data = 0:60 m ixture could be described quantitafor the tively to a certain extent [10]. How ever, surprisingly, the theory predicts two di erent scenarios. The speed-up of the dynam ics reported in Ref. [6] was found only for sufciently large size disparity, say a size ratio . 0**:**65. For & 0.8, the opposite e ect was predicted, i.e., m ixing slows down the dynam ics and the ideal-glass critical packing fractions decreases. The data reported in this letter con m s these astonishing predictions.

We perform standard constant-energy molecular dynam ics [11] simulation for a binary mixture of 1237 hardsphere particles. The two species (A and B) have diam – eters d_A and d_B respectively, with d_A d_B . The masses of the two species are taken as equal, thus all particles have the same therm all velocity, denoted as v_{th} . Units of length and time are chosen such that $d_A = 1$ and $v_{th} = 1$. We use the size ratio $= d_B = d_A$ as a control parameter specifying the size disparity. To model a large size disparity, we choose = 0:60. To model systems of small size disparity we study the value = 0:83. For each of the two values, we study two di erent values of the relative packing fraction of the small species, $x = '_B = ('_A + '_B)$, where ' = (=6)% d³. For the = 0:60 system, we have studied x = 0:10 and x = 0:20, corresponding to a fraction of B particles equal to 34% and 54%. For the

= 0.83 system, we have studied the cases x = 0.276and x = 0.37, corresponding to having 40% and 50% of B particles. For each and x values, we study several values of the total packing fraction $' = '_{A} + '_{B}$, covering a region where dynam ics slows down by 4 decades. D uring the long simulation runs, we check for crystallization by monitoring the time evolution of the pressure and by visual inspection of the con gurations. We also evaluate the structure factor to make sure that no crystalline peaks have developed.

The ' dependence of the di usion coe cient, evaluated from the long tim e lim it of the m ean square displacem ent [12], is shown in Fig. 1. Note that the variation of the di usivities extends over m ore than four decades.

It is important to separate mixing e ects already observed in norm al liquid states from mixing e ects which are peculiar of the structural relaxation dynamics. In hard-spheres, mixing e ects for the norm al-liquid dynam ics are qualitatively explained within Enskog's kinetic theory. The di usion constant D of the hard-sphere system (HSS) is expressed in terms of Enskog's collision rate , $D^{E} = 3v_{th}^{2} = (2)$. This rate modi es Boltzm ann's collision rate for dilute gases by the contact value g_{AA} $= 4^{P} v_{th}^{2} \otimes_{A} g_{AA} d_{A}^{2}$ of the pair-distribution function: [12]. Enskog's form ula is readily generalized to m ixtures with equal constituents' masses, giving $D^E = 3v_{tb}^2 = (2)$, $% g d^2 \cdot H ere g denotes the$ with $= 4^{-1}$ v_{th}^2 pair-distribution function of the colliding pair and for the distance at contact, d = (d + d) = 2. Results for D^E are included in Fig. 1 as lines for $\prime < 0.39$, using values from Percus-Yevick theory for g . The relevant norm al-liquid m ixing e ects for the HSM, as given by the Enskog theory, can be sum marized in the following: (i) D_A decreases and D_B increases upon decreasing ; (ii) both D $_{\rm A}\,$ and D $_{\rm B}\,$ decrease upon increasing x; (iii) all D $_{\rm A}\,$ values are smaller than all the $D_{\rm B}$ values. A detailed discussion of norm alm ixing e ects for an approxim ation to HSM has been published in Ref. [13].

In agreem ent with the pioneer work of A lder et al. for the HSS [14], the data for D are described correctly on a 30% accuracy level by the Enskog theory for ' < 0:40. U pon increasing ' beyond 0:40, the cage e ect starts dom inating and leads to a suppression of D and D below the corresponding Enskog values. Still we nd that up to the large value ' = 0:48 there are only norm al-liquid m ixing e ects since the log D -versus-' curves do not exhibit crossings.

Increasing ' further, the curves in Fig.1 get a stronger bending, i.e., the cage e ect enhances. If ' reaches




FIG.1: Diusivities D , = A; B, for binary hard-sphere m ixtures with size ratios $= d_B = d_A = 0.60$ (open symbols) = 0:83 (lled symbols); packing contributions of the and smaller spheres $x = '_B = '$ as in the legend. (a) Results for the two dierent with = A (circles) and = B (squares). The solid lines for ' < 0.39 are the Enskog results for D_B (= 0:6), D_B (= 0:83), D_A (= 0:83), and D_A (= 0:6), from top to bottom . D otted lines dem onstrate power-law ts with exponents (;x), see text for details. The inset show sD = D $^{1=}$ to demonstrate the extrapolation to zero di usivity. (b) Variation of D $_{\rm A}\,$ upon changes of x for both values; circles (triangles) refer to the higher (lower) x value studied.

0:56, the di usivities are about one order of magnitude smaller than in a conventional liquid. W ithin the interval 0:48 < ' < 0:56, there occur crossings of the curves, i.e. the mixing properties change qualitatively. The crossing observed in Fig. 1 demonstrates a rst remarkable feature of structural relaxation, viz., di erently from Enskog predictions, di usion of B particles for = 0:83 becomes smaller than D_A for the = 0:60 case. For ' > 0:56, a new pattern evolves, showing the mixing e ects for the fully developed structural relaxation. To support the association of the slowing down of the dynamics to the approach towards a glass transition, the




FIG.2: Normalized density-correlation functions for binary hard-sphere m ixtures at a total packing fraction ' = 0.600 for particles with diam eter ratio = 0.60. The full and dashed lines present m ixtures with relative packing contributions of the sm aller particles B of x = 0.20 and x = 0.10, respectively. U pper and lower panel exhibit the results for wave vectors $qd_A = 5$ and $qd_A = 10$, respectively. The results for the ones for the particles B have been shifted down by 0.3 relative to the ones for the particles A.

di usivities have been tted by a power-law function, $D = (;x)['^{c}(;x) ']^{(x)}$. A s shown in Fig.1, the power-law t can account well for the di usivity changes over m ore than two orders of m agnitude. For the four m ixtures studied, we nd that the critical packing fraction of species A deviates from that of species B by not m ore than 0.0008, so that we can con dently conclude, in agreem ent with M CT predictions, that each m ixture exhibits a single critical packing fraction 'c(;x). In the asymptotic lim it '! 'c(;x), the M CT exponent should be the same for D_A and D_B . This is not the case for our t results, as also found in previous studies [15], possibly because of preasymptotic correction e ects.

The presence of crossings in the di usivity curves on entering the region where structural relaxation becomes relevant clearly shows that dynamics for = 0.83 is signi cantly shower than the one for = 0.60. Moreover

di erently from the regime of norm al-liquid dynamics where the diusivities D (;x) exhibit the same trend with changes of x for = 0.83 and = 0.60 there is a qualitatively di erent x-dependence of the long-time dynam ics for sm all and large size disparity, as recently predicted by MCT [9]. As shown in Fig. 1b, while at = 0:83 dynam ics becom es slower on increasing x, the opposite behavior is observed at = 0:60. This second remarkable feature the plasticization phenom enon alluded to before and exhibited by the = 0.60 sim ulation data is clearly shown in Fig.1b. Consistently with these ndings, at = 0:60, the critical packing fraction increases on going from x = 0:10 (" (x = 0:10) = 0:6139 0:0004) to x = 0.20 ($'^{c}(x = 0.20) = 0.6169 = 0.0004$). The critical packing fractions for the system with = 0.83 show the opposite trend: they decrease on increasing x from x = 0.276 ($'^{c}(x = 0.276) = 0.5881$ 0.0004) to x = 0.37 $('^{c}(x = 0.37) = 0.5877 \quad 0.0004)$. This third rem arkable nding m eans that m ixing stabilizes the glass for the system with small size disparity and destabilizes it for large size disparities. Dierent from the plasticization e ect discovered by W illiam s and van M egen for = 0.60, for = 0:83 an increase of the concentration of the smaller m inority particles leads to a slowing down of the longtim e density- uctuation dynam ics. The di erent scenarios for the long-tim e relaxation scales are dem onstrated also in Figures 2 and 3. They show the x dependence of the density auto-correlation functions (q;t) for = A and B at two di erent wave vectors, below $(qd_A = 5)$ and above $(qd_A = 10)$ the rst peak of the structure factor $(qd_A = 7)$, respectively.

W ithin MCT, the ideal glass states are characterized by arrest of the density uctuations. W ithin the liquid state, for ' below but close to '°, the (q;t)-versuslogt curves exhibit a plateau at f° (q), the so-called critical D ebye-W aller factor. The curves are the atter the sm aller '° ', and the length of the plateau increases. This plateau, an outstanding feature of structural relaxation, is clearly exhibited in Figs. 2 and 3. The whole structural-relaxation process consists of two steps. The rst step deals with the relaxation towards the plateau f° (q). The second step is the long-time process dealing with the decay of (q;t) from the plateau to zero. The three mixing features discussed above concerned the time scale of the second relaxation step.

The results for $_{A}$ (q;t) shown in Figs. 2 and 3 for $qd_{A} = 5$ exhibit a fourth remarkablem ixing e ect: upon increasing x, the plateaus of the correlators increase. A c-companied with this is a attening of the $_{A}$ (q;t)-versus-logt curve in the plateau region. An indication of the same e ect is also shown by the $_{B}$ (q;t). These ndings are in accord with the two m ixing anom alies discovered in the experimental study of the colloidal = 0:60 system [6]. Our results show that they are structural-relaxation phenomena that occur also in system s with Newtonian dynamics, and that they occur for small as well as for




FIG. 3: As in Fig. 2, but for a packing fraction ' = 0.582, size ratio = 0.83, and relative packing contributions of sm all particles x = 0.37 (full lines) and x = 0.276 (dashed lines).

large size disparity of the particles. We also note that on increasing x at = 0.6, the increase of the height of the plateau value, associated with the speed-up of the dynamics, forces the relaxation curves for the same ' but for di erent x to cross at the beginning of the second relaxation step. For = 0.83, an increase of x at xed ' generates a slower dynamics and hence there is no crossing of the curves in the time interval for the second relaxation step.

To conclude, we have studied structural relaxation of four binary hard-spherem ixtures via m olecular dynam ics simulations over dynam ic ranges extending over about ve orders of magnitude. Surprising mixing e ects for the slow dynam ics have been identied. In particular, we have shown that increasing the mixing percentage of the smaller minority particles can lead to a speeding up as well as to a slowing down of the long-time decay processes, depending on whether the size disparity is large or small, respectively. There is also an increase of the plateau of the density-autocorrelation functions for small and intermediate wave vectors relation functions, which pose a challenge to theories of the glass transition, show in particular, that the description of a glass-form ing m ixture by an e ective one-component liquid cannot be possible for all properties of interest. The reported results also con rm the conclusions arrived at in a lightscattering study of a quasi-binary colloidal suspension [6], and predictions within m ode-coupling theory [9]. Finally, our work suggests that m ode-coupling theory can contribute to an understanding of qualitative trends for the m icroscopic details in the dynam ics of glass-form ing liquids.

W G. and Th.V. thank their colleagues from the University of Rome for their kind hospitality during the time this work was performed. Our collaboration was supported by DYGLAGEMEM and the Deutsche Forschungsgem einschaft through DFG grant G o 154/12-1. GF., FS. and P.T. acknow ledge support from MIUR Prin and Firb and INFM Pra-Genfit. We thank S.Buldyrev for providing us the MD code for HSM.

On leave from : Physik-D epartment, Technische Universitat M unchen, 85747 G arching, G erm any

- Proceedings of the 4th International Discussion M eeting on Slow Relaxations in Complex Systems, J. Non-Cryst. Solids, 307 (310 (2002).
- [2] Proceedings of the 3rd W orkshop on Non-Equilibrium Phenomena in Supercooled Fluids, Glasses, and Amorphous M aterials, J. Phys.: Condens. M atter, 15(11) (2003).
- [3] J. N. Roux, J.-L. Barrat, and J.-P. Hansen, J. Phys.: Condens. M atter 1, 7171 (1989).
- [4] W.Kob (2003), cond-m at/0212344, Les Houches Summer School of Theoretical Physics, Session LXXVII, in print.
- [5] S.I.Henderson, T.C. Mortensen, G.M. Underwood, andW. van Megen, Physica A 233, 102 (1996).
- [6] S.R.W illiam s and W .van Megen, Phys. Rev. E 64, 041502 (2001).
- [7] W. Gotze, in Liquids, Freezing and Glass Transition, edited by J.P.Hansen, D.Levesque, and J.Zinn-Justin (North Holland, Am sterdam, 1991), Les Houches Summer Schools of Theoretical Physics, pp.287{503.
- [8] W . van Megen, Transp. Theory Stat. Phys. 24, 1017 (1995).
- [9] W .Gotze and Th.Voigtmann, Phys. Rev. E 67, 021502 (2003).
- [10] Th. Voigtm ann (2003), to be published.
- [11] D. C. Rapaport, The Art of Molecular Dynamics Simulation (Cambridge University Press, Cambridge, UK, 1997).
- [12] J.P. Boon and S. Yip, Molecular Hydrodynamics (M cG raw Hill, New York, 1980).
- [13] H.M. Schaink and C.Hoheisel, Phys. Rev. A 45, 8559 (1992).
- [14] B. J. Alder, D. M. Gass, and T. E. Wainwright, J.Chem. Phys. 53, 3813 (1970).
- [15] W . Kob and H. C. Andersen, Phys. Rev. Lett. 73, 1376 (1994).