

Dynamical domain walls and spin-Peierls order in doped antiferromagnets: evidence from exact diagonalization of small clusters

R. Eder¹ and Y. Ohta²

¹Institut für Festkörperphysik, Forschungszentrum Karlsruhe, 76021 Karlsruhe, Germany

²Department of Physics, Chiba University, Chiba 263-8522, Japan

(dated: April 14, 2024)

The hole-doped 2D t - J model is studied by exact diagonalization on a 5×4 cluster which, unlike the standard tilted square clusters, can in principle accommodate an antiphase domain wall. For hole concentration 10% and $J=t=0.5$ the ground state energy/site is lower than the conventional tilted square 20 - 20 -cluster. In the ground state two holes form a loosely bound pair pinned to an antiphase domain wall. The dynamical density correlation function shows sharp quasiparticle-like peaks, reminiscent of the holons' in 1D chains, which suggest the existence of soliton-like propagating domain walls. The dynamical correlation function of the bond-singlet operator has a low-energy peak structure characteristic of columnar spin-Peierls order, the dynamical spin correlation function shows an intense and isolated 'resonance peak' near (π, π) .

PACS numbers: 74.20.Mn, 74.25.Dw

The presence of stripe-like structures [1] which were predicted early on by mean-field studies of Hubbard-like models [2], is the basic idea underlying much of the recent theoretical work [3, 4, 5, 6, 7] on cuprate superconductors. An interesting question is whether the 'pure' 2D t - J model, which still represents the simplest theoretical description of the CuO_2 planes, develops any stripe-like structures, either in its ground state or in low-energy excited states. The most conclusive evidence for the spontaneous formation of such stripes in the 2D t - J model is probably the density-matrix renormalization group (DMRG) study of White and Scalapino [8] and the related work by Martins et al. [9]. Exact diagonalization (ED) of small clusters [10] on the other hand, so far has not produced really compelling evidence for hole stripes, at least in the physical parameter regime. Evidence for stripe-like hole-density correlations from ED calculations has been reported by Pelovshek and Xotos [11], but only for relatively large values of $J=t=1.5$. This is astonishing, because alle effective interactions in the t - J model are expected to be short ranged, whence any strong tendency to form stripe-like structures should make itself felt even in small lattices - unless it is suppressed by finite-size effects.

As already stressed by White and Scalapino [8] a potential source of such finite-size effects are the boundary conditions imposed by the cluster geometry. Most studies to date have been performed on tilted-square clusters [10] with periodic boundary conditions (PBC), which are chosen such as to accommodate the Neel-type order parameter in the undoped system. On the other hand, such a geometry is precisely the wrong one when an antiphase domain wall is present in the ground state. The mere geometry of the cluster then can frustrate the domain wall, thus artificially enforcing a homogeneous ground state. An ideal system to check this would be the 5×4 -cluster with PBC, which on one hand is appropriate to accommodate an antiphase domain wall parallel to the y -direction and on the other hand can be compared di-




FIG. 1: Difference ΔE_0 between the GS energy of the 5×4 and the 20×20 cluster, as function of $J=t$.

rectly to the tilted square 20 -site cluster. As will be demonstrated, this cluster shows very clear and unambiguous evidence for a stripe-like domain wall not only in the ground state, but also in the form of excited states corresponding to a soliton-like propagating domain wall. The 2D t - J model thus has a very strong intrinsic tendency to form stripes - if it is allowed to do so. Moreover, the formation of stripes seems intimately related to spin-Peierls-like columnar singlet order [6, 12].

The Hamiltonian of the t - J model reads

$$H = t \sum_{\langle hi; ji \rangle} c_i^\dagger c_j + H_{sc} + J \sum_{\langle hi; ji \rangle} S_i \cdot S_j - \frac{n_i n_j}{4}$$

Here $\langle hi; ji \rangle$ denotes summation over nearest neighbor pairs, $c_i^\dagger = c_i + (1 - n_i) \tau_y$ and S_i and n_i denotes the operators of electron spin and density at site i , respectively. We begin with a comparison of the ground state (GS) energy for the 5×4 cluster and the tilted square 20 -site cluster. One might expect that this can give an indication, as to whether a given cluster has the proper geometry to describe the system. Inappropriate boundary conditions introduce 'frustration' thereby raising the energy. For example, at half filling the GS energy in 5×4 is $1.165J$ /site, whereas in the square-shaped cluster it is $1.191J$ /site, the higher energy for 5×4 obviously

arXiv:cond-mat/0304244v1 [cond-mat.str-el] 24 Apr 2003

being due to the frustration of the (quasi-) Neel order along the odd-numbered side of this rectangular cluster. Interestingly enough, this changes when holes are added. Figure 1 compares the GS energy of the two clusters with 2 holes as function of $J=t$. For $J=t > 0.5$ the rectangular cluster indeed has the lower energy, indicating that for the doped case the different boundary conditions are at least as reasonable to represent the bulk system as is the square cluster. All data to be presented below have been obtained for the value $J=t = 0.5$ and 2 holes. Table I shows the static density and spin correlation functions for the 5×4 cluster with 2 holes. These are defined as the GS expectation values $g_D(\mathbf{R}) = \langle \prod_j n_{j+n_j+\mathbf{R}} \rangle$ and $g_S(\mathbf{R}) = \langle \prod_j S_j \cdot S_{j+\mathbf{R}} \rangle$. There is pronounced anisotropy in $g_D(\mathbf{R})$, particularly so at short distances, which immediately suggests a 'hole stripe' in y-direction. The fact that the hole density correlation is strongly directional might seem to suggest that this state is related to the p-like pairing states, which usually form the first excited state above the $d_{x^2-y^2}$ -like ground state (and sometimes in fact the ground state itself) [13] of two holes in square clusters. These p-like pairs, however, are spin triplets whereas the present ground state is a spin singlet which moreover is even under reflection by both the x- and y-axis. This state therefore is qualitatively very different from those seen so far in square-shaped clusters. Pronounced anisotropy is also seen in the spin-correlation function, where in particular the nearest neighbor spin correlation in y-direction (i.e. parallel to the stripe) exceeds the one in x-direction by 50%. While some anisotropy in the correlation functions is to be expected solely due to the rectangular shape of the 5×4 cluster, the edges differ in length by only 20% which seems to be rather small to explain the strong anisotropy.

To further clarify the nature of the ground state, Figure 2 shows the 'representatives' of those basis states, which have the largest weight in the GS wave function, labelled by their coefficients in the GS. Taken together, these 9 states exhaust 14.1% of the GS wave function - which is a lot, given that there are all in all 59000 basis states for this system. For clarity we mention that a basis state for the ED scheme is created from the 'representative' by a) translating it by all N different lattice vectors of the cluster, b) applying all N^0 point group operations of the little group of the total momentum \mathbf{k} c) applying the time reversal operator and d) adding up the resulting $2N \times N^0$ states with the proper phase factors so as to create a state with prescribed momentum, point group symmetry and

	$g_D(\mathbf{R})$			$g_S(\mathbf{R})$		
$R_y = 2$	0.298	0.139	0.035	0.127	-0.110	0.032
$R_y = 1$	0.239	0.144	0.041	-0.299	0.110	-0.040
$R_y = 0$	2.000	0.049	0.018	0.675	-0.198	0.034
	0	1	2	0	1	2
	R_x			R_x		

TABLE I: Static correlation functions for the 4×5 cluster.




FIG. 2: Hole-spin configurations of the representatives with the largest weight in the GS wave function. Dashed and solid boxes indicate the hops and spin flips to create the state from the 'seed state' in the top left panel.

parity under time reversal. All states in Figure 2 show a domain wall separating two perfect Neel states. In the configuration with the largest weight (top left) the two holes form a pair parallel to the domain wall. All subsequent states can be generated from this 'seed state' either by hole hopping, by a quantum spin flip along the domain wall, or by a combination of both. This is very much reminiscent of 'string picture' theories for holes populating a domain wall between two Neel states [14, 15] and in particular highlights the importance of charge fluctuations transverse to the stripe [14, 15]. Taken together the data presented so far show that the 5×4 cluster has a qualitatively new (as compared to the tilted square clusters) ground state: a loosely bound hole pair pinned to an antiphase domain wall of the staggered magnetization, with strong quantum spin fluctuations along the fault line. Having established the nature of the ground state, we consider the excitation spectra of the system. As a first step, we address the existence of dynamical domain walls. Here the appropriate quantity to look at is the dynamical density correlation function (DDCF), defined as

$$D(\mathbf{q}; \omega) = \frac{1}{N} \sum_{\mathbf{R}} \frac{\langle \prod_j n_{j+\mathbf{R}} \rangle \langle \prod_j n_{j+\mathbf{R}} \rangle}{(\mathcal{E} - \mathcal{E}_0 - i0^+)}; \quad (1)$$

where $n_{\mathbf{q}} = \frac{1}{N} \sum_j n_j e^{i\mathbf{q} \cdot \mathbf{R}_j}$ is the Fourier transform of the electron-density operator n_j . Figure 3 compares $D(\mathbf{q}; \omega)$ for the 5×4 and the tilted square 20-site cluster. Whereas the DCF for the tilted square cluster is essentially incoherent [16], the DCF for the rectangular cluster shows intense and well defined peaks at the lower edge of the spectra, particularly so at $\mathbf{q} = (2\pi/5; 0)$ and $\mathbf{q} = (4\pi/5; 0)$. In fact, it is tempting to compare the DCF for the 4×5 cluster to the one for 1D chains. Figure 3 also shows the DCF for an 11-site t-J chain with one hole. The DCF here really is an almost ideal single-peak spectrum, with a dispersion that can be fitted very accurately by the expression $\omega(\mathbf{q}) = 2t(1 + \cos(\mathbf{q}))$. Obviously




FIG. 3: Left: DCF for the 4×5 and $\frac{p}{20} \times \frac{p}{20}$ cluster. The upper/lower momentum label refer to $4 \times 5 / \frac{p}{20} \times \frac{p}{20}$. Right: DCF for a single hole in a 1-D 11-site ring at $J=\tau=0.5$.

these peaks originate from the particle-hole excitations of the single holon' in the system. Since the holon is nothing but a domain wall, this suggests that the sharp peaks in the 4×5 DCF form momenta $(q_x; 0)$ also originate from the propagation of the domain wall as a whole. The stronger q -dependence of the peak weight in the 2D cluster is probably due to the finite width of the domain wall: whereas the holon really is a point-like object, making its density structure factor $g_0(q)$ at in q -space, the domain wall in 2D has some extension in x -direction, whence its structure factor must have a q_x -dependence.

Next, we focus on the anisotropic spin correlations. Read and Sachdev [17] have proposed a spontaneous spin-Peierls-like dimerization to occur as a general feature of $S = \frac{1}{2}$ antiferromagnets in 2D – which would explain the anisotropy of $g_S(R)$. The appropriate quantity to check this hypothesis is $B(q; !)$, the dynamical correlation function (defined as in (1)) of the bond-singlet operator $B_{\mu q} = \frac{1}{N} \sum_j S_j S_{j+e_{\mu}} e^{iq \cdot R_j}$. Thereby

$\mu \in \{x, y\}$ denotes the direction of the singlet-bond. A also of interest is $S(q; !)$, the dynamical correlation function of the spin operator $S_q = \frac{1}{N} \sum_j S_j e^{iq \cdot R_j}$. These spectra are shown in Figure 4. To begin with, $B(q; !)$ in 4×5 has a prominent low energy peak for both singlet directions, $\mu = x, y$, at $q = (2\pi/5; 0)$. This peak (as well as the similar peaks at $(0; \pi/2)$ and $(4\pi/5; 0)$) is simply a replica of the intense low energy peak seen at the same q in the DCF. Just as the density operator itself, $B_{\mu q}$ is a number-conserving spin singlet, whence $B(q; !)$ probes exactly the same manifold as the DCF. Since the bond singlet operator (partially) couples to fluctuations of the electron density, $B(q; !)$ must pick up the signal from the propagating domain wall. This interpretation is confirmed by the fact, that the excitation energies $E = E_0$ of the respective peaks




FIG. 4: Left: $B(q; !)$ and $S(q; !)$ in the 4×5 cluster with two holes. Right: The same correlation functions for the $\frac{p}{20} \times \frac{p}{20}$ cluster. $S(q; !)$ is multiplied by 0.5.

in the DCF and $B(q; !)$ agree to computer accuracy, i.e. these peaks originate from the same manifold. More interesting therefore is the second high-intensity and low energy peak, at $(0; \pi)$. This peak appears only in $B_y(q; !)$, that means for bonds parallel to the domain wall. The wave vector $(0; \pi)$ for bonds in y -direction corresponds exactly to the Spin-Peierls order found by Read and Sachdev [17] and proposed to be a general feature of doped antiferromagnets by various workers [6, 12]. This peak has a lower excitation energy than the dominant peak in the spin correlation function $S(q; !)$ at $(\frac{4\pi}{5}; \pi)$, indicating that spin-Peierls order is the most likely instability of the system. $S(q; !)$ itself is quite remarkable, especially when we compare it to the tilted square cluster. There $S(q; !)$ shows a series of low energy peaks near $q = (\pi; \pi)$, which suggest a relatively smooth dispersion with a shallow minimum at the incommensurate wave vector $(\frac{3\pi}{5}; \frac{4\pi}{5})$. $S(q; !)$ in 4×5 , on the other hand, consists of a rather diffuse high-energy continuum and an isolated low-energy peak at $(\frac{4\pi}{5}; \pi)$. The dispersion of the spin excitations (if it exists) must be rather discontinuous near this momentum. Such a spin-excitation spectrum is very much reminiscent of the 'resonance peaks' observed in neutron scattering experiments on the superconducting state [18]. In the framework of the spin-Peierls scenario, the peak should be interpreted as follows: starting from a columnar spin-Peierls state with singlets in y -direction, a Néel-ordered state can be generated by condensation of (Bosonic) bond-triplet excitations [19] with momentum $Q = (\pi; 0)$. In the present case of an antiphase domain wall, this should be replaced by $Q = (\frac{4\pi}{5}; 0)$. The spin operator $S(\frac{4\pi}{5}; \pi)$ then creates and annihilates the condensed triplet-excitations [20], so that the intense low-energy peak at $(\frac{4\pi}{5}; \pi)$ is simply the fingerprint of the (quasi)-condensate of triplets, which creates




FIG. 5: Mechanism for alignment of hole pairs.

antiphase Neel-like spin correlations out of the columnar spin-Peierls state. To conclude our discussion of the spin dynamics, we note that $B_y(q; !)$ for the 20×20 cluster also does show a low-energy peak at $(0; !)$ – the tendency towards spin-Peierls instability thus really seems to be quite general[6, 12, 17]. Since the geometry of square cluster is unfavourable for an antiphase domain wall, however, this does not make itself felt as clearly as in the 5×4 cluster.

In summary, it has been shown that a rectangular cluster of the t-J model whose geometry does not explicitly frustrate an antiphase domain wall, shows rather different behavior than the tilted square clusters used conventionally for exact diagonalization. The GS in this cluster shows the clear and unambiguous signatures of a stripe-like domain wall, a comparison of the GS energies shows, that such rectangular clusters are at least as well suited to describe the bulk system, as are the square clusters. Clearly, this does not prove the existence of stripes in the thermodynamic limit – just as a square-shaped cluster will tend to suppress a domain wall, the rectangular one will favour it – but given the previous evidence for stripes found by White and Scalapino[8] on much larger systems, it is quite plausible that the stripes in the rect-

angular cluster have a similar structure.

Stripe-like structures in the rectangular cluster then appear not only in the ground state – rather, there is evidence for excited states where apparently the finite total momentum of the state is carried by a soliton-like propagating domain wall – which is very much reminiscent of the ‘meandering domain wall’ scenario put forward by Zaanen et al.[5]. The data also show the presence of quasistatic columnar singlet order[6, 12] in the system, which seems to be either a prerequisite for or an immediate consequence of the stripe formation. Taken together, the above results suggest the following scenario for doped antiferromagnets: the spontaneous breaking of the point group symmetry of the lattice[17] by formation of bond-singlet order produces ladder-like patterns in the spin background of the system, which then serve as ‘race-tracks’ for hole-pair-like solitons. Condensation of bond-triplets with a condensation amplitude that changes sign across a soliton introduces strong antiferromagnetic correlations with opposite staggered magnetization on the two sides of the soliton. Under these circumstances, it becomes energetically favorable for the solitons on different ladders to form a line, because then the staggered magnetizations on the different ‘ladders’ can align. Solitons which propagate away from the stripe create a track of magnetic frustration (see Figure 5), much as a single hole in an antiferromagnet, which creates an effective potential that aligns the pairs. In this way, one arrives at the picture of a ‘wiggling domain wall’[5], made of loosely bound hole pairs in a background provided by the columnar singlets. An alternative point of view, which also fits the data very well (see especially Figure 2), would be the accumulation of hole pairs along the fault line separating two Neel states with opposite staggered magnetization, as proposed early on by White and Scalapino[21].

-
- [1] J. M. Tranquada, B. J. Sternlieb, J. D. Axe, Y. Nakamura, and S. Uchida, Nature 375.
- [2] D. Poilblanc and T. M. Rice, Phys. Rev. B 39, 9749 (1989); J. Zaanen and O. Gunnarsson, Phys. Rev. B 40, 7391 (1989).
- [3] U. Low et al., Phys. Rev. Lett. 72, 1918 (1994).
- [4] H. T sunetsugu, M. Troyer, and T. M. Rice, Phys. Rev. B 51, 16456 (1995).
- [5] J. Zaanen, O. Y. Osman, and W. van Saarloos, Phys. Rev. B 58, R11 868 (1998); J. Zaanen, O. Y. Osman, H. V. Kuis, Z. Nussinov, and J. Tworzydło, Philos. Mag. B 81, 1485 (2001).
- [6] M. Voja and S. Sachdev, Phys. Rev. Lett. 83, 3916 (1999); K. Park and S. Sachdev, Phys. Rev. B 64, 184510 (2001).
- [7] A. H. Castro-Neto, Phys. Rev. B 64, 104509 (2001).
- [8] S. R. White and D. J. Scalapino, Phys. Rev. Lett. 80, 1272 (1998); 81, 3227 (1998).
- [9] G. B. Martins et al., Phys. Rev. Lett. 84, 5844 (2000); G. B. Martins, C. Gazza, and E. Dagotto, Phys. Rev. B 62, 13926 (2000).
- [10] E. Dagotto, Rev. Mod. Phys. 66, 763 (1994).
- [11] P. Prelovsek and X. Zotos, Phys. Rev. B 47, 5984 (1993).
- [12] O. P. Sushkov, Phys. Rev. B 63, 174429 (2001).
- [13] J. Hasegawa and D. Poilblanc, Phys. Rev. B 40, 9035 (1989); P. W. Leung, Phys. Rev. B 65, 205101 (2002).
- [14] P. W. Robel and R. Eder, Phys. Rev. B 62, 4048 (2000).
- [15] A. L. Chemsyhev, S. R. White, and A. H. Castro Neto Phys. Rev. B 65, 214527 (2002).
- [16] T. Tohyama, P. Horsch, and S. M. Aekawa Phys. Rev. Lett. 74, 980 (1995); R. Eder, Y. Ohta, and S. M. Aekawa Phys. Rev. Lett. 74, 5124 (1995). M. Troyer, H. T sunetsugu, and T. M. Rice Phys. Rev. B 53, 251 (1996).
- [17] N. Read and S. Sachdev, Phys. Rev. Lett. 62, 1694 (1989).
- [18] H. A. Mook, M. Yethiraj, G. Aeppli, T. E. Mason, and T. Amstrong, Phys. Rev. Lett. 70, 3490 (1993).
- [19] S. Sachdev and R. N. Bhatt, Phys. Rev. B 41, 9323 (1990).
- [20] See e.g. Eq(3.11) in S. Gopalan, T. M. Rice and M.

Sigrist, Phys. Rev. B 49, 8901 (1994).

[21] S. R. White and D. J. Scalapino, cond-mat/9610104.