

X-ray diffraction of a disordered charge density wave

Alberto Rosso and Thierry Giamarchi

Université de Genève, DPMC, 24 Quai Ernest Ansermet, CH-1211 Genève 4, Switzerland

We study the X-ray diffraction spectrum produced by a collectively pinned charge density wave (CDW), for which one can expect a Bragg glass phase. The spectrum consists of two asymmetric divergent peaks. We compute the shape of the peaks, and discuss the experimental consequences.

The statics and dynamics of disordered elastic objects govern the physics of a wide range of systems, either periodic, such as vortex flux lines [1] and charge density waves (CDW) [2], or involving propagating interfaces, such as domain walls in magnetic [3] or ferroelectric [4] systems, contact lines of liquid menisci on rough substrates [5] and propagation of cracks in solids [6]. It was recently shown that periodic systems have unique properties, quite different from the ones of the interfaces. If topological defects (i.e. dislocations etc.) in the crystal are excluded, displacements grow only logarithmically [7, 8, 9], instead of the power-law growth as for interfaces. The positional order is only algebraically destroyed [9, 10] leading to divergent Bragg peaks and a nearly perfect crystal state. Quite remarkably, it was shown that for weak disorder this solution is *stable* to the proliferation of topological defects, and thus that a thermodynamically stable phase having both glassy properties and quasi-long range positional order exists [10]. This phase, nicknamed Bragg glass, has prompted many further analytical and experimental studies (see e.g. [11, 12] for reviews and further references). Although its existence can be tested indirectly by the consequences on the phase diagram of vortex flux lines, the most direct proof is to measure the predicted algebraic decay of the positional order. Such a measurement can be done by means of diffraction experiments, using either neutrons or X-rays on the crystal. Neutron diffraction experiments have recently provided unambiguous evidence [13] of the existence of the Bragg glass phase for vortex lattices.

Another periodic system in which one can expect a Bragg glass to occur are charge density waves [2], where the electronic density is spatially modulated. Disorder leads to the pinning of the CDW [14]. In such systems very high resolution X-rays experiments can be performed [15]. The resolution is in principle much higher than the one that can be achieved by neutrons for vortex lattices, consequently CDW systems should be prime candidates to check for the existence of a Bragg glass state. However, compared to the case of vortex lattices the interpretation of the spectrum is much more complicated for two main reasons: (i) the phase of the CDW is the object described by an elastic energy, whereas the X-rays probe the displacements of the atoms in the crystal lattice (essentially a cosine of the phase); (ii) since the impurities substitute some atoms of the crystal, the very presence of the impurities changes the X-ray spectrum.

This generates non-trivial terms of interference between disorder and atomic displacements [15, 16]. It is thus necessary to make a detailed theoretical analysis for the diffraction due to a pinned CDW. The study of the spectrum has been carried out so far either for strong pinning or at high temperatures [15, 16, 17, 18].

In this paper we focus on the low temperature limit where a well formed CDW exists and on weak disorder, for which one expects to be in the Bragg glass regime. We show that the diffraction spectrum consists in two asymmetric peaks. In contrast to previous assumptions [16], we show that the asymmetry is present also in the weak pinning limit. The peaks are power-law divergent, with an anisotropy in shape. This form is consistent with the Bragg glass behavior [10]. The asymmetry is a subdominant power-law too, with an exponent that we determine. We also briefly discuss the role of unscreened Coulomb interaction for the CDW on the diffraction spectrum.

The general expression [19] for the total diffraction intensity in a crystal is given by

$$I(q) = \frac{1}{V} \sum_{i,j} e^{-iq(R_i - R_j)} \left\langle \overline{f_i f_j e^{-iq(u_i - u_j)}} \right\rangle, \quad (1)$$

where u_i is the atom displacement from the equilibrium position $R_j = ja$, with a indicating the lattice constant, f_i the atomic scattering factor and $\overline{\langle \dots \rangle}$ denotes the double average over the disorder and over the thermal fluctuations. As an example let us first consider the case of fixed atoms ($u_i = 0$). We obtain:

$$I(q) = \bar{f}^2 \sum_K \delta(q - K) + \Delta f^2 N_I, \quad (2)$$

where $\Delta f = f_I - f$ is the difference between the impurity I and the host atom scattering factors, $N_I = n_I(1 - n_I)$, with n_I is the impurity concentration and \bar{f} is the average scattering factor. The usual Bragg peaks, in correspondence to the reciprocal lattice vectors K , arise from the first term in (2), the second term is responsible for a background intensity, called Laue scattering, due to the disorder.

In a second stage we take into account displacements of the atoms related to the presence of a CDW. To this purpose, we consider an electron density characterized by a sinusoidal modulation:

$$\rho(x) = \rho_0 \cos(Qx + \phi(x)). \quad (3)$$

ϕ is the phase of the charge density wave and $Q = 2k_F$, where k_F is the Fermi wave vector. The associated Hamiltonian writes:

$$H = \int d^d x \frac{c}{2} (\nabla \phi(x))^2 \pm V_0 \int d^d x \Sigma(x) \rho(x), \quad (4)$$

where d is the dimension of the space. The first term in the Hamiltonian (4) represents the elasticity. The elasticity is in fact anisotropic [20] along the Q -direction:

$$H_{\text{el.}} = \int dx d^{d-1} y \frac{c_1}{2} (\partial_x \phi)^2 + \frac{c_2}{2} (\partial_y \phi)^2, \quad (5)$$

where $x \parallel Q$, and $c_1 \gg c_2$. The compression along x corresponds to an increase of electric charge density and thus pays the price of Coulomb repulsion, while distortions along the remaining $d-1$ directions are much easier. We are led back to (4) by redefining the spatial variables $x' = x/\sqrt{c_1}$ and $y' = y/\sqrt{c_2}$, with $c = (c_1 c_2^{d-1})^{\frac{1}{2}}$. The main effect in the diffraction spectrum is thus to make the *shape* of the peaks anisotropic, but this will not change the overall divergence. The local, but anisotropic, elasticity (5) is valid beyond the distance at which the Coulomb interaction between various parts of the CDW is screened. If this length is very large, or if one want to examine short range regime one should keep the q -dependence in the elastic constants. This leads to a more complicated behavior that we will only briefly discuss here and will be examined in details elsewhere [21]. The second term in the Hamiltonian (4) reflects the effect of the disorder on the electron density. The Gaussian random function $\Sigma(x)$ describes the impurity distribution and is characterized by the correlator $\overline{\Sigma(x)\Sigma(y)} = N_I \delta(x-y)$, V_0 is a positive constant which measures the impurity potential and finally the sign $+$ ($-$) is related to the repulsive (attractive) interaction between the electrons and the local impurity. In the following, we restrict our analysis to the repulsive case, ρ_0 is absorbed in V_0 and we define the disorder strength $D = V_0^2 N_I$.

A density modulation is accompanied by a lattice distortion u given at low temperature by

$$u(x) = \frac{u_0}{Q} \nabla \cos(Qx + \phi(x)). \quad (6)$$

We are interested in the behavior of the scattering intensity $I(q)$ near a Bragg peak ($q \sim K$). Since $|\delta q| = |q - K| \ll K$, we can take the continuum limit $i \rightarrow x$ and we obtain from (1):

$$I(q) = \int_r \left\langle \overline{f_{\frac{r}{2}} f_{-\frac{r}{2}} e^{-i\delta q(u(\frac{r}{2}) - u(-\frac{r}{2}))}} \right\rangle. \quad (7)$$

where $\int_r = \frac{1}{a^d} \int d^d r e^{-i\delta q r}$ and $f_{\frac{r}{2}} = \bar{f} + \Delta f a^{d/2} \Sigma(\frac{r}{2})$. In (7) we have applied the standard decomposition in center of mass R and relative r coordinates ($x = R + \frac{r}{2}$ and $y = R - \frac{r}{2}$). The integration over R has already been

performed because u vary slowly at the scale of the lattice spacing. Assuming that in the elastic approximation displacements remain small ($u_i \ll R_i$), one can expand (7) as powers of Ku_0 . Developing up to the second order we get [17] :

$$\begin{aligned} I(q) &= I_d + I_a + I_{\text{tripl.}}, \text{ with} \\ I_d &= \bar{f}^2 q^2 \int_r \overline{u(\frac{r}{2}) u(-\frac{r}{2})}, \\ I_a &= -iq \Delta f a^{d/2} \bar{f} \int_r \left\langle \overline{\Sigma(-\frac{r}{2}) u(\frac{r}{2}) - \Sigma(\frac{r}{2}) u(-\frac{r}{2})} \right\rangle, \\ I_{\text{tripl.}} &= -iq \Delta f^2 a^d \int_r \left\langle \overline{\Sigma(-\frac{r}{2}) \Sigma(\frac{r}{2}) (u(\frac{r}{2}) - u(-\frac{r}{2}))} \right\rangle. \end{aligned} \quad (8)$$

While the contribution I_d represents the intensity due to the atomic displacements alone, the contributions I_a and $I_{\text{tripl.}}$ are generated by the coupling between the disorder and the displacement. The presence of a CDW is signaled by the formation, around each Bragg peak, of two satellites at reciprocal vectors $K \pm Q$. In absence of disorder ($D = 0$ and $\phi \sim \text{const.}$) the displacement term has the form $I_d = f^2 q^2 u_0^2 \sum_K \delta(q + K \pm Q)$ and the other terms are vanishing: in this case the two satellites have the same intensity and the broadening is absent. To interpret the experimental findings [15, 16, 18], in particular to explain the measured strong asymmetry [25] between the peaks at $K + Q$ and at $K - Q$, we need to account for the effect of impurities.

In the literature the term I_a was evaluated by means of models [16, 17, 18] which describe the pinning by imposing a constant value ϕ_0 to the phase in (3) in proximity of each impurity, and $I_{\text{tripl.}}$ was conjectured to be negligible [17, 22]. In that approach, the observed satellite asymmetry is seen as a clear sign of the strong disorder; in fact, ϕ_0 is not constant and, for sufficiently large domains, one should have [16] $I_a \propto \overline{\cos(\phi_0)} \sim 0$. To go beyond this phenomenological approach and also deal with the weak disorder limit, in which one expects the Bragg glass, we use a Gaussian variational approach [10, 23]. We first perform the average over the disorder using the standard replica techniques. The replicated Hamiltonian corresponding to (4) is

$$H_{\text{eff.}} = \int d^d r \sum_a \frac{c}{2} (\nabla \phi_a)^2 - \frac{D}{T} \sum_{a,b} \cos(\phi_a(r) - \phi_b(r)), \quad (9)$$

where T is the temperature and the sum over the n replica has to be considered in the limit $n \rightarrow 0$. We stress that, moving from the Hamiltonian (4) to its replicated version we also need to change the correlation functions containing explicitly the disorder: we have, for example, $\left\langle \overline{\Sigma(-\frac{r}{2}) u(\frac{r}{2})} \right\rangle \rightarrow -\frac{D}{TV_0} \sum_a \langle \rho_a(-\frac{r}{2}) u_1(\frac{r}{2}) \rangle_{\text{eff.}}$. After




FIG. 1: Intensities of the different contributions to satellite peaks. The more divergent term, I_d , is symmetric. I_a and $I_{\text{tripl.}}$ are antisymmetric, with $I_a \gg I_{\text{tripl.}}$.

some manipulations and using (3), we obtain:

$$I_d = \bar{f}^2 q^2 u_0^2 \int_r [e^{-iQr} + c.c.] C_d(r), \quad (10)$$

$$I_a = -\bar{f} \Delta f q u_0 \sqrt{N_I a^d D} \int_r [e^{-iQr} - c.c.] C_a(r),$$

where $C_d(r) = \langle e^{i(\phi_1(\frac{r}{2}) - \phi_1(-\frac{r}{2}))} \rangle_{\text{eff.}}$ and $C_a(r) = \frac{1}{T^n} \sum_{a,b}^n \langle e^{i(\phi_a(\frac{r}{2}) - \phi_b(-\frac{r}{2}))} \rangle_{\text{eff.}}$ are the positional correlation functions controlling the behavior of each contribution. We notice that the intensity of the peaks at $q = Q + K$ and $q = K - Q$ is symmetric, as in the case of a pure system, for the displacement term I_d , but it is antisymmetric for I_a . The sum of these two terms leads to an asymmetry of the peaks. Fig. 1 show the behavior of the different contributions.

Following the method used in [10] for flux lines in presence of weak disorder, we can calculate the various terms in (10). We look for the best trial Gaussian Hamiltonian $H_0 = \int_q G_{ab}(q) \phi_a(q) \phi_b(-q)$ in replica space, which approximates (9). Defining

$$B_{ab}(r) = \langle (\phi_a(r) - \phi_b(0))^2 \rangle_0 \quad (11)$$

$$= 2T \int_q [\tilde{G}(q) - G_{ab}(q) \cos qr],$$

where \tilde{G} is the diagonal element of G_{ab} , and using the Gaussian approximation, the positional correlation functions become $C_a(r) = \frac{1}{nT} \sum_{a,b}^n e^{-\frac{B_{ab}(r)}{2}}$ and $C_d(r) = e^{-\frac{\tilde{B}(r)}{2}}$, where \tilde{B} is the diagonal element of B_{ab} . Two general classes of solutions exist for this problem: while the first class preserves the permutation symmetry of the replica (RS), the second class (RSB) breaks the replica symmetry. It has been shown [10] that the stable solution for $d > 2$ corresponds to the RSB class, while the RS solution remains valid at short distance. C_d is similar to the correlation calculated for flux lines [10] and will

be discussed later. To evaluate the contribution of the interference between disorder and displacement we factorize the antisymmetric term $C_a(r) = \chi(r) C_d(r)$. We first consider the RS approximation:

$$\chi(r) = \frac{1}{T} \left[1 - e^{-T \int_q G_c(q) \cos qr} \right], \quad (12)$$

where $G_c = \frac{1}{cq^2}$ is the connected part of G_{ab} . In $d = 3$ we estimate

$$C_a^{RS}(r) \sim \frac{2\pi^2}{cr} e^{-\frac{\tilde{B}(r)}{2}}. \quad (13)$$

The triplet term can be evaluated in an analogous way, but it gives non-zero contributions only considering higher order harmonic terms in the electron density. Equation (3) becomes: $\rho(x) = \rho_0 \cos(n(Qx + \phi(x)))$, with $n = 1, 2$. As we have already found for I_a , we get an antisymmetric term with a prefactor $\propto \Delta f^2 q u_0 N_i a^d D$ and a correlation $C_{\text{tripl.}} = \frac{1}{2nT^2} \sum_{a,b,c}^n \langle (e^{-i(\phi_c(\frac{r}{2}) - 2\phi_a(\frac{r}{2}) + \phi_b(-\frac{r}{2}))}) \rangle_{\text{eff.}}$. In $d = 3$ and at low temperature we finally obtain

$$C_{\text{tripl.}}^{RS}(r) \sim \frac{2\pi}{c^2 ar} e^{-\frac{\tilde{B}}{2}}. \quad (14)$$

It is interesting to evaluate, at this stage, the relative weight of the two antisymmetric terms in a satellite peak. We introduce the Fukuyama-Lee length (or Larkin-Ovchinnikov length) [14, 24] $R_a = (c^2/D)^{1/4-d}$ (for $d = 3$ $R_a = c^2/D$) such that ϕ varies on scale given by the length R_a . The ratio of the two intensity peaks is:

$$\frac{I_{\text{tripl.}}}{I_a} = -\frac{\Delta f}{\pi \bar{f}} \sqrt{N_I} \sqrt{\frac{a}{R_a}}. \quad (15)$$

For weak disorder $R_a \gg a$ it follows that $I_a \gg I_{\text{tripl.}}$ and we thus need only to consider I_a and I_d .

Since for $d = 3$ the RS solution is unstable, to obtain the correct physics one has to look for the RSB method. Within this scheme [10], the off diagonal elements of $G_{ab}(q)$ are parameterized by $G(q, v)$ where $0 < v < 1$ and the solution is characterized by a variational break-point v_c . The form of the symmetric part is given in [10]:

$$C_d(r) \sim e^{-\frac{\phi_T^2}{2}} \left(\frac{l}{r} \right)^\eta \quad (16)$$

where $\phi_T^2 \simeq \frac{2T}{\pi c a}$ measures the strength of thermal fluctuations, and $\eta \sim 1$ is the Bragg glass exponent in $d = 3$. At low temperature one has $l \sim R_a$. The algebraic behavior of (16) is controlled by small v ($v < v_c$). Values of v above the breaking point ($v > v_c$) give the small distance contribution. Finally one finds $v_c = \frac{1}{8} \phi_T^2 \frac{a}{l}$.

To fully characterize the spectrum it still remains to evaluate $\chi(r)$ in the RSB scenario :

$$\chi(r) = \frac{1}{T} \left[1 - \int_0^1 dv e^{-T \int_q (\tilde{G}(q) - G(q, v)) \cos qr} \right]. \quad (17)$$




FIG. 2: Ratio between the RSB and RS solutions for $\chi(r)$. At large distance this ratio tends towards a constant value b , where $b \sim 0.018$. This means that the RSB solution affects $\chi(r)$ only by a multiplicative factor.

Restricting to the case $d = 3$, we write:

$$\tilde{G}(q) - G(q, v) = \frac{1}{c} \left[\frac{1}{q^2 + l^{-2}} + \frac{2}{l^2} \int_{v/v_c}^1 dt \frac{1}{(q^2 + (\frac{t}{l})^2)^2} \right]. \quad (18)$$

By integrating (18) over q and with some manipulations, (17) becomes:

$$\chi(r) = \frac{v_c}{T} \left[1 - \int_0^1 dz \exp \left(-8\pi^3 \int_z^1 \frac{dt}{t} e^{-rt/l} \right) \right] \quad (19)$$

The low temperature behavior ($l \sim R_a$) of this term is sketched in Fig. 2. As for the Replica Symmetry case, we have $C_a(r) \propto \frac{1}{r} e^{-\frac{\tilde{B}(r)}{2}}$. We can now compare the two terms:

$$I_d(K+Q) = \bar{f}^2 K^2 u_0^2 \int_r \left(\frac{R_a}{r} \right)^\eta \quad (20)$$

$$I_a(K+Q) = -2\pi^2 \bar{f} \Delta f \sqrt{N_I} \sqrt{\frac{a}{R_a}} K u_0 \int_r \left(\frac{R_a}{r} \right)^\eta \frac{ba}{r}.$$

After executing the d -dimensional Fourier Transforms, we conclude that both terms are divergent: in particular, $I_d \propto \frac{1}{q^{d-\eta}}$ and $I_a \propto \frac{1}{q^{d-\eta-1}}$. This effect, shown in Fig. 1, is a clear sign of a quasi-long range positional ordered phase. We have found that the peak at $K+Q$ is smaller than the $K-Q$ one, as the potential between the impurity and CDW is repulsive (we would have the opposite asymmetry in case of an attractive potential). We observe that for an ideal infinite resolution experiment, the symmetric term would be dominant, since $C_d(r)$ decays to zero less rapidly than $C_a(r)$. However, if the divergence in (20) is cut by the finite resolution of the experiment both terms should be taken into account because I_d is quadratic in the small parameter Ku_0 whereas I_a is only linear.

The powerlaw lineshape is obtained for a short range elasticity. If the Coulomb interaction is unscreened, as

might be the case in fully gapped systems such as the blue bronzes, the dispersion of c_1 should be kept in (5). In that case $c_1(q) \sim q_x^2/q^2$, which leads to peaks diverging even faster than (20) [21].

On the experimental side few detailed diffraction spectra are available at the moment. One case is doped blue bronzes where the lineshape corresponding to the CDW has been obtained after subtraction of a Friedel oscillation contribution [15]. The observed asymmetry of the peaks would be compatible with both strong and weak pinning. However given the short correlation length extracted from the data, this particular experiment is most likely still in the strong pinning regime. It would thus be highly desirable to have more detailed analysis of the lineshapes either in this compound, for different impurity concentrations, or in less disordered systems, where one can expect a Bragg glass behavior.

We thank J.-P. Poujet and S. Ravy for stimulating discussions.

-
- [1] G. Blatter, M. V. Feigel'man, V. B. Geshkenbein, A. I. Larkin, and V. M. Vinokur, Rev. Mod. Phys. **66**, 1125 (1994).
 - [2] G. Grüner, Rev. Mod. Phys. **60**, 1129 (1988).
 - [3] S. Lemerle, J. Ferré, C. Chappert, V. Mathet, T. Giamarchi, and P. Le Doussal, Phys. Rev. Lett. **80**, 849 (1998).
 - [4] T. Tybell, P. Paruch, T. Giamarchi, and J. M. Triscone, Phys. Rev. Lett. **89**, 097601 (2002).
 - [5] S. Moulinet, C. Guthmann, and E. Rolley, Eur. Phys. J. E **8**, 437 (2002).
 - [6] H. Gao and J. R. Rice, J. Appl. Mech. **56**, 828 (1989).
 - [7] T. Nattermann, Phys. Rev. Lett. **64**, 2454 (1990).
 - [8] S. E. Korshunov, Phys. Rev. B **48**, 3969 (1993).
 - [9] T. Giamarchi and P. Le Doussal, Phys. Rev. Lett. **72**, 1530 (1994).
 - [10] T. Giamarchi and P. Le Doussal, Phys. Rev. B **52**, 1242 (1995).
 - [11] T. Nattermann and S. Scheidl, Adv. Phys. **49**, 607 (2000).
 - [12] T. Giamarchi and S. Bhattacharya, in *High Magnetic Fields: Applications in Condensed Matter Physics and Spectroscopy*, edited by C. Berthier et al. (Springer-Verlag, Berlin, 2002), p. 314, cond-mat/0111052.
 - [13] T. Klein et al., Nature **413**, 404 (2001).
 - [14] H. Fukuyama and P. A. Lee, Phys. Rev. B **17**, 535 (1978).
 - [15] S. Rouzière, S. Ravy, J.-P. Poujet, and S. Brazovskii, Phys. Rev. B **62**, R16231 (2000).
 - [16] S. Ravy and J.-P. Poujet, J. de Phys. IV **3**, 109 (1993).
 - [17] S. Ravy, J.-P. Poujet, and R. Comès, J. de Phys. I **2**, 1173 (1992).
 - [18] S. Brazovskii, J.-P. Poujet, S. Ravy, and S. Rouzière, Phys. Rev. B **55**, 3426 (1997).
 - [19] A. Guiner, *X-Ray diffraction, imperfect Crystals and Amorphous Bodies* (W. H. Freeman and Company, San Francisco, 1969).
 - [20] D. Feinberg and J. Friedel, in *Low-Dimensional Electronic Properties of Molybdenum Bronzes and Oxides*,

- edited by C. Schlenker (Kluwer Academic Publishers, Netherlands, 1989), p. 407.
- [21] A. Rosso and T. Giamarchi (2003), in preparation.
 - [22] J. M. Cowley, Surf. Sci. **298**, 336 (1993).
 - [23] M. Mézard and G. Parisi, J. de Phys. I **1**, 809 (1991).
 - [24] A. I. Larkin and Y. N. Ovchinnikov, J. Low Temp. Phys **34**, 409 (1979).
 - [25] The asymmetry of the two peaks can be very strong: it has been observed that the intensity of the lowest peak can even be smaller than the Laue scattering intensity. In this case one talks of white line in the spectrum.