

# A quantum generalization of the thermal viscous friction law

P. Shiktorov, E. Starikov, V. Gružinskis

Semiconductor Physics Institute, A. Goštauto 11, 2600 Vilnius, Lithuania

L. Reggiani

INFM - National Nanotechnology Laboratory, Dipartimento di Ingegneria dell'Innovazione, Università di Lecce, Via Arnesano s/n, 73100 Lecce, Italy

(March 22, 2022)

On the basis of the equivalence of the energy balance description at micro- and macro-level we propose a quantum generalization of the viscous friction law for a macroscopic Langevin equation describing thermal fluctuations without the zero point contribution. This equation recovers the classical case in the limit of  $\hbar \rightarrow 0$ . In particular it satisfies the quantum regression theorem and resolves several anomalies appearing in the quantum extension of the fluctuation dissipation theorem.

PACS numbers: 03.65.Ca, 05.30.-d, 05.40.+j

The classical Langevin equation for the damped harmonic oscillator with eigenfrequency  $\omega_s$  in the presence of friction and of a stochastic Langevin force  $f_C(t)$  in the time domain writes:

$$\frac{d^2}{dt^2}x + \gamma \frac{d}{dt}x + \omega_s^2 x = f_C(t) \quad (1)$$

Here  $\gamma$  is the viscous friction coefficient and the correlation function of the thermal Langevin force is given by  $\langle f_C(0)f_C(t) \rangle = 2kT\gamma\delta(t)$ . Equation (1) forms a basis for the fluctuation dissipation theorem (FDT) in its classical form:

$$S_{xx}^C(\omega) = \frac{2kT}{\omega} \text{Im}\{\alpha_x^C(\omega)\} \quad (2)$$

with  $S_{xx}^C$  the two-side spectral density of the displacement fluctuations in the frequency range  $[-\infty, \infty]$  and  $\alpha_x^C(\omega)$  the classical susceptibility (i.e. the Fourier transform of the response function) that for the damped harmonic oscillator writes:

$$\alpha_x^C(\omega) = \frac{1}{\omega_s^2 - \omega^2 - i\gamma\omega} \quad (3)$$

According to a recent comment of van Kampen<sup>1</sup>: *the shortcoming of the above Langevin scheme is that it cannot be extended to quantum mechanics; attempts have been made to simple generalize Eq. (1) so as to turn it into an equation of operators, but that did not work.*<sup>2</sup> Indeed, one way to perform such a generalization is to introduced a relaxation already at the level of the Heisenberg equations of motion for operators, the so called *quantum Langevin equations*.<sup>3,4</sup> However, as mentioned

by van Kampen, such an approach is internally contradictory since formally it implies that a set of quantum-mechanical operators and their commutation rules which describe the physical system are relaxing. This would be a severe violation of the laws of quantum mechanics since commutation relations must be valid at all times.<sup>4</sup>

To overcome this contradiction, a second way of proceeding is to introduce a relaxation for observables only. Accordingly, relaxation appears just in the macroscopic equations of motion for the observables, such as  $x = \text{Tr}\{\hat{\rho}\hat{x}\}$ , that are obtained after averaging over the statistical operator  $\hat{\rho}$ .<sup>4,5</sup> This second approach is free from the above contradiction since the quantum corrections to the Langevin scheme are formulated only at the macrolevel on the basis of the quantum FDT (QFDT) given usually by the well known Callen and Welton<sup>6</sup> expression. In full analogy with the classical case, here the friction remains viscous and the modifications concern with the Langevin force spectrum only. However, some contradictions appear in this scheme too. For example, when thermal fluctuations are considered this second approach enters in conflict with the quantum regression theorem (QRT). Usually such a conflict is announced in the form *"there is no quantum regression theorem"*<sup>7,8</sup>, despite of the fact that the QRT was derived by Lax<sup>9</sup> independently from the QFDT.

The aim of this letter is to remove these contradictions of the second approach by introducing a quantum viscous force law which generalizes the relaxation of observables given by Eq. (1) to the quantum case. To this purpose: (i) the classical viscous friction law is replaced by the quantum analog as

$$\gamma \frac{d}{dt}x \rightarrow \gamma \frac{2}{\beta} \sin\left(\frac{\beta}{2} \frac{d}{dt}\right)x \quad (4)$$

and (ii) the classical Langevin force correlator by the quantum analog:

$$\langle f_Q(0)f_Q(t) \rangle = \frac{2\hbar\gamma}{\pi\beta} \int_0^\infty \exp\left(-\frac{\beta\omega}{2}\right) \cos(\omega t) d\omega \quad (5)$$

where  $1/\beta = kT/\hbar$ .

The above quantum generalization of the Langevin scheme satisfies the QFDT in the form originally proposed by Nyquist<sup>10</sup> without the contribution of the zero-point energy of the thermal bath radiation:

$$S_{xx}^Q(\omega) = 2\hbar \text{sgn}\{\omega\} \bar{N}_T(\omega) \text{Im}\{\alpha_x^Q(\omega)\} \quad (6)$$

with the quantum susceptibility  $\alpha_x^Q(\omega)$ :

$$\alpha_x^Q(\omega) = \frac{1}{\omega_s^2 - \omega^2 - i\frac{\gamma}{\beta} \sinh(\frac{\omega\beta}{2})} \quad (7)$$

Here  $\bar{N}_T(\omega) = [\exp(\beta|\omega|) - 1]^{-1}$  is the average number of thermal photons with frequency  $\omega$ .

Below, we shall demonstrate that the above extended approach is consistent with the principle of detailed energy balance, while the conventional approach based on the classical viscous force and Callen-Welton expression<sup>6</sup> of the QFDT violate this principle.

*Micro- and macro-level of detailed energy balance description.* We shall consider a sufficiently large isolated system, with Hamiltonian  $\hat{H} = \hat{H}_S + \hat{H}_T + \hat{V}$ , which can be decomposed into two interacting subsystems. The first one is described by  $\hat{H}_S$  and corresponds to the physical system under test. The second subsystem is described by  $\hat{H}_T$  and represents the surrounding world. The interaction between these subsystems is described by the Hamiltonian  $\hat{V} = -\hat{x}\hat{f}$ , where  $\hat{x}$  and  $\hat{f}$  are operators representing dynamical variables pertaining to the S- and T-subsystems, respectively.

By using the standard procedure<sup>4</sup> to go from the density matrix of the whole system  $\hat{\rho}$  to the reduced density matrixes of subsystems  $\hat{\rho}_S = \text{Tr}_T\{\hat{\rho}\}$  and  $\hat{\rho}_T = \text{Tr}_S\{\hat{\rho}\}$  in the framework of the energy representation of  $\hat{\rho}_i = \hat{\rho}_i(\hat{H}_i)$  one obtains the following equation for the time evolution of the average energy  $\langle \hat{H}_i \rangle$  in the  $i$ -th subsystem ( $i = S, T$ ):

$$\begin{aligned} \frac{d}{dt} \left[ \langle \hat{H}_T \rangle \right] &= \frac{\pi}{\hbar} \sum_{m,n}^S \sum_{M,N}^T |x_{mn}|^2 |f_{MN}|^2 \\ &\times \left[ \frac{\omega_{mn}^S}{\omega_{MN}^T} \right] (\rho_m^S \rho_M^T - \rho_n^S \rho_N^T) \delta(\omega_{mn}^S + \omega_{MN}^T) \end{aligned} \quad (8)$$

where  $\rho_m^i$  is the probability to find the  $i$ -th subsystem in the state with eigenenergy  $E_m^i$ ,  $\omega_{mn}^i = (E_m^i - E_n^i)/\hbar$  the frequency of permitted transitions,  $x_{mn}$  and  $f_{MN}$  the matrix elements of the  $\hat{x}$  and  $\hat{f}$  operators, respectively. When  $\langle \hat{H}_i \rangle = \text{const}$ , from Eq. (8) one directly obtains the condition of microscopic detailed energy balance<sup>11</sup>

$$\rho_m^S \rho_M^T = \rho_n^S \rho_N^T \quad (9)$$

To formulate the conditions of energy balance at the macroscopic level of description, we replace the factor  $\delta(\omega_{mn}^S + \omega_{MN}^T)$  by  $\int \delta(\omega_{mn}^S - \omega) \delta(\omega_{MN}^T + \omega) d\omega$  and by using the definition of spectral density given by:

$$J_{ii}(\pm\omega) = 2\pi \sum_{m,n} \rho_n |i_{mn}|^2 \delta(\omega_{mn} \mp \omega) \quad (10)$$

which corresponds to the quantum correlation function  $\text{Tr}\{\hat{\rho}_i(\pm\tau)\hat{i}(0)\}$  of the dynamical operators  $\hat{i} = \hat{x}, \hat{f}$ , we

rewrite Eq. (8) as:

$$\begin{aligned} \frac{d}{dt} \langle \hat{H}_T \rangle &= -\frac{d}{dt} \langle \hat{H}_S \rangle = \\ &= \frac{1}{4\pi\hbar} \int_{-\infty}^{\infty} \omega [J_{xx}(-\omega)J_{ff}(\omega) - J_{xx}(\omega)J_{ff}(-\omega)] d\omega \end{aligned} \quad (11)$$

From Eq. (11) we obtain the condition of macroscopic detailed energy balance (MaDEB) as:

$$J_{xx}(-\omega)J_{ff}(\omega) = J_{xx}(\omega)J_{ff}(-\omega) \quad (12)$$

which requires to be fulfilled for any value of the current frequency  $\omega$ . We notice that the condition given by Eq. (12) is not the only form in which the MaDEB can be expressed. By using the expression for the imaginary part of the macroscopic susceptibility<sup>12,13</sup>:

$$\text{Im}\{\alpha_i(\omega)\} = \frac{1}{2\hbar} [J_{ii}(\omega) - J_{ii}(-\omega)] \quad (13)$$

which describes the dissipative part of the response of one subsystem to the action of the other one (for example, if  $i = x$  is the response then the action is given by  $f = \text{Tr}\{\hat{\rho}_T \hat{f}\}$  and *vice versa*) the MaDEB conditions can be rewritten in an equivalent form as:

$$Y_x^\mu(\omega) \text{Im}\{\alpha_f(\omega)\} = Y_f^\mu(\omega) \text{Im}\{\alpha_x(\omega)\} \quad (14)$$

where, in general, the weighted symmetric spectral density  $Y_i^\mu(\omega)$  is given by:

$$Y_i^\mu(\omega) = (1 - \mu)J_{ii}(-|\omega|) + \mu J_{ii}(|\omega|) \quad (15)$$

and the weighting factor  $\mu$  can be treated as an arbitrary parameter. Note, that to derive Eq. (14) one need to preserve the antisymmetric property of the term in the square brackets of Eq. (11) under all the transformations. Since by definition  $\text{Im}\{\alpha_i(\omega)\}$  is an antisymmetric function of frequency, to keep the antisymmetry of the whole expression,  $Y_i^\mu(\omega)$  must be treated as a symmetric function, i.e. as a function of the absolute value of frequency.

In the following we shall consider the formulation of MaDEB conditions by using the ratios of some characteristics of only one of the two subsystems. In such a representation, Eq. (12) takes the form:

$$\frac{J_{xx}(-\omega)}{J_{xx}(\omega)} = \frac{J_{ff}(-\omega)}{J_{ff}(\omega)} \equiv p(\omega) \quad \text{or} \quad J_{ii}(-\omega) = p(\omega)J_{ii}(\omega) \quad (16)$$

where  $p(\omega)$  is a factor common to both subsystems satisfying the relation  $p(-\omega) = p^{-1}(\omega)$ . The MaDEB condition given by Eq. (14) can be represented in the analogous forms:

$$Y_i^\mu(\omega) = g^\mu(\omega) \text{Im}\{\alpha_i(\omega)\} \quad (17)$$

where

$$g^\mu(\omega) = 2\hbar \text{sgn}\{\omega\} \left[ \frac{p(|\omega|)}{1 - p(|\omega|)} + \mu \right] \quad (18)$$

Here,  $g^\mu(\omega)$  depends on  $\mu$  and is determined by the frequency dependence of  $p(\omega)$  at  $\omega > 0$  only.

When the physical system interacts with the radiation bath described by a certain, not obligatory thermal, distribution of photon numbers in the radiation modes, the bath spectral densities  $J_{ff}(\pm\omega)$  can be represented as<sup>14</sup>:

$$J_{ff}(\pm|\omega|) = 2\pi \frac{G(\omega)\gamma^2(\omega)\hbar}{\omega} \left[ \overline{N}(\omega) + \frac{1}{2} \pm \frac{1}{2} \right] \quad (19)$$

where  $G(\omega)$  is the density of radiation modes in the bath and  $\gamma^2(\omega)$  the electro-dipole matrix element square. Substitution of Eq. (19) into the MaDEB condition given by Eq. (16) allows us to rewrite the latter in terms of the average number of photons in the radiation mode with frequency  $\omega$ :

$$\overline{N}(\omega) \equiv \text{Tr}\{\hat{\rho}_T \hat{n}_\omega\} = \frac{p(|\omega|)}{1 - p(|\omega|)} \quad (20)$$

Accordingly, the MaDEB condition given by Eq. (17) takes the form:

$$Y_x^\mu(\omega) = 2\hbar \text{sgn}\{\omega\} [\overline{N}(\omega) + \mu] \text{Im}\{\alpha_x(\omega)\} \quad (21)$$

By using Eq. (21), let us give an interpretation to the energy balance at the macro-level. As follows from Eq. (14) the energy balance description at the macro-level is formulated only in the framework of the notion of a classical absorption (i.e. the energy dissipation by one subsystem from another) which is described by  $\text{Im}\{\alpha_i(\omega)\}$ . With respect to each interacting subsystem such an energy transfer occurs in one direction, from outside to inside. The opposite process, which returns the energy back, i.e. the energy emitted by the subsystem is absent. This representation is due to the definition of  $\alpha_i(\omega)$  as the response of the system to a classical external force which, by definition, does not take into account the process of spontaneous emission. Therefore, with respect to a single subsystem, one side of Eq. (14) must be treated as the absorbed (dissipated) power and the other side as the emitted (returned) power. Indeed, a direct substitution of Eq. (19) into Eq. (14) gives us again Eq. (21). Here, the right-hand side, which contains  $\text{Im}\{\alpha_x(\omega)\}$ , describes the power gained from the radiation bath and dissipated in the system. The left-hand side, which is given identically by  $Y_x^\mu(\omega)$ , describes the power returned by the system to the bath. However, as follows from Eq. (21), at the MaDEB level the state of the bath radiation which is proportional to  $(\overline{N}(\omega) + \mu)$  and the spectrum of fluctuations  $Y_x^\mu(\omega)$  are formally uncertain since  $\mu$  can take arbitrary values.

*Equivalence between MiDEB and MaDEB.* As follows from the above consideration, micro- and macro-levels of the energy balance description can not be treated as entirely equivalent. Indeed, the former is formulated in

terms of statistical operators  $\hat{\rho}_i$  (see Eq. (9)), which describe the interacting subsystem microstates only. By contrast, the latter is based on characteristics which are by definition invariant with respect to these microstates since all the MaDEB conditions are formulated for the quantities statistically averaged over  $\hat{\rho}_i$ . Moreover, the MaDEB conditions can include the extra arbitrary parameter  $\mu$  which does not directly follow from MiDEB.

It is easy to see, that the energy balance descriptions based on Eqs. (8) and (11) will be equivalent if the following relation is satisfied at all the frequencies of permitted transitions between the subsystems:

$$\frac{\rho_m^S}{\rho_n^S} = \frac{\rho_N^T}{\rho_M^T} = p(\omega) \Big|_{\omega=\omega_{mn}=\omega_{NM}} \quad (22)$$

At thermal equilibrium, when  $\hat{\rho}_i \sim \exp(-\hat{H}_i/kT)$ , the common factor  $p(\omega) = \exp(-\hbar\omega/kT)$  becomes the universal function of the temperature and frequency only, so that the MaDEB condition given by Eq. (21) at  $\mu = 0$  and  $1/2$  coincides with the Nyquist<sup>10</sup> and Callen-Welton<sup>6</sup> versions of the QFDT, respectively. *This fact allow us to give to the QFDT an alternative physical interpretation with respect to the conventional one as a relation which describes the MaDEB between two interacting physical systems under thermal equilibrium.*

The explicit presence of the uncertainty introduced by  $\mu \neq 0$  in the MaDEB given by Eqs. (17) and (21) leads to a disagreement between the micro- and the macro-level of description since, in essence, it claims that the statistical and photon-number operators,  $\hat{\rho}_T$  and  $\hat{n}$ , respectively, are not sufficient to determine the average number of photons in the mode, i.e.  $\overline{N} \neq \text{Tr}\{\hat{\rho}_T \hat{n}\}$ . In accordance with the MiDEB and MaDEB equivalence condition given by Eq. (22),  $\mu$  can always be neglected by renormalizing the average number of photons in the bath  $N'(\omega) = \overline{N}(\omega) + \mu$ . Such a renormalization will change only the value of  $\hat{\rho}_S$ , i.e. a microstate, by transforming  $Y_x^\mu(\omega)$  to  $J_{xx}(-|\omega|)$ . In this case, only the photon part,  $\hbar\omega\overline{N}(\omega)$ , of the full energy of the field is involved in the dissipation process described by  $\text{Im}\{\alpha_x(\omega)\}$ , and the process of returning back the energy is accomplished by the spontaneous emission characterized by the spectral density  $J_{xx}(-|\omega|)$ . Thus, when  $\mu = 0$ , Eq. (21) describes the energy exchange between the two subsystems consistently with the microscopic picture.

As follows from Eq. (22), in the frequency regions where  $p(\omega) = 1$  the energy transitions are under saturation, that is  $\rho_m = \rho_n$  at  $\omega = \omega_{mn}$ . The saturated transitions can only appear jointly in both the interacting subsystems. In the opposite case, the MiDEB conditions are violated. By using Eq. (13) and Eq. (16), the imaginary part of the susceptibilities can be represented as:

$$\text{Im}\{\alpha_i(\omega)\} = \frac{1}{2\hbar} [1 - p(\omega)] J_{ii}(\omega) \quad (23)$$

As follows from Eq. (23), at the frequencies of the saturated transitions the macroscopic response of both the

subsystems must not contain the dissipative component, since  $Im\{\alpha_i(\omega)\} = 0$  at  $p(\omega) = 1$ .

If one of the subsystems is the radiation bath, then, as follows from Eq. (20), the dissipativeless interaction occurs at all the frequencies where  $\overline{N}(\omega) \rightarrow \infty$ . Note, that the state of the radiation in these modes is similar to the classical state with exactly determined phase<sup>14</sup>. Under thermal equilibrium the frequencies of the saturated transitions are determined by the condition  $exp(-\beta\omega) = 1$  and they correspond to the so-called Matsubara frequencies  $\omega = \Omega_k \equiv i2\pi\beta^{-1}k$  with  $k = 0, \pm 1, \pm 2, \dots$  placed at  $k \neq 0$  in the imaginary axis. The classical viscous friction law introduced usually to describe the macroscopic dissipation satisfies the MiDEB conditions of the dissipativeless interaction, however, on the real frequency axis only. Indeed on the real axis  $\overline{N}(\omega) \rightarrow \infty$  only at  $\omega = 0$  where the viscous friction law always leads to  $Im\{\alpha_x(\omega)\} = 0$ . However, this law fails by extending these MiDEB requirements to the whole complex frequency plane since it formally violates the energy balance at the Matsubara frequencies with  $k \neq 0$  where  $Im\{\alpha_x(\Omega_k)\} \neq 0$ . Such a violation is just the source of the QRT-QFDT conflict<sup>7,8</sup> and of unremovable divergences of the correlation functions of fluctuations when  $T \rightarrow 0$ .<sup>3,5</sup>

*The Langevin scheme.* In the framework of linear response theory, the spectral densities of fluctuations of the observable and the external force by which they are initiated are interrelated by the simple relation:

$$Y_x^\mu(\omega) = |\alpha_x(\omega)|^2 Y_f^\mu(\omega) \quad (24)$$

This allows us to rewrite MaDEB condition given by Eq. (17) in the form:

$$Y_f^\mu(\omega) = -g^\mu(\omega) Im\{\alpha_x^{-1}(\omega)\} \quad (25)$$

where  $Im\{\alpha_x^{-1}(\omega)\} = -Im\{\alpha_i(\omega)\}/|\alpha_i(\omega)|^2$  describes the relaxation law of fluctuations of the observable. For instance,  $Im\{\alpha_x^{-1}(\omega)\} = -\gamma\omega$  in the case of the classical viscous friction. In the framework of the Langevin approach the quantity  $Y_f^\mu(\omega)$  determines the spectral density of the Langevin force which initiates fluctuations of the observable described by the spectral density  $Y_x^\mu(\omega)$ .

To match the Langevin scheme with MiDEB it is necessary:

(i) To exclude all the extra contributions caused by the free parameter  $\mu$ . It means that the spectrum of fluctuations of the observable must coincide with the spontaneous emission spectrum of the system under test. This MiDEB requirement agrees with the opinion often meet in the literature<sup>15,16</sup> that zero-point fluctuations which corresponds to MaDEB condition with  $\mu = 1/2$  cannot be directly detected.

(ii) To satisfy the dissipativeless interaction requirement, the condition  $Im\{\alpha_x(\omega)\} = 0$  must hold in all the points of the complex frequency plane where  $p(\omega) = 1$ .

The former requirement is satisfied in a trivial way by assuming  $\mu = 0$ . One of the possible ways to satisfy the latter one is to modify the viscous friction law.

For this sake, by taking into account that by definition  $Im\{\alpha_x^{-1}(\omega)\}$  and  $Y_f^0(\omega)$  are odd and even functions of frequency, respectively, let us rewrite Eq. (25), which determines the spectral density of the Langevin force, in the form:

$$Y_f^0(\omega) = \left[ 2\hbar \frac{\gamma}{\beta} e^{-\frac{\beta|\omega|}{2}} \right] \times \left[ \frac{\beta}{\gamma} \frac{Im\{\alpha_x^{-1}(\omega)\}}{(e^{-\frac{\beta\omega}{2}} - e^{\frac{\beta\omega}{2}})} \right] \quad (26)$$

In the classical limit  $\hbar \rightarrow 0$ , the first factor in the r.h.s. of Eq. (26) gives the classical spectral density of the Langevin force,  $2kT\gamma$ , while the second factor is identically equal to unity for the case of a classical viscous friction, i.e. when  $Im\{\alpha_x^{-1}(\omega)\} = -\gamma\omega$ . By assuming that the second factor keeps the same value also at  $\hbar \neq 0$ , one obtains the quantum generalization of the thermal law of viscous friction in spectral representation as:

$$Im\{\alpha_x^{-1}(\omega)\} = \frac{\gamma}{\beta} (e^{-\frac{\beta\omega}{2}} - e^{\frac{\beta\omega}{2}}) \quad (27)$$

which satisfies the second requirement of MiDEB in the whole complex frequency plane. The corresponding time representation is given by expression (4). In so doing, the first factor in the r.h.s of Eq. (25) gives the quantum analog of the spectral density of the thermal Langevin force which corresponds to the correlation function given by Eq. (5).

In conclusion, we have proposed a thermal quantum viscous friction law at a macroscopic level. This law leads to a macroscopic quantum Langevin equation that does not include the zero point contribution in the spectral density of the fluctuating observable under thermal equilibrium conditions. The most relevant implications of this generalization are: (i) in the classical limit  $\hbar \rightarrow 0$  the quantum Langevin equation recovers the classical one as it should; (ii) it resolves several anomalies appearing in the quantum extensions of the fluctuation dissipation and regression theorems, such as the well known QRT-QFDT conflict, and various divergencies originated by the Callen-Welton form of the QFDT; (iii) the prediction of an exponential like decay of the susceptibility at  $\omega \gg kT/\hbar$ .

*Acknowledgments.* Partial support from European Commission through project No. IST2001-38899 is gratefully acknowledged.

<sup>1</sup> N.G. Van Kampen, *Fluctuation Noise Letters*, **1**, 3 (2001).

<sup>2</sup> R. Benguria and M. Kac, *Phys. Rev. Lett.*, **46**, 1 (1980).

<sup>3</sup> C.W. Gardiner, *Quantum Noise*, Springer Verlag (Berlin, 1981).

<sup>4</sup> P. Meystre, M. Sargent, *Elements of Quantum Optics* (Springer-Verlag, 1991).

<sup>5</sup> H. Grabert, U. Weiss, P. Talkner, *Z. Phys. B*, **55** 87 (1984).

- <sup>6</sup> H.B. Callen and T.A. Welton, *Phys. Rev.* **83**, 34 (1951).
- <sup>7</sup> P. Talkner, Ann. Phys. (N.Y.) **167**, 380 (1986).
- <sup>8</sup> G.W. Ford and R.F. O'Connell, Phys. Rev. Lett., **77**, 798 (1996).
- <sup>9</sup> M. Lax, Phys. Rev., **129**, 2342 (1963).
- <sup>10</sup> H. Nyquist, *Phys. Rev.* **32**, 110 (1928).
- <sup>11</sup> N.G. Van Kampen: Stochastic processes in physics and chemistry, North Holland, Amsterdam (1992)
- <sup>12</sup> D.N. Zubarev, *Nonequilibrium Statistical Thermodynamics*, (Nauka, Moscow, 1971)
- <sup>13</sup> R. Kubo, M. Toda, N. Hashitsume, *Statistical Physics II* (Spriger-Verlag, Berlin, 1985)
- <sup>14</sup> R. Loudon, *The Quantum Theory of Light* (Clarendon Press, Oxford, 1973)
- <sup>15</sup> C.H. Henry and F. Kazarinov, Rev. Mod. Phys. **68** 801 (1996).
- <sup>16</sup> U. Gavish, Y. Levinson and Y. Imry, Phys. Rev. **B62**, R10637 (2000).