

Comment on "Relevant Length Scale of Barchan Dunes"

In a recent experimental breakthrough, Hersen et al. [1] demonstrated that by changing the agitating medium from air to water, one can obtain, on laboratory scale, dunes that are downsized copies of desert dunes, thereby overcoming a major obstacle for their systematic study. Here we argue in two steps (i), (ii) that an alternative data analysis leads to some conclusions that are qualitatively and quantitatively different from Hersen et al.'s but justify their similarity hypothesis on different grounds.

It has been a long-standing debate whether the various shapes of desert dunes share the scale-invariance of the turbulent wind field that creates them. Careful field measurements of barchan dunes have recently provided compelling evidence that scale invariance is borne out only transverse to the wind direction, whereas it is systematically broken along the wind direction [2]. That the responsible characteristic scale is the windward distance  $\lambda_s$  over which the sand flux adapts to a sudden change in wind speed or sand supply, was already vaguely anticipated by Bagnold. But only recent theoretical developments succeeded in making this notion fully quantitative and established its crucial role for systematic shape variations and shape transitions of aeolian dunes [3].

(i) The analysis of similarities of aeolian and submarine dunes in Ref. [1] suffers from mixing transverse and longitudinal dune dimensions. From what was said above, a genuine test of similarity should address the longitudinal geometry directly, thereby avoiding inferences [2] from uncontrolled parameters affecting the transverse direction. Recombining the data from Ref. [1], we can infer the (a) aeolian and submarine height-length ( $H/L$ ) relations that suggest a considerably larger scale factor than estimated in [1]. Our best estimate for the latter is however obtained from analysing data [1, 4] for the dune migration speed  $V$  as follows.

(ii) The (trivial) fact that for a barchan dune, which loses practically no sand over its crest,  $V$  is proportional to the flux  $q_c$  over the crest divided by its height, is the origin of the popular relation  $V/H^{1/2}$  adopted in Ref. [1]. Note however that the latter would only hold for size-independent  $q_c$ , i.e. in the case of scale-invariant dune shape, inconsistent with the observation of broken scale invariance crucially underlying the similarity hypothesis by Hersen et al. Yet, one can show on quite general grounds [3] that (essentially)  $V/L^{1/2}$ ,  $L$  being the total length of either smooth heaps or barchan dunes. Fig. 1 demonstrates that this (but not  $V/H^{1/2}$ ) is compatible with both aeolian and submarine data. To superimpose the data, lengths had to be rescaled by  $6.2 \cdot 10^3$  (in line with our recombined  $H/L$  relations) instead of the  $1.7 \cdot 10^3$  suggested in [1], and times by  $2.3 \cdot 10^3$ .

Assuming  $\lambda_s^{\text{air}} = 0.8 \text{ m}$  [3] for the flux-weighted average of  $\lambda_s$  [5] this implies  $\lambda_s^{\text{water}} = 0.13 \text{ mm}$  equal to the grain diameter, as naturally expected for direct fluid entrainment.




FIG. 1: Submarine (black [1]) and aeolian (gray [4]; rescaled) migration speed data compared to  $H^{-1/2}$  (dotted [1]) failing for small dunes, and  $L^{-1/2}$  (solid [3]) compatible with the data.

ment. In conclusion, we agree with Ref. [1] that there is an intriguing similarity between submarine and aeolian dunes, but on somewhat different grounds. Further quantitative confirmation of this similarity by data for small submarine dunes, backing the highest submarine data point in Fig. 1, would be desirable.

Klaus Kroy<sup>1</sup> and Xiang Gou<sup>2</sup>

<sup>1</sup>Hahn-Meitner Institut

Geilicker Str. 100, 14109 Berlin, Germany

<sup>2</sup>Department of Building and Construction,  
City University of Hong Kong, Hong Kong

- 
- [1] P. Hersen, S. Douady, and B. Andreotti, Phys. Rev. Lett. 89, 264301 (2002).
  - [2] G. Sauermann, P. Rognon, A. Poliakov, and H. J. Herrmann, Geomorphology 36, 47 (2000).
  - [3] G. Sauermann, K. Kroy, and H. J. Herrmann, Phys. Rev. E 64, 031305 (2001); K. Kroy, G. Sauermann, and H. J. Herrmann, Phys. Rev. Lett. 88, 054301 (2002), and Phys. Rev. E 66, 031302 (2002).
  - [4] H. J. Finkel, J. Geol. 67, 614 (1959).
  - [5] While the average saltation length, the scale  $\lambda_{\text{drag}}$  introduced in Ref. [1], and  $\lambda_s$  are all proportional to each other on dimensional grounds, their absolute values are generally strongly diverse due to distinctly different dependencies on the wind speed [3], which prohibits their identification even for order-of-magnitude estimates.