

# Ground state, vibrational spectrum and deterministic transport of a chain of charged particles

S. I. Denisov and E. S. Denisova

Department of Mechanics and Mathematics, Sumy State University,  
2, Rimskiy-Korsakov Street, 40007 Sumy, Ukraine

## Abstract

We study the equilibrium, oscillatory and transport properties of a chain of charged particles which interact with each other via the Coulomb and power-like repulsive interactions. Exact, analytical expressions for the energy of the ground state, interparticle distances, and vibration spectrum are derived, and the stability criterion for a chain with equidistant particles is established. We also consider the phenomenon of deterministic transport of a chain in a ratchet potential subject to a longitudinal alternating electric field.

PACS numbers: 45.05.+x, 05.60.Cd, 45.90.+t

## I. INTRODUCTION

A chain of coupled oscillators is a universal mathematical model that describes a large variety of phenomena in physics, chemistry, biology and other sciences. Examples include energy localization and propagation,<sup>1,3</sup> discrete solitons,<sup>4,5</sup> chemical reactions,<sup>6</sup> nonequilibrium phase transitions,<sup>7,8</sup> stochastic resonance,<sup>9,10</sup> synchronization,<sup>11</sup> brain dynamics<sup>12</sup> and many others. A special case of that model, a chain of charged particles that interact via the Coulomb interaction, also has a variety of applications. In particular, if the charges have the same sign and are confined by a trapping potential, it describes the so-called one-dimensional Coulomb crystals, i.e., a crystallized structure of ions in a linear trap formed by external focusing fields.<sup>13,15</sup> If neighboring charges have opposite signs and a repulsive interaction exists between particles, then a such chain represents a relevant model for one-dimensional ionic crystals. It seems that those models can be useful also for the study of one-dimensional crystals within carbon nanotubes.<sup>16,17</sup>

During the last decade, much attention was devoted to the transport properties of particles in thermal ratchets, i.e., spatially periodic stochastic models exhibiting noise-induced macroscopic transport.<sup>18,27</sup> The ratchet effect or stochastic transport in asymmetric potentials is important for biological motors,<sup>28,29</sup> separation techniques,<sup>30</sup> surface smoothing<sup>31</sup> and vortex density reduction,<sup>32</sup> to name a few. Most studies of ratchets, including deterministic ones,<sup>33,34</sup> i.e., ratchets without internal and external noises, have been performed for a single particle. However, examples of coupled particles<sup>35,36</sup> and vortices<sup>37</sup> show that their collective interactions can significantly modify the transport properties from the single particle case. To our knowledge, the transport properties of a chain of strongly coupled charged particles in deterministic ratchets have not been studied at all.

In this paper, we study the equilibrium, oscillatory and transport properties of a chain where the neighboring particles have the same mass and opposite charges. Particles interact with each other via both the Coulomb interaction and a repulsive interaction that depends on the interparticle distance as  $|x_i - x_j|^r$  ( $x_i = x_i(t)$  is the coordinate of the  $i$ th particle,  $r > 1$ ). In Sec. II, we consider the ground state of such a chain depending on the exponent  $r$ . Three different states of a chain, namely, a periodic chain with equidistant particles, a periodic chain with non-equidistant particles, and a chain with infinite period, are analyzed, and the interparticle distances and the chain energies as functions of  $r$  are calculated. In

contrast to intuitive expectations, we show that there is a critical value of the exponent,  $r = r_2 \approx 2.799$ , such that for  $r > r_2$  the minimum energy occurs for a chain with equidistant particles, and for  $r < r_2$  the minimum energy occurs for a chain with infinite period.

Propagation of longitudinal waves in a chain with equidistant particles, its stability with respect to small displacement of particles from their equilibrium positions, and forced vibrations of particles under action of an alternating electric field are studied in Sec. III. We derive the set of equations that describe small oscillations of the particles, find the phonon spectrum and its asymptotics near the center and boundary of the Brillouin zone, and calculate the amplitude of forced vibrations. We show that the spectrum consists of one branch; on the boundary of the Brillouin zone it has a non-analyticity which is caused by the long-range action of the Coulomb interaction, and a chain is stable only if  $r > r_1 \approx 2.779$ .

The deterministic transport of a chain is studied in Sec. IV. We consider the case when the ratchet potential is charge independent, and a chain is driven by a longitudinal alternating electric field. We find the average velocity of a chain and analyze in detail its dependence on the parameters characterizing the interparticle interactions, ratchet potential and alternating electric field. Concluding remarks are contained in Sec. V, and some technical details are given in the Appendixes.

## II. GROUND STATE

We consider a chain of particles which have the same mass  $M$  and any neighboring particles have opposite charges, i.e.,  $q_{i+1} = -q$  and  $q_i = q$  (the index  $i$  labels the particles). We assume that in addition to the Coulomb interaction a repulsive interaction exists between the particles. In analogy to ionic crystals,<sup>38</sup> we choose a repulsive interaction that depends on a power of the interparticle distance  $|x_i - x_j|$ , and write the total interaction energy of a chain in the form

$$W_{\text{int}} = \frac{1}{2} \sum_{i \neq j} \frac{q_i q_j}{|x_i - x_j|} + \frac{b}{2} \sum_{i \neq j} \frac{1}{|x_i - x_j|^r} : \quad (2.1)$$

Here  $b$  is a dimensioned parameter characterizing the strength of the repulsive interaction, and the primes on the summation signs indicate the absence of self-interaction terms, i.e.,  $i \neq j$ . To prevent the unbounded contraction of a chain, we require  $r > 1$ .

If a chain is periodic in equilibrium (see Fig. 1), then the coordinates of the positive and negative charges are written as  $np + x_n^+$  and  $np + a + x_n^-$ , respectively. Here  $p$  is the chain

period,  $n$  ( $n = 0; 1; \dots; 1$ ) numbers the chain cells which contain two particles,  $a$  is the interparticle equilibrium distance in the cells ( $a < p$ ), and  $x_n^+ = x_n^+(t)$  and  $x_n^- = x_n^-(t)$  are the displacements of positive and negative charges from their equilibrium positions. With these definitions Eq. (2.1) reads

$$W_{int} = \frac{q^2 x_0^2}{2_{nm}} \frac{1}{j(n-m)p + a + x_n^+ - x_m^-} + \frac{1}{j(n-m)p + x_n^+ - x_m^-} + \frac{1}{j(n-m)p + x_n^- - x_m^+} + \frac{q^2 x_0^2}{2_{nm}} \frac{1}{j(n-m)p + x_n^+ + x_m^-} + \frac{b x_0^2}{2_{nm}} \frac{1}{j(n-m)p + x_n^+ - x_m^-} + \frac{1}{j(n-m)p + x_n^- - x_m^+} + \frac{b x_0^2}{2_{nm}} \frac{1}{j(n-m)p + x_n^- - x_m^+}; \quad (2.2)$$

where summation on  $n$  and  $m$  is taken over all integers. Note also that the particles cannot move past each other, and therefore the conditions  $a > x_n^+ - x_n^-$  and  $p - a > x_n^- - x_{n+1}^+$  must hold for all  $n$  and  $t$ .

In equilibrium  $x_n^+ = x_n^- = 0$ , and Eq. (2.2) is reduced to the form

$$W_{int}^{(0)} = N \frac{q^2}{2a} - N \frac{q^2 x^2}{2p} \sum_{n=1}^{\infty} \frac{1}{n} + \frac{1}{n+1} - \frac{2}{n} + N \frac{b}{p^r} \zeta(r) + N \frac{b x}{2p^r} \sum_{n=1}^{\infty} \frac{1}{j^2 + j^r}; \quad (2.3)$$

where  $N$  is the number of particles in a chain ( $N \neq 1$ ),  $\zeta = a/p$  ( $0 < \zeta < 1$ ), and  $\zeta(r) = \sum_{n=1}^{\infty} n^{-r}$  is the Riemann zeta function. Taking into account that the energy  $W_{int}^{(0)}$  is invariant with respect to the replacement of  $\zeta$  by  $1/\zeta$ , we conclude that such chains are equivalent. Therefore, the permissible values of  $\zeta$  can be restricted by the condition  $0 < \zeta \leq 1/2$ .

Defining the dimensionless parameter  $\eta = p/a$  ( $0 < \eta < 1$ ) and the length scale  $l = (b/q^2)^{1/(r-1)}$ , we can rewrite Eq. (2.3) as  $W_{int}^{(0)} = N (q^2/a) U_0(\eta; r)$ , where

$$U_0(\eta; r) = \frac{1}{2} - \frac{\eta^2}{2} \sum_{n=1}^{\infty} \frac{1}{n(n^2 - \eta^2)} + \frac{\zeta(r)}{r}$$

$$+ \frac{1}{2} \sum_{r=1}^{\infty} \frac{1}{j_1 + j_r} \quad (2.4)$$

is the dimensionless chain energy per particle (for another representation of  $U_0(\beta; \alpha)$  see Appendix A). The equilibrium values of  $\alpha$  and  $\beta$  are defined by the set of equations  $\partial U_0(\beta; \alpha) / \partial \alpha = 0$  and  $\partial U_0(\beta; \alpha) / \partial \beta = 0$ , which are written as

$$\frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{(n^2 - \alpha^2)^2} - \frac{1}{2} \sum_{n=1}^{\infty} \frac{n}{j_1 + j_{r+2}} = 0; \quad (2.5)$$

$$\frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n(n^2 - \alpha^2)} - \frac{1}{2} \sum_{r=1}^{\infty} \frac{1}{j_1 + j_r} = 0; \quad (2.6)$$

Analysis shows that these equations have three solutions:  $\alpha = \alpha_s(r)$  and  $\beta = \beta_s(r)$  ( $s = 1, 2, 3$ ). Below we find  $\alpha_s(r)$  and  $\beta_s(r)$ , and calculate the chain equilibrium energies  $U_s(r) = U_0(\alpha_s(r); \beta_s(r))$ . For illustrative purposes we introduce also the parameters  $\alpha_s(r) = \alpha_s(r) / \alpha_s(r)$ , which define the equilibrium values of the parameter  $\alpha = \alpha_s(r) / \alpha_s(r) = 1$ .

#### A. First solution

For  $\alpha = 1/2$  ( $\beta = 2\alpha$ ), i.e., for a chain with equidistant particles, the series in Eqs. (2.4) { (2.6) are either standard series or can be reduced to them. Indeed, using standard series,<sup>39</sup> we find with linear accuracy in  $\alpha = 1/2$  ( $\beta = 1$ ):

$$\sum_{n=1}^{\infty} \frac{1}{(n^2 - \alpha^2)^2} = \frac{1}{2} \left[ \frac{1}{3} - \frac{1}{6} \right]; \quad (2.7)$$

$$\sum_{n=1}^{\infty} \frac{n}{j_1 + j_{r+2}} = \frac{1}{2} (r+1) (2^{r+2} - 1) (r+2); \quad (2.8)$$

$$\sum_{n=1}^{\infty} \frac{1}{n(n^2 - \alpha^2)} = \frac{1}{4} (2 \ln 2 - 1) - \frac{1}{8} (3 - 4 \ln 2); \quad (2.9)$$

$$\sum_{n=1}^{\infty} \frac{1}{j_1 + j_r} = \frac{1}{2} (2^r - 1) (r); \quad (2.10)$$

According to these formulas,  $U_0(1/2; \beta)$  is written as

$$U_0(1/2; \beta) = \frac{1}{2} \ln 2 + \frac{1}{2} \sum_{r=1}^{\infty} \frac{1}{j_1 + j_r} (r); \quad (2.11)$$

at  $\beta = 1/2$  Eq. (2.5) is satisfied identically, i.e.,  $\phi_1(r) = 1/2$ , and Eq. (2.6), which at  $\beta = 1/2$  is equivalent to the equation  $dU_0(1/2; r)/dr = 0$ , yields

$$\phi_1(r) = 2 \frac{r}{\ln 2} \frac{1}{r^{1/(r-1)}} : \quad (2.12)$$

Substituting  $\phi = \phi_1(r)$  into Eq. (2.11), we find for the equilibrium chain

$$U_1(r) = -\ln 2 \frac{r}{r-1} \frac{1}{r^{1/(r-1)}} \frac{\ln 2}{r} : \quad (2.13)$$

[Note that  $U_1(r)$  is the minimum value of  $U_0(1/2; r)$  only if  $r > 1$ .]

The plots of the functions  $\phi_1(r)$ ,  $\phi_2(r)$ , and  $U_1(r)$  are shown in Figs. 2, 3 and 4, respectively. To gain more insight into the behavior of these functions near  $r = 1$  and  $r = \infty$ , we define the limiting values of those functions as  $r \rightarrow 1$  and their asymptotics as  $r \rightarrow \infty$ . Taking into account that  $\lim_{x \rightarrow 0} x^x = 1$ ,  $\lim_{x \rightarrow 0} (1+x)^{1/x} = e$ ,  $\phi_1(1) = 1/2$ , and  $\phi_2(r) \sim (r-1)^{-1}$  as  $r \rightarrow 1$ , we obtain from Eq. (2.12)  $\phi_1(1) = 1/2$  and  $\phi_1(r) = 2e(\ln 2)^{1/(r-1)} \rightarrow 1$  as  $r \rightarrow \infty$ , and from Eq. (2.13)  $U_1(1) = -\ln 2$  and  $U_1(r) \sim e^{1/(r-1)} (\ln 2)^{1/(r-1)} \rightarrow 0$  as  $r \rightarrow \infty$ . Thus, if  $r \rightarrow 1$  then the functions  $\phi_1(r)$  and  $U_1(r)$  go extremely fast to infinity and zero, respectively. From a physical point of view, this behavior of  $\phi_1(r)$  and  $U_1(r)$  is caused by the unlimited growth of the total repulsive energy of a chain as  $r \rightarrow 1$ .

## B. Second solution

If  $\beta \neq 1/2$ , then we eliminate  $\phi$  from Eq. (2.5) with the help of Eq. (2.6) and obtain the equation

$$\frac{1}{2} - 4 \sum_{n=1}^{\infty} \frac{n}{(n^2 - 1/2)^2} \phi_2(r) + \sum_{n=1}^{\infty} \frac{1}{n + j^r} = \frac{1}{2} + 2 \sum_{n=1}^{\infty} \frac{1}{n(n^2 - 1/2)} \sum_{n=1}^{\infty} \frac{n + j^{r+2}}{n + j^{r+2}} = 0; \quad (2.14)$$

which defines the equilibrium values of  $\phi$ . According to the results of the previous subsection, Eq. (2.14) always has the solution  $\phi = 1/2$ , i.e.,  $\phi_1(r) = 1/2$ . To determine if Eq. (2.14) has solutions on the interval  $(0; 1/2)$ , we find first the asymptotics of its left-hand side, denoted by  $L_r(\phi)$ , as  $\phi \rightarrow 0$  and  $\phi \rightarrow 1/2$ . If  $\phi \rightarrow 0$ , then

$$L_r(\phi) = \left[ \frac{1}{2} - 4 \sum_{n=1}^{\infty} \frac{n}{(n^2 - 1/2)^2} \right] \left[ \frac{1}{r+2} \phi(r) \right] \left[ \frac{1}{1+2} \sum_{n=1}^{\infty} \frac{n}{(n^2 - 1/2)} \right]^{(r+1)}; \quad (2.15)$$

and we obtain different asymptotics of  $L_r(\epsilon)$  for different values of  $r$ :

$$L_r(\epsilon) \sim \begin{cases} 6(3)^{-(r-1)}; & r > 3; \\ 2(3)^{-2}; & r = 3; \\ + 4(r)^{-2}; & r < 3; \end{cases} \quad (2.16)$$

Using Eqs. (2.7)–(2.10), as  $\epsilon \rightarrow 0$  we obtain

$$L_r(\epsilon) \sim 2^{r+3} (r) F(r); \quad (2.17)$$

where

$$F(r) = 4(1 - 2^{-r-2})(r+1) \ln 2 - \frac{(r+2)}{(r)} \quad (2.18)$$

is a monotone increasing function of  $r$ , and  $F(r) = 0$  at  $r = r_1 \approx 2.779$ . Therefore, in the left vicinity of the point  $\epsilon = 0$ , the conditions  $L_r(\epsilon) > 0$  and  $L_r(\epsilon) < 0$  hold for  $r < r_1$  and  $r > r_1$ , respectively. Using this fact and Eq. (2.16), we conclude that if  $r_1 < r < 3$ , then  $L_r(\epsilon)$  near the points  $\epsilon = 0$  and  $\epsilon = 1/2$  has different signs, and so Eq. (2.14) has at least one solution  $\epsilon_2(r) \in (0; 1/2)$  satisfying the conditions  $\epsilon_2(r) \rightarrow 0$  as  $r \rightarrow r_1$  and  $\epsilon_2(r) \rightarrow 0$  as  $r \rightarrow 3$ . Numerical analysis shows that such a solution is unique, and Eq. (2.6) yields

$$\epsilon_2(r) = \frac{2}{4} r \frac{2^{-r} + \sum_{n=1}^{\infty} \frac{1}{n^{r+2}(r)^n}}{\frac{1}{2^{-r}} + 2 \sum_{n=1}^{\infty} \frac{1}{n^{r+2}(r)^n}} : \quad (2.19)$$

Since a chain is transformed to a chain with equidistant particles as  $r \rightarrow r_1$ , the equalities  $\epsilon_2(r_1) = \epsilon_1(r_1)$ ,  $\epsilon_2(r_1) = \epsilon_1(r_1)$ , and  $U_2(r_1) = U_1(r_1)$  hold. If  $r \rightarrow 3$ , then according to Eq. (2.19)  $\epsilon_2(r) \rightarrow \frac{1}{3}$ , and we obtain  $\epsilon_2(3) = \frac{1}{3}$ ,  $U_2(3) = \frac{1}{3}$ . The plots of the functions  $\epsilon_2(r)$ ,  $\epsilon_1(r)$ , and  $U_2(r)$ , which are defined for  $r \in [r_1; 3]$ , are also shown in Figs. 2–4. As is seen from Fig. 4,  $U_2(r) > U_1(r)$  for  $r_1 < r < 3$ , i.e., the equilibrium chain with equidistant particles is energy-wise preferred compared to the equilibrium chain with non-equidistant particles.

### C. Third solution

A further solution of Eqs. (2.5) and (2.6) corresponds to the case when the period of the chain is infinite and the distance between particles in the cells is finite, i.e., when  $\epsilon \rightarrow 0$ ,

$\neq 1$ , and  $\neq 0$ . In this case

$$U_0(r; \epsilon) - U_0(r) = \frac{1}{2} + \frac{1}{2r}; \quad (2.20)$$

Eq. (2.6) is satisfied identically, and Eq. (2.5) is reduced to the equation  $dU_0(r)/dr = 0$ . Its solution  $r = r_3(r)$ ,  $r_3(r) = r^{1/(r-1)}$ , defines the equilibrium distance between the particles, and the equilibrium energy per particle is given by

$$U_3(r) = \frac{r-1}{2r} - \frac{1}{r^{1/(r-1)}}; \quad (2.21)$$

The functions  $r_3(r)$  and  $U_3(r)$  monotonically decrease as  $r$  grows from 1 to  $\infty$  [ $U_3(r_3(r)) = 0$  as  $r \rightarrow 1$ ,  $U_3(3) = -1/3$ ,  $U_3(1) = -1/2$ ,  $r_3(1) = e$ ,  $r_3(3) = 3/2$ ,  $r_3(1) = 1$ ], and their plots are shown in Figs. 3 and 4. Comparing the functions  $U_1(r)$ ,  $U_2(r)$ , and  $U_3(r)$ , we conclude that for  $r > r_2$  [ $r = r_2 = 2.799$  is the solution of the equation  $U_1(r) = U_3(r)$ ] the minimum energy occurs for a chain with equidistant particles [ $p = 2a$ ,  $a = 1/r_3(r)$ ], and for  $r < r_2$  the minimum energy occurs for a chain with infinite period [ $p = 1$ ,  $a = 1/r_3(r)$ ]. Note also that a chain with non-equidistant particles has a slightly larger energy than other chains.

Next we study the vibration properties of a chain with equidistant particles.

### III. VIBRATION PROPERTIES

#### A. Dispersion relation

Assuming that the condition  $|x_n| \ll a$  holds for all particles, we expand the chain energy  $W_{int}$  in a power series of  $x_n = a$  with quadratic accuracy:  $W_{int} = W_{int}^{(0)} + W_{int}^{(1)} + W_{int}^{(2)}$ , where the upper indices indicate the power of  $x_n$ . Straightforward calculations based on Eq. (2.2) with  $p = 2a$  and arbitrary value of  $a$  yield  $W_{int}^{(0)} = N(q^2=1)U_0(1/2; 2a=1)$ ,  $W_{int}^{(1)} = 0$ , and

$$W_{int}^{(2)} = \frac{1}{4} \sum_{n,m} B_{2(n-m)} (x_n^+ - x_m^+)^2 + (x_n - x_m)^2 + \frac{1}{2} \sum_{n,m} B_{2(n-m)-1} (x_n^+ - x_m)^2; \quad (3.1)$$

where

$$B_n = \frac{(n-1)^2 2q^2}{a^3 n^3} + \frac{br(r+1)}{a^{r+2} n^{r+2}}; \quad (3.2)$$



For the case considered here that  $a$  has an equilibrium value, i.e.,  $a = a_1(r)$ , we represent  $B_n$  in the form  $B_n = M^{-2} \tilde{B}_n$ , where  $M^2 = \alpha^2 M l^3$ ,

$$\tilde{B}_n = \frac{1}{l^3(r)} \left[ 2 \frac{(l-1)^2}{j^3} + \frac{(r+1) \ln 2}{(r) j^2} \right]; \quad (3.3)$$

and according to Eq. (2.12)  $a_1(r) = [r(r) \ln 2]^{1/(r-1)}$ .

From Eq. (3.1), we obtain the system of equations,  $M \ddot{x}_n + \partial W_{int}^{(2)} / \partial x_n = 0$ , that describe free longitudinal vibrations of particles near their equilibrium positions in the harmonic approximation, in the form

$$\begin{aligned} \ddot{x}_n^+ + \sum_m^X \tilde{B}_{2(n-m)+1} (x_n^+ - x_m^+) \\ + \sum_{m \in n}^X \tilde{B}_{2(n-m)} (x_n^+ - x_m^+) = 0; \\ \ddot{x}_n + \sum_m^X \tilde{B}_{2(n-m)+1} (x_n - x_m) \\ + \sum_{m \in n}^X \tilde{B}_{2(n-m)} (x_n - x_m) = 0; \end{aligned} \quad (3.4)$$

Since  $n$  ranges over the integers, this system contains infinitely many coupled equations. We seek wave solutions, which in the complex representation have the form

$$x_n = x \exp(i2nak - i\omega t); \quad (3.5)$$

Here  $x_+$  and  $x_-$  are the complex amplitudes,  $k$  is the wave number belonging to the Brillouin zone, i.e., to the region  $[-2a; 2a]$  of reciprocal space, and  $\omega$  is the wave frequency. Substituting solution (3.5) into Eqs. (3.4) and using the definition  $\omega = ak = (2\pi/l) [l=2; l=2]$ ,

$$D = 2 \sum_{p=1}^X \tilde{B}_p; \quad (3.6)$$

$$P(\omega) = 2 \sum_{p=1}^X \tilde{B}_{2p} \cos(2p\theta); \quad (3.7)$$

$$S(\omega) = \sum_p^X \tilde{B}_{2p-1} \exp(-i2p\theta); \quad (3.8)$$

we obtain the system of equations for  $x_+$  and  $x_-$

$$\begin{aligned} x_+ [-\omega^2 + D - P(\omega)] - x_- S(\omega) &= 0; \\ x_- S(\omega) + x_+ [-\omega^2 + D - P(\omega)] &= 0; \end{aligned} \quad (3.9)$$

where  $\omega = \omega_0$  is the dimensionless wave frequency, and the asterisk denotes complex conjugation. This system has non-trivial solutions, i.e., propagation of phonons along a chain can occur, if its determinant equals zero,<sup>41</sup>

$$[\omega^2 - D + P(\omega)]^2 - \mathcal{J}(\omega)^2 = 0; \quad (3.10)$$

Solving Eq. (3.10) with respect to  $\omega$ , we find the dispersion relation for phonons:  $\omega_{1,2}(\omega) = \omega_{1,2}(\omega)$ , where

$$\omega_{1,2}(\omega) = \sqrt{D - P(\omega) \pm \mathcal{J}(\omega)}; \quad (3.11)$$

and the plus and minus sign in front of  $\mathcal{J}(\omega)$  correspond to  $\omega_1(\omega)$  and  $\omega_2(\omega)$ , respectively.

## B. Analysis of the spectrum

Since  $\omega_{1,2}(\omega) = \omega_{1,2}(\omega)$ , we can restrict the permissible values of the dimensionless wave number by the condition  $\omega \in [0; 1/2]$ . According to Eq. (3.11), in this interval the dimensionless phonon spectrum consists of two branches,  $\omega_1(\omega)$  and  $\omega_2(\omega)$ . It is easy to see that on the interval  $[0; 1/2]$  the spectrum is given by

$$\omega(\omega) = \begin{cases} \omega_1(\omega); & 0 \leq \omega \leq 1/2; \\ \omega_2(1/2 - \omega); & 1/2 < \omega \leq 1; \end{cases} \quad (3.12)$$

and  $\omega(\omega)$  is continuous at  $\omega = 1/2$ . Indeed, for  $\omega = 1/2$  Eq. (3.8) yields  $S(1/2) = P_p(1/2) \mathcal{B}_{2p-1}$ . Replacing in this formula  $p$  by  $p+1$  and using the condition  $\mathcal{B}_n = \mathcal{B}_{-n}$ , we obtain  $S(1/2) = P_p(1/2) \mathcal{B}_{2p-1}$ . This implies that  $S(1/2) = 0$ , and Eq. (3.11) yields  $\omega_1(1/2) = \omega_2(1/2)$ . Hence, the phonon spectrum on the interval  $[0; 1/2]$  is represented by the single branch  $\omega(\omega)$ . Note that the region  $[-1/2; 1/2]$  is the dimensionless Brillouin zone corresponding to a chain with a primitive cell which contains one particle.

To study the behavior of the phonon spectrum  $\omega(\omega)$ , we use Eq. (3.3) to represent Eqs. (3.6)–(3.8) in the form

$$D = \frac{1}{2} \frac{1}{\omega^2} \ln 2 \frac{(r+2)}{\omega(r)} \quad (3.13)$$

$$P(\omega) = \frac{1}{2} \frac{1}{\omega^2} \frac{(r+1) \ln 2}{\omega(r)} \quad (3.14)$$

$$S(\omega) = \frac{4e^{-i}}{2} \frac{(r+1) \ln 2}{\omega(r)} \quad (3.15)$$

where

$$s(\epsilon) = \sum_{p=1}^{\infty} \frac{x^{2p}}{p^s}; \quad (3.16)$$

$$s(\epsilon) = \sum_{p=1}^{\infty} \frac{x^{2p-1}}{(2p-1)^s}; \quad (3.17)$$

[Recall that the functions  $P(\epsilon)$  and  $S(\epsilon)$  are defined for  $j = 1, 2$ .] Unfortunately, the analytical summation of the series in Eqs. (3.14) and (3.15) can be performed only for special cases, such as  $\epsilon = 0$  or  $\epsilon = 1$ . We calculate  $\epsilon(\epsilon)$  only in the centre ( $\epsilon = 0$ ) and on the boundary ( $\epsilon = 1$ ) of the Brillouin zone which, according to Eq. (3.12), requires us to determine  $P(0)$  and  $S(0)$ . Using the definition of the Riemann zeta function and the standard series  $\sum_{p=1}^{\infty} (2p-1)^{-s} = (1-2^{-s})\zeta(s)$ , we obtain

$$P(0) = \frac{1}{2^{3/2}} \sum_{r=1}^{\infty} \frac{\ln 2 - \frac{(r+2)}{r}}{r} + \quad (3)$$

and  $D = P(0) = S(0) = F(r) = 2^{3/2} \sum_{r=1}^{\infty} \frac{1}{r}$ , where  $F(r)$  is defined by Eq. (2.18). If  $F(r) > 0$ , i.e.,  $r > r_1$ , then

$$\epsilon(0) = 0; \quad \epsilon(1) = \frac{q}{F(r) = 2^{3/2}}; \quad (3.19)$$

Note that Eqs. (3.9) yield  $x_+ = x_-$  for  $\epsilon = 0$ , and  $x_+ = -x_-$  for  $\epsilon = \epsilon(1)$ . This means that neighboring particles in a chain perform acoustical vibrations at  $\epsilon = 0$  and optical ones at  $\epsilon = 1$ .

To find for  $r > r_1$  the asymptotics of  $\epsilon(\epsilon)$  as  $\epsilon \rightarrow 0$  and  $\epsilon \rightarrow 1$ , we need to calculate the series (3.16) and (3.17) with quadratic accuracy in  $\epsilon$  as  $\epsilon \rightarrow 0$ . This is not a problem for the series  $\sum_{r=2}^{\infty} \frac{1}{r^2}$  and  $\sum_{r=2}^{\infty} \frac{1}{r^3}$  that arise from the repulsive interaction of particles, because expanding the cosines in a power series of  $\epsilon$  to an accuracy of  $\epsilon^2$  we immediately obtain

$$\sum_{r=2}^{\infty} \frac{1}{r^2} = \frac{\pi^2}{6} - \frac{1}{2}; \quad (3.20)$$

$$\sum_{r=2}^{\infty} \frac{1}{r^3} = \frac{\pi^2}{12} - \frac{1}{2}; \quad (3.21)$$

As to the series  $\sum_{r=1}^{\infty} \frac{1}{r^2}$  and  $\sum_{r=1}^{\infty} \frac{1}{r^3}$  that arise from the Coulomb interaction, their calculation by this method is impossible since divergencies occur. To avoid this difficulty, we use their integral representations obtained in Appendix B. These representations have the advantage that the double integrals in Eqs. (B3) and (B6) can be easily calculated with arbitrary

accuracy as  $\epsilon \rightarrow 0$ . Specifically, if we use the approximate formulas  $\ln [2 \sin (\gamma/2)] \approx \ln \gamma$  and  $\ln [\tan (\gamma/2)] \approx \ln (\gamma/2) - (\gamma/4)$ , then Eqs. (B3) and (B6) yield

$$\epsilon_3(\gamma) = \epsilon_3(0) - \frac{1}{2} \beta \ln^2(2\gamma); \quad (3.22)$$

$$\epsilon_3(\gamma) = \frac{7}{8} \epsilon_3(0) - \frac{1}{8} \beta \ln^2 \frac{\gamma}{2}; \quad (3.23)$$

Substituting series (3.20)–(3.23) into Eqs. (3.14) and (3.15), we find the main terms of the asymptotic expansion of  $\epsilon(\gamma)$  as  $\epsilon \rightarrow 0$ ,

$$\epsilon(\gamma) \approx \frac{\epsilon_{\text{cut}}}{\epsilon_1(r)} \frac{(r-1) \ln 2}{3}; \quad (3.24)$$

and as  $\epsilon \rightarrow 1$ ,

$$\epsilon(\gamma) \approx \epsilon(1) + \frac{1}{\epsilon_1(r)} (1-\gamma)^2 j \ln(1-\gamma); \quad (3.25)$$

According to these asymptotics, the long-wave part of the vibrational spectrum (for  $\epsilon \rightarrow 1$ ) has the usual form, i.e.,  $\epsilon(\gamma) \propto \gamma^2$ . On the boundary of the Brillouin zone, the function  $\epsilon(\gamma)$  has a local minimum,  $\epsilon^0(1) = d\epsilon(\gamma)/d\gamma|_{\gamma=1} = 0$ , and the short-wave part of the spectrum (for  $\epsilon \rightarrow 1$ ) exhibits a non-analytic dependence on  $\gamma$ . As Eqs. (3.22) and (3.23) show, the non-analyticity of  $\epsilon(\gamma)$  at  $\epsilon \rightarrow 1$  is caused by the long-range action of the Coulomb interaction.

The dependence of  $\epsilon(\gamma)$  on  $\gamma$  for different values of  $r$  is shown in Fig. 5. As is seen from this figure,  $\epsilon(\gamma)$  is a non-monotonic function of  $\gamma$  which has a unique maximum on the interval  $(1/2; 1)$ . As  $r$  decreases, its local minimum  $\epsilon(1)$  also decreases, and  $\epsilon(1) > 0$  for  $r > r_1$ . Since the inequality  $\epsilon(1) > 0$  implies that  $\epsilon(\gamma) > 0$  for all  $\gamma \in [0, 1]$ , the condition  $r > r_1$  is the stability criterion of a chain with respect to small displacements of particles from their equilibrium positions. Note that though for  $r_1 < r < r_2$  a chain with infinite period is energy-wise preferred compared to a chain with equidistant particles, it is the latter which is stable. If  $r < r_1$ , then a region exists on the interval  $[0; 1]$  where  $\epsilon^2(\gamma) < 0$ , and therefore such chains are unstable. Finally, if  $r = r_1$ , then  $\epsilon(1) = 0$ ,  $\epsilon^0(1) = -1$ , and the asymptotic of  $\epsilon(\gamma)$  as  $\epsilon \rightarrow 1$  has the form

$$\epsilon(\gamma) \approx \frac{\epsilon_{\text{cut}}}{\epsilon_1(r_1)} \frac{2}{3} (1-\gamma)^2 j \ln(1-\gamma); \quad (3.26)$$

In this case the optical displacements of particles do not lead to an appearance of restoring forces. Therefore, to determine if a chain is stable or not at  $r = r_1$ , it is necessary to go beyond the harmonic approximation.

### C. Forced vibrations

We briefly consider also the forced vibrations of particles under the action of the longitudinal alternating electric field  $E(t) = E \cos \omega_0 t$ , where  $E$  and  $\omega_0$  are the amplitude and frequency of the alternating field, respectively. In such field the electric force  $qE \cos \omega_0 t$  acts on positive charges, and the force  $-qE \cos \omega_0 t$  acts on negative charges. If the friction force  $\gamma \dot{x}_n$  ( $\gamma$  is a damping coefficient) also acts on each particle, then the forced vibrations of particles are described by the following system of motion equations:

$$\begin{aligned} \ddot{x}_n^+ + 2\gamma \dot{x}_n^+ + \sum_m^X B_{2(n-m)+1} (x_n^+ - x_m^+) \\ + \sum_{m \neq n}^X B_{2(n-m)} (x_n^+ - x_m^+) = A \cos \omega_0 t; \\ \ddot{x}_n^- + 2\gamma \dot{x}_n^- + \sum_m^X B_{2(n-m)+1} (x_n^- - x_m^-) \\ + \sum_{m \neq n}^X B_{2(n-m)} (x_n^- - x_m^-) = -A \cos \omega_0 t \end{aligned} \quad (3.27)$$

( $\gamma = \gamma_0 M$ ,  $A = qE/M$ ). Symmetry of these equations suggests that we seek their solution in the form  $x_n^+ = x(t)$  and  $x_n^- = -x(t)$ . Taking into account that  $\sum_{p=1}^P B_{2p-1} = 0$ , in this case Eqs. (3.27) are reduced to the ordinary differential equation for a forced harmonic oscillator

$$\ddot{x}(t) + 2\gamma \dot{x}(t) + \omega_0^2(1) x(t) = A \cos \omega_0 t \quad (3.28)$$

Its steady-state solution, which describes the forced vibrations of particles, has the form  $x(t) = x_0 \cos(\omega_0 t - \varphi)$ , where

$$x_0 = \frac{A}{[\omega_0^2(1) - \omega_0^2]^2 + 4\gamma^2 \omega_0^2} \quad (3.29)$$

is the amplitude displacement of particles from their equilibrium positions, and  $\tan \varphi = 2\gamma \omega_0 / [\omega_0^2(1) - \omega_0^2]$ . According to Eq. (3.29), if  $\omega_0(1) > \omega_0$ , then the dependence of  $x_0$  on the driving frequency  $\omega_0$  has resonance character;  $\omega_0 = [\omega_0^2(1) - 2\gamma^2]^{1/2}$  is the resonance frequency, and  $x_0 = A/2[\omega_0^2(1) - 2\gamma^2]$  is the resonance amplitude.

#### IV . DETERMINISTIC TRANSPORT

In this section we study the transport properties of a chain whose particles are driven by the longitudinal alternating electric field  $E(t) = E h(t)$ ,

$$h(t) = \begin{cases} 1; & 2k < t < 2k + 1; \\ 0; & 2k + 1 < t < 2k + 2; \end{cases} \quad (4.1)$$

and interact with the sawtooth (ratchet) potential  $V(x)$  that induces the force field  $f(x) = -dV(x)/dx = -g(x)$ , where

$$g(x) = \begin{cases} 0; & md + c < x < md + d; \\ 1; & md < x < md + c; \end{cases} \quad (4.2)$$

Here  $k$  and  $m$  run over all integers,  $\tau$  is the half-period of  $E(t)$ ,  $f_0 = j m \ln f(x) j$  and  $d$  is the period of  $V(x)$  (see Fig. 6). The dynamics of particles in such a chain is described by the system of equations (3.27) in which the right-hand sides  $A \cos \omega_0 t$  and  $-A \cos \omega_0 t$  must be replaced by  $A h(t) + R g(x_n^+)$  and  $-A h(t) + R g(x_n^-)$  ( $R = f_0/M$ ), respectively. To simplify the problem, we assume that the condition  $a = Ld$  ( $L$  is a natural number) holds, so that the equilibrium positions of all particles coincide with the minima of  $V(x)$ . In this case, under the action of the electric field  $E(t)$  all positively charged particles and all negatively charged particles move identically, i.e.,  $x_n^+ = w(t)$  and  $x_n^- = u(t)$ , and we obtain

$$\begin{aligned} w + 2\pi \omega_0^{-1} (1) (w - u) &= A h(t) + R g(w); \\ u + 2\pi \omega_0^{-1} (1) (u - w) &= -A h(t) + R g(u); \end{aligned} \quad (4.3)$$

Due to the periodicity of  $h(t)$  and  $g(x)$ , the steady-state solution of Eqs. (4.3) satisfies the conditions

$$w(t + 2\pi \omega_0^{-1}) = u(t) + Kd; \quad u(t + 2\pi \omega_0^{-1}) = w(t) + Kd; \quad (4.4)$$

where  $K$  is a whole number. It follows from these conditions that

$$w(t + 2\pi \omega_0^{-1}) - w(t) = 2Kd; \quad u(t + 2\pi \omega_0^{-1}) - u(t) = 2Kd; \quad (4.5)$$

i.e., for the period of  $h(t)$  all particles are displaced the same distance  $\Delta = 2Kd$ . This fact permits us to define the average velocity of a chain as

$$v = \frac{w(2\pi \omega_0^{-1}) - w(0)}{2\pi \omega_0^{-1}} = K \frac{d}{\tau}; \quad (4.6)$$

To find  $v$ , we need to calculate  $\dot{x}$  and in the general case solve Eqs. (4.3). These equations have the important feature that we need to consider only the ratchet potentials with  $c > d=2$ . Indeed, the transformation  $c \rightarrow d - c$  leads to the inverted potential  $V_{in}(x) = V(-x)$  which gives rise to the reduced force field  $g_{in}(x) = -g(-x)$ . According to Eqs. (4.3), in such a potential the displacements  $w_{in}(t)$  and  $u_{in}(t)$  of positive and negative charges from their equilibrium positions are governed by the equations of motion

$$\begin{aligned} \ddot{w}_{in} + \frac{1}{2} \ddot{w}_{in} + \frac{1}{2} (1) (\dot{w}_{in} - \dot{u}_{in}) &= A h(t) - R g(-w_{in}); \\ \ddot{u}_{in} + \frac{1}{2} \ddot{u}_{in} + \frac{1}{2} (1) (\dot{u}_{in} - \dot{w}_{in}) &= -A h(t) - R g(u_{in}); \end{aligned} \quad (4.7)$$

Comparing these equations with Eq. (4.3), we obtain  $w_{in}(t) = -u(t)$  and  $u_{in}(t) = -w(t)$ . This means that if for  $c > d=2$  the solution of equations (4.3) is known, then their solution for  $c < d=2$  is known too. Using Eqs. (4.5) and (4.6), we find the chain velocity in the inverted potential  $v_{in} = -v$ , i.e., in the potentials  $V_{in}(x)$  and  $V(x)$  a chain drifts in the opposite directions with the same velocity. If  $c = d=2$  then  $V_{in}(x) = V(x)$ , both conditions  $v_{in} = -v$  and  $v_{in} = v$  hold, and  $v_{in} = v = 0$ . Hence, to study the transport properties of a chain, we indeed can consider only those ratchet potentials for which  $c > d=2$ . In this case, a chain can drift only along the positive direction of the  $x$  axis and so  $K = 0$ .

Next we study the deterministic transport of a chain in the overdamped limit where Eqs. (4.3) are reduced to

$$\begin{aligned} 2 \ddot{w} + \frac{1}{2} \ddot{w} + \frac{1}{2} (1) (\dot{w} - \dot{u}) &= A h(t) + R g(w); \\ 2 \ddot{u} + \frac{1}{2} \ddot{u} + \frac{1}{2} (1) (\dot{u} - \dot{w}) &= -A h(t) + R g(u); \end{aligned} \quad (4.8)$$

Since our aim is to find the chain velocity  $v$ , we need to solve Eqs. (4.8) only for  $t \in (0; 2\pi)$ . To simplify the problem further, we assume that the conditions  $L = K = 1$  and  $1 < A=R < c=(d-c)$  hold. The former implies that the interparticle distance  $a$  is much larger than the chain displacement  $\delta$ , which, in turn, is much larger than the potential period  $d$ . The latter shows that for  $h(t) > 0$  ( $h(t) < 0$ ) positive (negative) charges can overcome the potential barriers of  $V(x)$  and they move along the positive direction of the  $x$  axis, whereas negative (positive) charges stay in the minima of  $V(x)$ . After  $h(t)$  changes sign, the particles at rest begin to move, and the moving particles move to the appropriate minima of  $V(x)$  and stop there. Since the maximal distance that the moving particles travel after  $h(t)$  changes sign does not exceed  $c$ , i.e., this distance is much less than  $a$ , we assume that the moving

particles reach the corresponding minimum of  $V(x)$  immediately as  $h(t)$  changes sign. Within this approximation, the dynamics of positive and negative charges is described separately, and Eqs. (4.8) yield  $u(t) = 0$  and

$$2 \frac{dw}{dt} + \frac{1}{2} w^2 = A + Rg(w) \quad (4.9)$$

$w(0) = -2, w(t) = -2$  for  $t \in (0; \tau)$ , and  $w(t) = 2$  and

$$2 \frac{du}{dt} + \frac{1}{2} u^2 = A + Rg(u) \quad (4.10)$$

$u(t) = 0, u(t) = 2$  for  $t \in (\tau; 2\tau)$ .

To calculate  $\tau$  we can use either Eq. (4.9) or Eq. (4.10). We take Eq. (4.9) as our starting point and write it in the form

$$\frac{dw}{dt} = \begin{cases} \frac{1}{2} (2 - w)^2 & \text{for } 0 \leq t < \tau \\ \frac{1}{2} (w - 2)^2 & \text{for } \tau \leq t \leq 2\tau \end{cases} \quad (4.11)$$

where  $\frac{1}{2} = \frac{1}{2} (1 - \frac{c}{d})$ ,  $\frac{1}{2} = \frac{1}{2} (A - R) = \frac{1}{2}$ ,  $\frac{2}{2} = \frac{1}{2} (A + R) = \frac{1}{2} (d - c) = \frac{1}{2}$ , and  $m = K; K + 1; \dots; K - 1$ . The solution of Eq. (4.11) depends on  $\tau$  and actually defines  $\tau$ , because  $w(0) = -2$  and  $w(t) = -2$ . We use this feature to derive the following equation (see Appendix C)

$$e^{t_{2K}} = \frac{\frac{2 - \frac{1}{2}}{2d} + 1}{\frac{2 - \frac{1}{2}}{2d} + 1} \cdot \frac{\frac{2 - \frac{1}{2}}{2d} + \frac{c}{d} + 1}{\frac{2 - \frac{1}{2}}{2d} + \frac{c}{d} + 1} \cdot \frac{\frac{2 - \frac{2}{2}}{2d} + \frac{c}{d} + 1}{\frac{2 - \frac{2}{2}}{2d} + \frac{c}{d} + 1} \cdot \frac{\frac{2 - \frac{2}{2}}{2d} + \frac{c}{d} + 1}{\frac{2 - \frac{2}{2}}{2d} + \frac{c}{d} + 1} \quad (4.12)$$

for the instant  $t = t_{2K}$  that corresponds to a displacement of positive charges of  $x = 2Kd$ . An increase of  $\tau$  leads to the step-wise increase of  $t_{2K}$  and  $K$ , and from Eq. (4.12) we obtain

$$m_{\max} K = m \ln \left( \frac{1 + (d - c)}{d} \right)^{\frac{1}{2}} ; \frac{2}{d} ; \quad (4.13)$$

where  $[x]$  denotes the integer part of  $x$ .



Replacing in Eq. (4.12)  $t_{2K}$  by  $\tau$ , we obtain the approximate equation that defines  $\tau$  as a function of  $\tau$ . For  $\tau \gg 1$  it yields  $K \approx \max K$ , and so in this case the chain velocity  $v$  is given by

$$v = \frac{2d}{\tau} \max K : \quad (4.14)$$

It follows from this formula that  $v \rightarrow 0$  as  $\tau \rightarrow \infty$  and, according to Eq. (4.13),  $v$  depends weakly on  $d$  and  $v \propto \tau^{-2}$  (1). The last proportionality shows that the chain velocity strongly depends on the interparticle interactions, and  $v$  is a decreasing function of  $r$ . [Note that the permissible values of  $r$  are not too close to  $r_1$ , since our assumptions imply that  $\max K \ll L$  holds.]

If  $K \approx \max K$ , then, using the asymptotic formula<sup>40</sup>

$$\frac{(z+x)}{(z+y)} = z^{x-y} \left( 1 + \frac{1}{2z} (x-y)(x+y-1) \right) \quad (4.15)$$

( $z \gg 1$ ), we reduce the equation for  $\tau$  to the form

$$e^{-\tau} = 1 + \frac{c}{d_1} + \frac{1}{d_2} \left( 1 - \frac{c}{d} \right) : \quad (4.16)$$

Solving Eq. (4.16) with respect to  $\tau$  and taking into account that in this case  $\tau \gg 1$ , we derive the formula

$$v = \frac{1}{\tau} \frac{2d}{(d_1 - c) + d_2 c} ; \quad (4.17)$$

which shows that for  $1 \ll K \approx \max K$  the chain velocity does not depend on the chain characteristics. Since  $1 < A=R < c=(d-c)$ , for a given ratio  $A=R = \eta$  ( $\eta > 1$ ) the parameter  $c=d$  must belong to the interval  $(\eta/( \eta + 1); 1)$ . On this interval,  $v$  is a decreasing function of  $c=d$ , and according to Eq. (4.17)  $v \rightarrow 1/2 = (\eta + 1)$  as  $c=d \rightarrow \eta/( \eta + 1)$ , and  $v \rightarrow 1$  as  $c=d \rightarrow 1$ .

## V. CONCLUSIONS

We have studied the ground state, vibration properties and deterministic transport of a chain of charged particles interacting via both the Coulomb interaction and a repulsive interaction, which is characterized by the power  $r$ . We have shown that a chain with equidistant particles has minimum energy for  $r > r_2 \approx 2.799$ . For a such chain, we have calculated its energy and period as functions of  $r$ , derived and analyzed in detail its phonon

spectrum, and established that it is stable with respect to small displacements of particles from their equilibrium positions only if  $r > r_1$  [2:779].

We have studied also the deterministic transport of a chain in the case when the particles are acted upon by both a longitudinal alternating electric field and a force field generated by a ratchet potential. We have derived the set of equations that describes the chain transport, analyzed the general properties of its solution, and calculated the chain velocity in the overdamped limit. If the period of the electric field is large enough, then the chain velocity strongly depends on the interparticle interactions. Otherwise, i.e., if the period of the electric field is small enough, the chain velocity does not depend on them.

#### ACKNOWLEDGMENT

We are grateful to Werner Horsthemke for a critical reading of the manuscript and his valuable comments.

#### APPENDIX A: ANOTHER FORM FOR $U_0(\cdot; \cdot)$

Using the representation<sup>40</sup>

$$\psi(z) = \lim_{k \rightarrow \infty} \left[ \ln k - \frac{1}{z} \sum_{n=1}^k \frac{1}{n+z} \right] \quad (\text{A } 1)$$

for  $\psi(z) = d \ln \Gamma(z) / dz$  ( $\Gamma(z)$  is the gamma function) and the definition

$$\zeta(r; z) = \sum_{n=0}^{\infty} \frac{1}{(n+z)^r} \quad (\text{A } 2)$$

of the generalized zeta function  $\zeta(r; z)$ , we find

$$\sum_{n=1}^{\infty} \frac{1}{n} + \frac{1}{n+1} - \frac{2}{n} = 2 \quad (1) \quad (2) \quad (3) \quad (\text{A } 3)$$

and

$$\sum_{n=1}^{\infty} \frac{1}{n^2 + j^2} = \zeta(2; j) + \zeta(2; 1-j): \quad (\text{A } 4)$$

Substituting Eqs. (A 3) and (A 4) into Eq. (2.3) and taking into account that  $\zeta(1) = (1-j) + 1 = 1$ ,<sup>40</sup> we obtain the desired representation

$$U_0(\cdot; \cdot) = \frac{1}{2} [2 \zeta(1) - \zeta(2) - \zeta(2)]$$

$$+ \frac{1}{2} \left[ \phi(r) + \phi(r; \infty) + \phi(r; 1) \right] : \quad (\text{A } 5)$$

## APPENDIX B : INTEGRAL REPRESENTATION FOR $\phi_3(\cdot)$ AND $\phi_3(\cdot)$

We proceed from the standard series<sup>39</sup>

$$\sum_{p=1}^{\infty} \frac{\cos(pz)}{p} = -\ln 2 \sin \frac{z}{2} \quad (\text{B } 1)$$

( $0 < z < \pi$ ). Integrating Eq. (B 1) twice, we find

$$\sum_{p=1}^{\infty} \frac{1}{p^3} \cos(pz) = \int_0^z \int_0^x dy \ln 2 \sin \frac{y}{2} ; \quad (\text{B } 2)$$

and using the definition (3.16), we obtain

$$\phi_3(\cdot) = \phi_3(0) + \int_0^z \int_0^x dy \ln 2 \sin \frac{y}{2} : \quad (\text{B } 3)$$

To find the integral representation for  $\phi_3(\cdot)$ , we use the identity

$$\sum_{p=1}^{\infty} \frac{\cos(pz)}{p} = \sum_{p=1}^{\infty} \frac{\cos(2pz)}{2p} + \sum_{p=1}^{\infty} \frac{\cos[(2p-1)z]}{2p-1} \quad (\text{B } 4)$$

and the series (B 1) to obtain

$$\sum_{p=1}^{\infty} \frac{\cos[(2p-1)z]}{2p-1} = \frac{1}{2} \ln \tan \frac{z}{2} : \quad (\text{B } 5)$$

Integrating Eq. (B 5) by the same way as Eq. (B 1) and using the definition (3.17), we find

$$\phi_3(\cdot) = \frac{7}{8} \phi_3(0) + \frac{1}{2} \int_0^z \int_0^x dy \ln \tan \frac{y}{2} : \quad (\text{B } 6)$$

## APPENDIX C : DERIVATION OF EQUATION (4.12)

Let  $t_k$  ( $k = 0; 1; \dots; 2K$ ,  $t_0 = 0$ ,  $t_{2K} = \pi$ ) and  $t_k$  ( $k = 0; 1; \dots; 2K-1$ ,  $t_k < t_k < t_{k+1}$ ) be the times at which the positive charges reach the minima and maxima of  $V(x)$  during the half-period, i.e.,  $w(t_k) = 2 + kd$  and  $w(t_k) = 2 + kd + c$ . Solving Eq. (4.11), we find

$$w(t) = \frac{1}{2} \left( \frac{1}{2} + \frac{1}{2} \right) kd e^{-(t-t_k)} \quad (\text{C } 1)$$

for  $t \in (t_k; t_{k+1})$ , and

$$w(t) = \frac{1}{2} \left( \frac{1}{c} + \frac{1}{d} \right) e^{-\frac{1}{2} \left( \frac{1}{c} + \frac{1}{d} \right) (t - t_k)} \quad (C2)$$

for  $t \in (t_k; t_{k+1})$ . At  $t = t_k$ , Eq. (C1) leads to

$$e^{-\frac{1}{2} \left( \frac{1}{c} + \frac{1}{d} \right) (t_k - t_k)} = \frac{1}{\frac{1}{c} + \frac{1}{d}}; \quad (C3)$$

and Eq. (C2) at  $t = t_{k+1}$  yields

$$e^{-\frac{1}{2} \left( \frac{1}{c} + \frac{1}{d} \right) (t_{k+1} - t_k)} = \frac{1}{\frac{1}{c} + \frac{1}{d}}. \quad (C4)$$

Multiplying Eqs. (C3) and (C4), substituting the result

$$e^{-\frac{1}{2} \left( \frac{1}{c} + \frac{1}{d} \right) (t_{k+1} - t_k)} = \frac{1}{\frac{1}{c} + \frac{1}{d}} \quad (C5)$$

into the identity  $e^{-\frac{1}{2} \left( \frac{1}{c} + \frac{1}{d} \right) (t_{k+1} - t_k)} = \sum_{k=0}^{2K-1} e^{-\frac{1}{2} \left( \frac{1}{c} + \frac{1}{d} \right) (t_{k+1} - t_k)}$ , and using the formula  $\sum_{k=0}^{2K-1} (x - k) = (x) - (x - 2K)$  and condition  $x = 2Kd$ , we obtain Eq. (4.12).

---

Electronic address: denisov@ssu.sumy.ua

- <sup>1</sup> J. Ford, Phys. Rep. 213, 271 (1992).
- <sup>2</sup> J. Szeftel and P. Laurent, Phys. Rev. E 57, 1134 (1998).
- <sup>3</sup> R. Reigada, A. Samiñento, and K. Lindenberg, Phys. Rev. E 64, 066608 (2001).
- <sup>4</sup> S. Flach and C. R. Willis, Phys. Rep. 295, 181 (1998).
- <sup>5</sup> D. Henning and G. P. Tsironis, Phys. Rep. 307, 333 (1999).
- <sup>6</sup> Y. Kuramoto, Chemical Oscillations, Waves, and Turbulence (Springer-Verlag, Berlin, 1984).
- <sup>7</sup> C. Van den Broeck, J. M. R. Parrondo, and R. Toral, Phys. Rev. Lett. 73, 3395 (1994).
- <sup>8</sup> J. H. Li and Z. Q. Huang, Phys. Rev. E 53, 3315 (1996).
- <sup>9</sup> J. F. Lindner, B. K. Meadows, W. L. Ditto, M. E. Inchiosa, and A. R. Bulsara, Phys. Rev. Lett. 75, 3 (1995).
- <sup>10</sup> S. K. Han, W. S. Kim, and H. Kook, Phys. Rev. E 58, 2325 (1998).
- <sup>11</sup> A. Pikovsky, M. Rosenblum, and J. Kurths, Synchronization: A Universal Concept in Nonlinear Science (Cambridge University Press, Cambridge, 2001).
- <sup>12</sup> H. Haken, Brain Dynamics: Synchronization and Activity Patterns in Pulse-Coupled Neural Nets with Delays and Noise (Springer-Verlag, Berlin, 2002).
- <sup>13</sup> D. H. E. Dubin, Phys. Rev. E 55, 4017 (1997).
- <sup>14</sup> D. H. E. Dubin and T. M. O'Neil, Rev. Mod. Phys. 71, 87 (1999).
- <sup>15</sup> G. Morigi and J. Eschner, Phys. Rev. A 64, 063407 (2001).
- <sup>16</sup> K. Hirahara, K. Suenaga, S. Bandow, H. Kato, T. Okazaki, H. Shinohara, and S. Tjima, Phys. Rev. Lett. 85, 5384 (2000).
- <sup>17</sup> R. R. Meyer, J. Sloan, R. E. Dunin-Borkowski, A. I. Kirkland, M. C. Novotny, S. R. Bailey, J. L. Hutchison, and M. L. H. Green, Science 289, 1324 (2000).
- <sup>18</sup> M. O. Magnasco, Phys. Rev. Lett. 71, 1477 (1993).
- <sup>19</sup> R. D. Astumian and M. Bier, Phys. Rev. Lett. 72, 1766 (1994).
- <sup>20</sup> C. R. Doering, W. Horsthemke, and J. Riordan, Phys. Rev. Lett. 72, 2984 (1994).
- <sup>21</sup> C. R. Doering, Nuovo Cimento 17D, 685 (1995).
- <sup>22</sup> R. Bartussek, P. Reimann, and P. Hanggi, Phys. Rev. Lett. 76, 1166 (1996).
- <sup>23</sup> R. D. Astumian, Science 276, 917 (1997).

- <sup>24</sup> J.M .R .Parrondo, Phys.Rev.E 57, 7297 (1998).
- <sup>25</sup> M .N .Popescu, C .M .A rizm endi, A .L .Salas-Brito, and F .Fam ily, Phys.Rev.Lett. 85, 3321 (2000).
- <sup>26</sup> L.C ao and D .-j.W u, Phys.Lett.A 291, 371 (2001).
- <sup>27</sup> P .Reim ann, Phys.Rep. 361, 57 (2002).
- <sup>28</sup> J.H oward, Nature (London) 391, 240 (1998).
- <sup>29</sup> R .D .Vale and R .A .M illigan, Science 288, 88 (2000).
- <sup>30</sup> J.Rousselet, L .Salom e, A .A jdari, and J.Prost, Nature (London) 370, 446 (1994).
- <sup>31</sup> I.D erenyi, C .Lee, and A .-L .Barabasi, Phys.Rev.Lett. 80, 1473 (1998).
- <sup>32</sup> C .-S.Lee, B .Janko, I.D erenyi, and A .-L .Barabasi, Nature (London) 400, 337 (1999).
- <sup>33</sup> A .Sarm iento and H .Larralde, Phys.Rev.E 59, 4878 (1999).
- <sup>34</sup> J.L.M ateos, Phys.Rev.Lett. 84, 258 (2000).
- <sup>35</sup> Z.C sahok, F .Fam ily, and T .Vicsek, Phys.Rev.E 55, 5179 (1997).
- <sup>36</sup> J.Buceta, J.M .Parrondo, C .Van den Broeck, and F .J.de la Rubia, Phys.Rev.E 61, 6287 (2000).
- <sup>37</sup> C .J.O lson, C .Reichhardt, B .Janko, and F .Nori, Phys.Rev.Lett. 87, 177002 (2001).
- <sup>38</sup> N.W .Ashcroft and N.D .M ermin, Solid State Physics (Holt, Rinehart & W inston, New York, 1976).
- <sup>39</sup> A .P.P nudnikov, Yu.A .B rychkov, and O .I.M arichev, Integrals and Series (G ordon & B reach, New York, 1989), Vol. 1.
- <sup>40</sup> H .Bateman and A .Erdelyi, Higher Transcendental Functions (M cG raw-H ill, New York, 1953), Vol. 1.
- <sup>41</sup> P.Lankaster, Theory of Matrices (A cadem ic P ress, New York, 1969).




FIG .1: Part of the equilibrium chain of charged particles.




FIG .2: Plots of  $\beta_1(r)$  and  $\beta_2(r)$ ; for the third solution  $\beta_3(r) = 1$ . The function  $\beta_2(r)$  increases from  $\beta_2(r_1) = \beta_1(r_1) = 4.959$  to  $\beta_2(3) = 1$ .




FIG . 3: Plots of  $\gamma_s(r)$ . All functions  $\gamma_s(r)$  decrease monotonically, and  $\gamma_1(r_1) = \gamma_2(r_1) = 2.420$ ,  $\gamma_2(3) = \gamma_3(3) = \frac{1}{3}$ .




FIG . 4: Plots of  $U_1(r)$  and  $U_3(r)$  [figure (a)], and  $U_1(r)$  and  $U_2(r)$  [figure (b)]. The functions  $U_3(r)$  and  $U_2(r)$  are very close to each other [ $U_3(r) < U_2(r)$  for  $r_1 < r < 3$ ,  $U_3(3) = U_2(3) = \frac{1}{3}$ , and  $U_2(r_1) - U_3(r_1) = 0.001$ ], therefore the plot of  $U_3(r)$  in figure (b) is not shown.






FIG . 5: Dependence of the phonon spectrum  $\tilde{\omega}(\kappa)$  on  $\kappa$  for  $r = 4$  (curve 1),  $r = 3$  (curve 2), and  $r = r_1$  (curve 3). Since for  $r = r_1$ , the function  $\tilde{\omega}^0(\kappa)$  goes to infinity very slowly [as  $\sqrt{\ln(1-\kappa)}$ ] as  $\kappa \rightarrow 1$ , this feature of the spectrum is not represented on the chosen scale of the  $\kappa$  axis.




FIG . 6: Illustrative plots of the reduced alternating electric field  $h(t)$  [figure (a)], the ratchet potential  $V(x)$  [figure (b)], and the reduced force field  $g(x)$  [figure (c)].