

Comment on "The nature of slow dynamics in a minimal model of frustration limited domains" by P.L. Geissler and D.R. Reichman: In a recent paper Geissler and Reichman<sup>1</sup> investigated a model with the Hamiltonian

$$H = \int dr \frac{1}{2} \dot{r}^2 + k_0^2 r^2 + k_0^2 \phi^2 + \frac{1}{4!} \phi^4 \quad (1)$$

with scalar field  $\phi(r)$  in three dimensions using Monte Carlo simulations.<sup>2</sup> The parameters of the model are the typical wave number,  $k_0$ , for modulations of  $\phi$ , the "segregation strength",  $\phi$ , between positive and negative  $\phi$ , and the interaction strength,  $\phi^4$ . Many years ago by Brazovskii<sup>3</sup> argued that this model may undergo a fluctuation induced first order transition into a lamellar state.

We recently argued that systems described by Eq.1 may also undergo a self-generated glass transition, i.e. can become non-ergodic without the presence of quenched disorder<sup>4</sup>. This result was based on our earlier work<sup>5,6</sup> which demonstrated such glassiness in a related model discussed in the context of frustration limited domain formation by Kivelson et al.<sup>7</sup>. The qualitative features of that theory were confirmed in mode-coupling calculations and Monte Carlo simulations of the corresponding lattice model by Grousson et al.<sup>8,9</sup>. The theory is based on a replica formalism and two key assumptions were made in Refs.4,5,6: i) Glassiness can be studied within the mean-field replica formalism with one step replica symmetry breaking which was shown to be marginally stable. ii) The self-consistent screening approximation (SCSA), including de Gennes narrowing, for the summation of the perturbation series in  $\phi$  can be used. Recently we have been able to avoid this second assumption and found a complete solution of the mean-field replica equations by generalizing the dynamical mean-field approach (DMFT) of correlated electron models<sup>10</sup> to the case of glassy systems.<sup>11</sup> Given that our theory still relies on the applicability of the mean-field replica formalism, computer simulations like those of Refs.1,8 remain crucial for a final judgment on the existence of self-generated glassiness in uniformly frustrated systems of this type.

Geissler and Reichman performed Monte-Carlo (MC) calculations and compared the results with two analytical approaches, one, a Hartree theory and the other, a mode coupling theory performed self-consistently up to second order in  $\phi$ . For  $\phi = 1$ ,  $k_0 = 0.5$  and between 0.14 and 0, good agreement between the MC and Hartree approaches was obtained, whereas the primitive mode-coupling theory yields glassy dynamics already for

0.1, which was not seen in the Monte Carlo simulations. Thus, the Monte Carlo simulations could not confirm the glassy dynamics found in mode coupling calculations, but rather gave only fast, "liquid-like" relaxations. Geissler and Reichman concluded that Eq.1 likely does not exhibit glassy dynamics.

In Fig.1 we compare our results for the dynamic transition between the liquid and glass state obtained within




FIG. 1: Phase segregation strength  $\phi$  at the dynamic transition between the liquid and glass state as function of the coupling constant  $\lambda$ , obtained by using 2<sup>nd</sup>-order perturbation theory, the self-consistent screening approximation (SCSA) of Refs.4,5,6 and the dynamical mean-field theory (DMFT) of Ref.11. The diamonds indicate the parameters of the Monte Carlo calculations by Geissler and Reichman where only liquid-like behavior was found. The SCSA and DMFT calculations are consistent with the Monte Carlo results whereas the 2<sup>nd</sup>-order calculation (similar to the mode coupling theory of Ref.1) strongly overestimates glassiness.

the replica formalism and solved by using three different techniques: a self-consistent second order theory (similar to the mode coupling theory of Ref.1), the SCSA used in Refs.4,5,6 and the complete solution of the replica mean-field problem (DMFT) of Ref.11. In the second order calculation we determined the liquid state correlation function within Hartree theory, which seems to be closest to the procedure of Ref.1, who used Monte Carlo results for the liquid structure factor and found that it is similar to the one obtained within the Hartree approach. The diamonds are the results of Geissler and Reichman (Figs.5 and 6 of Ref.1) where MC simulations give liquid-like behavior. It is obvious from this plot that the results by Geissler and Reichman indeed demonstrate that a second order mode coupling theory gives a poor approximation for the location of the glass transition. However, their results cannot be used to rule out the glass transition we proposed. Within the complete solution of the replica mean-field theory this transition should occur only for  $\lambda$ -values smaller than 0.2. In fact Fig. 2 of Geissler and Reichman is not inconsistent with a very slow relaxation rate for  $\lambda < 0.2$ . Clearly, just as for any phase transition, even the best solution of the mean-field approach might yield a phase boundary which is different from what the essentially exact Monte Carlo simulation finds. The key aspect of our theory was that

even a simple mean field theory leads to a proliferation of metastable states and to an entropy crisis. Thus, self-generated randomness emerges according to the random first order transition scenario.<sup>12</sup>

That the parameters studied in Ref.1 are outside the region relevant for glassiness is consistent with their findings that no indication for the first order Brazovskii transition was detected either. We argued in Ref.4 that the glass transition occurs due to the same type of fluctuations and for similar parameters to the fluctuation induced first order transition proposed by Brazovskii<sup>3</sup>. The absence of the glass transition could therefore most convincingly be demonstrated by showing the impossibility of achieving cooling rates that exceed nucleation rates.

In summary, the recent Monte Carlo calculations by Geissler and Reichman<sup>1</sup> are not conclusive in ruling out the self-generated glass transition proposed in Ref.5. Nevertheless, if performed for parameters relevant for glassy behavior Monte Carlo calculations like theirs

would be very useful in judging whether Eq.1 can be considered as a minimal model for glassiness or whether additional ingredients are needed. In addition they offer a powerful tool to study glassy dynamics beyond the limitations of replica mean field theory.

This work has been supported by Research Corporation and Ames Laboratory, operated for the U.S. Department of Energy by Iowa State University under Contract No. W-7405-Eng-82 (J.S. and S.W.) and NSF (P.G.W.), Grant No. CHE-9530680.

Jörg Schmalian, Sangwook Wu

Department of Physics and Astronomy and

Ames Laboratory, Iowa State University, Ames, IA 50011

Peter G. Wolynes,

Department of Chemistry and Biochemistry,

University of California at San Diego, La Jolla, CA 92093

<sup>1</sup> P. L. Geissler and D. R. Reichman, cond-mat/0304254

<sup>2</sup> In Ref.[1] there is no factor  $\frac{1}{2}$  in front of the term quadratic in  $\phi$ . Since this factor is needed to derive Eqs. 7 and 9 from Eq.2 of Ref.[1], we assume that the missing pre-factor is a misprint of Ref.[1].

<sup>3</sup> S. A. Brazovskii, Sov. Phys. JETP 41, 85 (1975).

<sup>4</sup> S.W. Wu, H. Westfahl Jr., J. Schmalian and P. G. Wolynes, Chem. Phys. Lett. 359, 1 (2002).

<sup>5</sup> J. Schmalian and P. G. Wolynes, Phys. Rev. Lett. 85, 836 (2000).

<sup>6</sup> H. Westfahl Jr., J. Schmalian and P. G. Wolynes, Phys. Rev. B 64, 174203 (2001).

<sup>7</sup> D. Kivelson, S. A. Kivelson, X. L. Zhao, Z. Nussinov, and G. Tarjus, Physica A 219, 27 (1995).

<sup>8</sup> M. G. Rousson, G. Tarjus, and P. Viot, Phys. Rev. Lett. 86, 3455 (2001); *ibid* Phys. Rev. E 65, 065103(R) (2002).

<sup>9</sup> M. G. Rousson, V. K. Rakoviack, G. Tarjus, P. Viot, Phys. Rev. E 66, 026126 (2002).

<sup>10</sup> A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. 68, 13-125 (1996).

<sup>11</sup> S. Wu, J. Schmalian, G. Kotliar and P. G. Wolynes, cond-mat/0305404.

<sup>12</sup> T. R. Kirkpatrick and P. G. Wolynes, Phys. Rev. A 35, 3072 (1987); *ibid* Phys. Rev. B 36, 8552 (1987), T. R. Kirkpatrick and D. Thirumalai, and P. G. Wolynes, Phys. Rev. A 40, 1045 (1989), T. R. Kirkpatrick and D. Thirumalai, Phys. Rev. Lett. 58, 2091 (1987).