

Preliminary Preparations: Scattering Problem of Double-layered Sphere Containing Left-handed Media

Jian-Qi Shen^y, Yi Jin & Long Chen

Centre for Optical and Electromagnetic Research, State Key Laboratory of Modern Optical Instrumentation
Zhejiang University, Hangzhou 310027, P. R. China
(December 25, 2021)

In this paper, we present Mie coefficients of double-layered sphere and consider the scattering problem, including the topics on field distribution, electromagnetic cross section, extinction spectra as well as some potential peculiar properties arising from the presence of left-handed media. The formulation presented here can be easily generalized to cases of multiple-layered spheres.

Keywords: Mie coefficients, double-layered sphere, scattering problem, left-handed media
PACS: 78.35.+c

I. INTRODUCTION

More recently, a kind of artificial composite media (the so-called left-handed media) having a frequency band where the effective permittivity (ϵ) and the effective permeability (μ) are simultaneously negative attracts attention of many researchers in various fields such as materials science, condensed matter physics, optics and electromagnetism [1-5]. Veselago first considered this peculiar medium and showed that it possesses a negative index of refraction [6]. It follows from the Maxwell's equations that in this medium the Poynting vector and wave vector of electromagnetic wave would be antiparallel, i. e., the vector \mathbf{k} , the electric field \mathbf{E} and the magnetic field \mathbf{H} form a left-handed system; thus Veselago referred to such materials as "left-handed", and correspondingly, the ordinary medium in which \mathbf{k} , \mathbf{E} and \mathbf{H} form a right-handed system may be termed the "right-handed" medium. Other authors call this class of materials "negative-index media (NIM)", "double negative media (DNM)" [5] and Veselago's media. It is readily verified that in such media having both ϵ and μ negative, there exist a number of peculiar electromagnetic properties, for instance, many dramatically different propagation characteristics stem from the sign change of the group velocity, including reversals of both the Doppler shift and the Cherenkov radiation, anomalous refraction, modified spontaneous emission rates and even reversals of radiation pressure to radiation tension [2]. In experiments, this artificial negative electric permittivity media may be obtained by using the array of long metallic wires (ALMWS), which simulates the plasma behavior at microwave frequencies, and the artificial negative magnetic permeability media may be built up by using small resonant metallic particles, e. g., the split ring resonators (SRRs), with very high magnetic polarizability [3,7,9].

The extinction properties of a sphere (single-layered) with negative permittivity and permeability is investigated by Ruppin. Since recently Wang and Asher developed a novel method to fabricate nanocomposite SiO_2 spheres (100 nm) containing homogeneously dispersed Ag quantum dots (2–5 nm) [10], which may have potential applications to design and fabrication of photonic crystals, it is believed that the absorption and transmittance of double-layered sphere deserves consideration. In this paper, both the Mie coefficients and the electromagnetic field distributions in the two-layered sphere containing left-handed media irradiated by a planar electromagnetic wave are presented.

II. METHODS

It is well known that in the absence of electromagnetic sources, the characteristic vectors such as \mathbf{E} , \mathbf{B} , \mathbf{D} , \mathbf{H} and Hertz vector in the isotropic homogeneous media agree with the same differential equation

$$\nabla \times \nabla \times \mathbf{r} - \nabla (\nabla \cdot \mathbf{r}) + k^2 \mathbf{r} = 0: \quad (2.1)$$

This paper will be submitted nowhere else for the publication, just uploaded at the e-print archives.

^yE-mail address: jqshen@coer.zju.edu.cn

The three independent vector solutions to the above equation is [11]

$$L = r \times a; \quad M = r \times a \times r; \quad N = \frac{1}{k} r \times \nabla M \quad (2.2)$$

with a being a constant vector, where the scalar function satisfies $r^2 + k^2 = 0$. It is verified that the vector solutions M , N and L possess the following mathematical properties

$$M \cdot L = a = \frac{1}{k} r \cdot N; \quad L \cdot M = 0; \quad r \cdot L = 0; \quad r \cdot L = r^2 = -k^2; \quad r \cdot M = 0; \quad r \cdot N = 0; \quad (2.3)$$

Set $M = m \exp(-i\omega t)$ and $N = n \exp(-i\omega t)$, the vector wave functions M and N can be expressed in terms of the following spherical vector wave functions [11]

$$\begin{aligned} m_{\theta m n} &= \frac{m}{\sin \theta} z_n(kr) P_n^m(\cos \theta) \sin^m \theta i_2 - z_n(kr) \frac{\partial P_n^m(\cos \theta)}{\partial \theta} \cos^m \theta i_3; \\ n_{\theta m n} &= \frac{n(n+1)}{kr} z_n(kr) P_n^m(\cos \theta) \cos^m \theta i_1 + \frac{1}{kr} \frac{\partial}{\partial r} [r z_n(kr)] \frac{\partial}{\partial \theta} P_n^m(\cos \theta) \sin^m \theta i_2 \\ &\quad - \frac{m}{kr \sin \theta} \frac{\partial}{\partial r} [r z_n(kr) P_n^m(\cos \theta) \sin^m \theta i_3; \end{aligned} \quad (2.4)$$

In what follows we treat the Mie coefficients of double-layered sphere irradiated by a plane wave.

A. Definitions

We consider a double-layered sphere with interior radius a_1 and external radius a_2 having relative permittivity (permability) ϵ_1 (μ_1) and ϵ_2 (μ_2), respectively, placed in a medium having the relative permittivity ϵ_0 and permability μ_0 . Suppose that the double-layered sphere irradiated by the following plane wave with the electric amplitude E_0 along the z -direction of Cartesian coordinate system [11]

$$\begin{aligned} E_i &= a_x E_0 \exp(ik_0 z - i\omega t) = E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} m_{o1n}^{(1)} i_n^{(1)}; \\ H_i &= a_y \frac{k_0}{\omega} E_0 \exp(ik_0 z - i\omega t) = \frac{k_0}{\omega} E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} m_{e1n}^{(1)} + i n_{o1n}^{(1)}; \end{aligned} \quad (2.5)$$

where

$$\begin{aligned} m_{e1n}^{(1)} &= \frac{1}{\sin \theta} j_n(k_0 r) P_n^1(\cos \theta) \cos \theta i_2 - j_n(k_0 r) \frac{\partial P_n^1(\cos \theta)}{\partial \theta} \sin \theta i_3; \\ n_{e1n}^{(1)} &= \frac{n(n+1)}{k_0 r} j_n(k_0 r) P_n^1(\cos \theta) \sin \theta i_1 + \frac{1}{k_0 r} [k_0 r j_n(k_0 r)]^0 \frac{\partial P_n^1(\cos \theta)}{\partial \theta} \cos \theta i_2 - \frac{1}{k_0 r \sin \theta} [k_0 r j_n(k_0 r)]^0 P_n^1(\cos \theta) \sin \theta i_3 \end{aligned} \quad (2.6)$$

with the primes denoting differentiation with respect to their arguments and k_0 being $\frac{\omega}{c}$. In the region $r < a_1$, the wave function is expanded as the following series

$$\begin{aligned} E_t &= E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} a_n^t m_{o1n}^{(1)} - i b_n^t n_{e1n}^{(1)}; \\ H_t &= \frac{k_1}{\omega} E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} b_n^t m_{e1n}^{(1)} + i a_n^t n_{o1n}^{(1)}; \end{aligned} \quad (2.7)$$

Note that here the propagation constant in $m_{o1n}^{(1)}$ and $n_{e1n}^{(1)}$ which have been defined in (2.6) is replaced with k_1 , i.e., $\frac{\omega}{c_1}$. In the region $a_1 < r < a_2$, one may expand the electromagnetic wave amplitude as

$$\begin{aligned} E_m &= E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} a_n^m m_{o1n}^{(1)} - i b_n^m n_{e1n}^{(1)} + a_n^m m_{o1n}^{(3)} - i b_n^m n_{e1n}^{(3)}; \\ H_m &= \frac{k_2}{\omega} E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} b_n^m m_{e1n}^{(1)} + i a_n^m n_{o1n}^{(1)} + b_n^m m_{e1n}^{(3)} + i a_n^m n_{o1n}^{(3)}; \end{aligned} \quad (2.8)$$

Note that here in order to obtain $m_{e1n}^{(3)}$ and $n_{o1n}^{(3)}$, the spherical Bessel functions j_n in the spherical vector function $m_{e1n}^{(1)}$ and $n_{o1n}^{(1)}$ is replaced by the Hankel functions $h_n^{(1)}$, namely, the explicit expressions for $m_{e1n}^{(3)}$ and $n_{o1n}^{(3)}$ can also be obtained from (2.6), so long as we replace the spherical Bessel functions j_n in (2.6) with the Hankel functions $h_n^{(1)}$. Apparently, here the propagation constant in $m_{e1n}^{(1)}$ and $n_{o1n}^{(1)}$ should be replaced with k_2 , i.e., $P \frac{1}{2} \frac{1}{c}$.

In the region $r > a_2$ the reflected wave is written [11]

$$\begin{aligned} E_r &= E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} a_n^r m_{o1n}^{(3)} - i b_n^r n_{e1n}^{(3)} ; \\ H_r &= \frac{k_0}{0!} E_0 \exp(-i\omega t) \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} b_n^r m_{e1n}^{(3)} + i a_n^r n_{o1n}^{(3)} ; \end{aligned} \quad (2.9)$$

where the propagation vector in $m_{e1n}^{(3)}$ and $n_{o1n}^{(3)}$ is replaced by k_0 , i.e., $P \frac{1}{0} \frac{1}{c}$.

B. Boundary conditions

At the boundary $r = a_1$, the boundary condition is given as follows

$$i_1 E_m = i_1 E_t; \quad i_1 H_m = i_1 H_t; \quad (2.10)$$

where the unit vectors of Cartesian coordinate system agree with $i_1 = i_3, i_2 = i_3, i_3 = i_1, i_3 = i_2$.

It follows from $i_1 E_m = i_1 E_t$ that

$$\begin{aligned} a_n^m j_n(N_2 a_1) + a_n^m h_n^{(1)}(N_2 a_1) &= a_n^t j_n(N_1 a_1); \\ N_1 b_n^m j_n(N_2 a_1) + N_1 b_n^m h_n^{(1)}(N_2 a_1) &= N_2 b_n^t j_n(N_1 a_1); \end{aligned} \quad (2.11)$$

where $N_1 = k_0 a_1, N_2 = \frac{k_1}{k_0}$.

In the similar manner, it follows from $i_1 H_m = i_1 H_t$ that

$$\begin{aligned} a_n^m j_n(N_2 a_1) + a_n^m h_n^{(1)}(N_2 a_1) &= a_n^t j_n(N_1 a_1); \\ N_2 b_n^m j_n(N_2 a_1) + N_2 b_n^m h_n^{(1)}(N_2 a_1) &= N_1 b_n^t j_n(N_1 a_1); \end{aligned} \quad (2.12)$$

At the boundary $r = a_2$, in the same fashion, it follows from $i_1 (E_i + E_r) = i_1 E_m$ that

$$\begin{aligned} j_n(N_2 a_2) + a_n^r h_n^{(1)}(N_2 a_2) &= a_n^m j_n(N_2 a_2) + a_n^m h_n^{(1)}(N_2 a_2); \\ N_2 [a_n^r j_n(N_2 a_2) + N_2 b_n^r h_n^{(1)}(N_2 a_2)] &= b_n^m j_n(N_2 a_2) + b_n^m h_n^{(1)}(N_2 a_2); \end{aligned} \quad (2.13)$$

where $N_2 = k_0 a_2, N_2 = \frac{k_2}{k_0}$.

In the meanwhile, it follows from $i_1 (H_i + H_r) = i_1 H_m$ that

$$\begin{aligned} a_n^r j_n(N_2 a_2) + [a_n^r h_n^{(1)}(N_2 a_2) + [a_n^r j_n(N_2 a_2)]^0] &= 0 a_n^m j_n(N_2 a_2) + a_n^m h_n^{(1)}(N_2 a_2); \\ 2 b_n^r h_n^{(1)}(N_2 a_2) + a_n^r j_n(N_2 a_2) &= N_2 b_n^m j_n(N_2 a_2) + N_2 b_n^m h_n^{(1)}(N_2 a_2); \end{aligned} \quad (2.14)$$

C. Calculation of Mie coefficients

1. Mie coefficient a_n^r

According to the above boundary conditions, one can arrive at the following matrix equation

[illegible]

For convenience, the 4×4 matrix on the left handed side of (2.15) is denoted by

$$A = \begin{pmatrix} 0 & A_{11} & A_{12} & A_{13} & 0 \\ A_{21} & A_{22} & A_{23} & 0 & C \\ 0 & A_{32} & A_{33} & A_{34} & A \\ 0 & A_{42} & A_{43} & A_{44} & \end{pmatrix} \quad (2.16)$$

whose determinant is

$$\det A = A_{11}A_{22}A_{33}A_{44} - A_{11}A_{22}A_{34}A_{43} - A_{11}A_{32}A_{23}A_{44} + A_{11}A_{42}A_{23}A_{34} \\ + A_{21}A_{12}A_{33}A_{44} + A_{21}A_{12}A_{34}A_{43} + A_{21}A_{32}A_{13}A_{44} - A_{21}A_{42}A_{13}A_{34} : \quad (2.17)$$

So, the M ie coefficient a_n^r is of the form

$$a_n^r = \frac{\det A_r}{\det A} \quad (2.18)$$

with

$$A_r = \begin{pmatrix} 0 & 1 \\ A_{11} & A_{12} & A_{13} & 0 \\ A_{21} & A_{22} & A_{23} & 0 \\ 0 & A_{32} & A_{33} & j_n(z) \\ 0 & A_{42} & A_{43} & z [z j_n(z)]^0 \end{pmatrix} \quad (2.19)$$

whose determinant is

$$\det A_r = A_{11}A_{22}A_{33} \left[2 \left(\frac{j_n}{2} \right)^0 \right] A_{11}A_{22} \frac{j_n}{2} (2) A_{43} + A_{11}A_{32}A_{23} \left[2 \left(\frac{j_n}{2} \right)^0 \right] + A_{11}A_{42}A_{23} \frac{j_n}{2} (2) \\ A_{21}A_{12}A_{33} \left[2 \left(\frac{j_n}{2} \right)^0 \right] + A_{21}A_{12} \frac{j_n}{2} (2) A_{43} + A_{21}A_{32}A_{13} \left[2 \left(\frac{j_n}{2} \right)^0 \right] A_{21}A_{42}A_{13} \frac{j_n}{2} (2): \quad (2.20)$$

2. M i e c o e c i e n t b_n^r

According to the above boundary conditions, one can arrive at the following matrix equation

$$\begin{aligned}
& \begin{array}{ccccccc}
0 & & & & h & & i_0 \\
N_2 [N_1 \ 1 \ j_h (N_1 \ 1)]^0 & N_1 [N_2 \ 1 \ j_h (N_2 \ 1)]^0 & N_1 [N_2 \ 1 h_n^{(1)} (N_2 \ 1)]^0 & 0 & 1 & 0 & 1 \\
N_1 [2 \ j_h (N_1 \ 1)] & N_2 [1 \ j_h (N_2 \ 1)] & h [N_2 \ 1 h_n^{(1)} (N_2 \ 1)]_{i_0} & h & 0 & i_0 & 0 \\
0 & N_2 [2 \ j_h (N_2 \ 2)]^0 & N_2 [2 h_n^{(1)} (N_2 \ 2)] & N_2 [2 h_n^{(1)} (2)] & A & \begin{array}{c} \text{at} \\ \text{B} \\ \text{C} \\ \text{D} \\ \text{E} \\ \text{F} \end{array} & \begin{array}{c} b_h^t \\ b_h^m \\ b_h^c \\ b_h^r \end{array} \\
0 & N_2 [0 \ j_h (N_2 \ 2)] & N_2 [0 h_n^{(1)} (N_2 \ 2)] & 2 h_n^{(1)} (2) & & & N_2 [2 \ j_h (2)]^0 \\
& & & & & & 2 \ j_h (2)
\end{array} = \begin{array}{c} \text{at} \\ \text{B} \\ \text{C} \\ \text{D} \\ \text{E} \\ \text{F} \end{array} : \quad (2.21)
\end{aligned}$$

For convenience, the 4×4 matrix on the left handed side of (2.21) is denoted by

$$\mathbf{B} = \begin{pmatrix} 0 & B_{11} & B_{12} & B_{13} & 0 & 1 \\ B_{21} & B_{22} & B_{23} & 0 & C & C \\ 0 & B_{32} & B_{33} & B_{34} & A & A \\ 0 & B_{42} & B_{43} & B_{44} & & \end{pmatrix} \quad (2.22)$$

whose determinant is

$$\begin{aligned} \det B = & B_{11}B_{22}B_{33}B_{44} - B_{11}B_{22}B_{34}B_{43} - B_{11}B_{32}B_{23}B_{44} + B_{11}B_{42}B_{23}B_{34} \\ & B_{21}B_{12}B_{33}B_{44} + B_{21}B_{12}B_{34}B_{43} + B_{21}B_{32}B_{13}B_{44} - B_{21}B_{42}B_{13}B_{34} : \end{aligned} \quad (2.23)$$

So, the Mie coefficient b_n^r is of the form

$$b_n^r = \frac{\det B_r}{\det B} \quad (2.24)$$

with

$$B_r = \begin{pmatrix} 0 & 1 \\ B_{11} & B_{12} & B_{13} & 0 \\ B_{21} & B_{22} & B_{23} & 0 \\ 0 & B_{32} & B_{33} & N_2 [2j_n(\rho)]^0 \\ 0 & B_{42} & B_{43} & 2j_n(\rho) \end{pmatrix} \begin{matrix} C \\ A \end{matrix} \quad (2.25)$$

whose determinant is

$$\begin{aligned} \det B_r = & B_{11}B_{22}B_{33} 2j_n(\rho) - B_{11}B_{22}N_2 [2j_n(\rho)]^0 B_{43} - B_{11}B_{32}B_{23} 2j_n(\rho) + B_{11}B_{42}B_{23}N_2 [2j_n(\rho)]^0 \\ & B_{21}B_{12}B_{33} 2j_n(\rho) + B_{21}B_{12}N_2 [2j_n(\rho)]^0 B_{43} + B_{21}B_{32}B_{13} 2j_n(\rho) - B_{21}B_{42}B_{13}N_2 [2j_n(\rho)]^0 : \end{aligned} \quad (2.26)$$

III. REDUCED TO THE CASE OF SINGLE-LAYERED SPHERE

In order to see whether the above Mie coefficients in the case of double-layered sphere is correct or not, we consider the reduction problem of double-layered case to the single-layered one when the following reduction conditions are satisfied: $A_{11} = A_{12}, A_{21} = A_{22}, N_1 = N_2, \rho_1 = \rho, \rho_2 = \rho$.

By lengthy calculation, we obtain

$$\begin{aligned} \det A = & (A_{11}A_{23} - A_{21}A_{13})(A_{42}A_{34} - A_{32}A_{44}) \\ & - h_0 h_n^{(1)}(\rho) [N_2 2j_n(N_2 \rho)]^0 h_n^{(1)}(\rho) \end{aligned} \quad (3.1)$$

and

$$\det A_r = (A_{11}A_{23} - A_{21}A_{13}) [N_2 2j_n(N_2 \rho)]^0 j_n(\rho) - j_n(N_2 \rho) 2 [2j_n(\rho)]^0 : \quad (3.2)$$

Thus

$$a_n^r = \frac{[N_2 2j_n(N_2 \rho)]^0 j_n(\rho) - j_n(N_2 \rho) 2 [2j_n(\rho)]^0}{j_n(N_2 \rho) 2 h_n^{(1)}(\rho) - [N_2 2j_n(N_2 \rho)]^0 h_n^{(1)}(\rho)} ; \quad (3.3)$$

which is just the Mie coefficient a_n^r of single-layered sphere [11].

In the same fashion, when the reduction conditions $B_{11} = B_{12}, B_{21} = B_{22}, N_1 = N_2, \rho_1 = \rho, \rho_2 = \rho$ are satisfied, one can arrive at

$$\begin{aligned} \det B = & (B_{11}B_{23} - B_{21}B_{13})(B_{42}B_{34} - B_{32}B_{44}) \\ & - h_0 h_n^{(1)}(\rho) [N_2 2j_n(N_2 \rho)]^0 2h_n^{(1)}(\rho) - N_2^2 [N_2 2j_n(N_2 \rho)]^0 h_n^{(1)}(\rho) \end{aligned} \quad (3.4)$$

and

$$\det B_r = (B_{11}B_{23} - B_{21}B_{13}) N_2^2 [N_2 2j_n(N_2 \rho)]^0 [2j_n(\rho)]^0 - [N_2 2j_n(N_2 \rho)]^0 2j_n(\rho) : \quad (3.5)$$

So,

$$b_n^r = \frac{N_2^2 [N_2 2j_n(N_2 \rho)]^0 [2j_n(\rho)]^0 - [N_2 2j_n(N_2 \rho)]^0 2j_n(\rho)}{[N_2 2j_n(N_2 \rho)]^0 2h_n^{(1)}(\rho) - N_2^2 [N_2 2j_n(N_2 \rho)]^0 h_n^{(1)}(\rho)} ; \quad (3.6)$$

which is just the Mie coefficient b_n^r of single-layered sphere [11].

This, therefore, means that the Mie coefficients of double-layered sphere presented here is right.

IV . ALL THE M IE COEFFIC IENTS INSIDE THE DOUBLE-LAYERED SPHERE

A . M ie coe cients a_n^t , a_n^m and a_n^m

The 4 4 matrix A_t is

$$A_t = \begin{pmatrix} 0 & 0 & A_{12} & A_{13} & 0 & 1 \\ 0 & 0 & A_{22} & A_{23} & 0 & C \\ 0 & j_h(\rho) & A_{32} & A_{33} & A_{34} & C \\ 0 & 2[2j_h(\rho)]^0 & A_{42} & A_{43} & A_{44} & C \end{pmatrix} \quad (4.1)$$

whose determinant is

$$\det A_t = j_h(\rho) A_{12} A_{23} A_{44} - j_h(\rho) A_{22} A_{13} A_{44} - 2[2j_h(\rho)]^0 A_{12} A_{23} A_{34} + 2[2j_h(\rho)]^0 A_{22} A_{13} A_{34} : \quad (4.2)$$

So,

$$a_n^t = \frac{\det A_t}{\det A} : \quad (4.3)$$

The 4 4 matrix A_m is

$$A_m = \begin{pmatrix} 0 & A_{11} & 0 & A_{13} & 0 & 1 \\ 0 & A_{21} & 0 & A_{23} & 0 & C \\ 0 & 0 & j_h(\rho) & A_{33} & A_{34} & C \\ 0 & 2[2j_h(\rho)]^0 & A_{43} & A_{44} & C & C \end{pmatrix} \quad (4.4)$$

whose determinant is

$$\det A_m = A_{11} j_h(\rho) A_{23} A_{44} + A_{11} 2[2j_h(\rho)]^0 A_{23} A_{34} + A_{21} j_h(\rho) A_{13} A_{44} - A_{21} 2[2j_h(\rho)]^0 A_{13} A_{34} : \quad (4.5)$$

So,

$$a_n^m = \frac{\det A_m}{\det A} : \quad (4.6)$$

The 4 4 matrix A_m is

$$A_m = \begin{pmatrix} 0 & A_{11} & A_{12} & 0 & 0 & 1 \\ 0 & A_{21} & A_{22} & 0 & 0 & C \\ 0 & 0 & A_{32} & j_h(\rho) & A_{34} & C \\ 0 & 0 & A_{42} & 2[2j_h(\rho)]^0 & A_{44} & C \end{pmatrix} \quad (4.7)$$

whose determinant is

$$\det A_m = A_{11} A_{22} j_h(\rho) A_{44} - A_{11} A_{22} A_{34} 2[2j_h(\rho)]^0 - A_{21} A_{12} j_h(\rho) A_{44} + A_{21} A_{12} A_{34} 2[2j_h(\rho)]^0 : \quad (4.8)$$

So,

$$a_n^m = \frac{\det A_m}{\det A} : \quad (4.9)$$

It is readily verified that under the reduction conditions $A_{11} = A_{12}$, $A_{21} = A_{22}$, $N_1 = N_2$, $\rho_1 = \rho_2$, $\rho_1 = \rho_2$, one can arrive at

$$\det A_t = \det A_m ; \quad a_n^t = a_n^m ; \quad \det A_m = 0 ; \quad a_n^m = 0 ; \quad (4.10)$$

which means that the above Mie coefficient can be reduced to the those of single-layered sphere.

B.M ie coe cients b_n^t , b_n^m and b_n^n

The 4 4 matrix B_t is

$$B_t = \begin{pmatrix} 0 & 0 & B_{12} & B_{13} & 0 & 1 \\ B_{21} & 0 & B_{22} & B_{23} & 0 & C \\ 0 & N_2 [\dot{\alpha}_h (\alpha)]^0 & B_{32} & B_{33} & B_{34} & A \\ 0 & \dot{\alpha}_h (\alpha) & B_{42} & B_{43} & B_{44} & 0 \end{pmatrix} \quad (4.11)$$

whose determinant is

$$\det B_t = N_2 [\dot{\alpha}_h (\alpha)]^0 B_{12} B_{23} B_{44} - N_2 [\dot{\alpha}_h (\alpha)]^0 B_{22} B_{13} B_{44} - \dot{\alpha}_h (\alpha) B_{12} B_{23} B_{34} + \dot{\alpha}_h (\alpha) B_{22} B_{13} B_{34} : \quad (4.12)$$

So,

$$b_n^t = \frac{\det B_t}{\det B} : \quad (4.13)$$

The 4 4 matrix B_m is

$$B_m = \begin{pmatrix} 0 & B_{11} & 0 & B_{13} & 0 & 1 \\ B_{21} & 0 & B_{23} & 0 & C & 0 \\ 0 & N_2 [\dot{\alpha}_h (\alpha)]^0 & B_{33} & B_{34} & A & 0 \\ 0 & \dot{\alpha}_h (\alpha) & B_{43} & B_{44} & 0 & 0 \end{pmatrix} \quad (4.14)$$

whose determinant is

$$\det B_m = B_{11} N_2 [\dot{\alpha}_h (\alpha)]^0 B_{23} B_{44} + B_{11} \dot{\alpha}_h (\alpha) B_{23} B_{34} + B_{21} N_2 [\dot{\alpha}_h (\alpha)]^0 B_{13} B_{44} - B_{21} \dot{\alpha}_h (\alpha) B_{13} B_{34} : \quad (4.15)$$

So,

$$b_n^m = \frac{\det B_m}{\det B} : \quad (4.16)$$

The 4 4 matrix B_n is

$$B_n = \begin{pmatrix} 0 & B_{11} & B_{12} & 0 & 0 & 1 \\ B_{21} & B_{22} & 0 & 0 & C & 0 \\ 0 & B_{32} & N_2 [\dot{\alpha}_h (\alpha)]^0 & B_{34} & A & 0 \\ 0 & B_{42} & \dot{\alpha}_h (\alpha) & B_{44} & 0 & 0 \end{pmatrix} \quad (4.17)$$

whose determinant is

$$\det B_n = B_{11} B_{22} N_2 [\dot{\alpha}_h (\alpha)]^0 B_{44} - B_{11} B_{22} B_{34} \dot{\alpha}_h (\alpha) - B_{21} B_{12} N_2 [\dot{\alpha}_h (\alpha)]^0 B_{44} + B_{21} B_{12} B_{34} \dot{\alpha}_h (\alpha) : \quad (4.18)$$

So,

$$b_n^n = \frac{\det B_n}{\det B} : \quad (4.19)$$

It is readily verified that under the reduction conditions $B_{11} = B_{12}, B_{21} = B_{22}, N_1 = N_2, \alpha_1 = \alpha_2, \dot{\alpha}_1 = \dot{\alpha}_2$, one can arrive at

$$\det B_t = \det B_m ; \quad b_n^t = b_n^m ; \quad \det B_n = 0 ; \quad b_n^n = 0 ; \quad (4.20)$$

which means that the above M ie coe cient can be reduced to the those of single-layered sphere.

In this section we briefly discuss several cases with left-handed media involved.

(i) If the following conditions $h_n^{(1)}(N_{21}) = 0$, $h_n^{(1)}(N_{21}) = 0$, $h_n^{(1)}(N_{22}) = h_n^{(1)}(2)$, $h_{i_0}^{(1)}(N_{22}) = h_{i_0}^{(1)}(2)$, $N_2 = 2 = 2 = 1$ and $\epsilon_0 = +1$ are satisfied, then it is easily verified that

$$\det A = 0; \det B = 0 \text{ and } a_n^r = 1; b_n^r = 1: \quad (5.1)$$

Thus in this case the extinction cross section of the two-layered sphere containing left-handed media is rather large.

(ii) If the following conditions $h_n^{(1)}(N_{21}) = 0$, $h_n^{(1)}(N_{21}) = 0$, $h_n^{(1)}(N_{22}) = j_n(2)$, $h_n^{(1)}(N_{22}) = [j_n(2)]^0$, $N_2 = 2 = 2 = 1$ and $\epsilon_0 = +1$ are satisfied, then it is easily verified that

$$\det A_r = 0; \det B_r = 0 \text{ and } a_n^r = 0; b_n^r = 0: \quad (5.2)$$

Thus in this case the extinction cross section of the two-layered sphere containing left-handed media is negligibly small.

(iii) The case with $N_1 = 1 = 1 = 1$, $N_2 = 2 = 2 = 0$, $N_0 = 0 = 0 = +1$ is of physical interest, which deserves consideration by using the Mie coefficients presented above.

Based on the calculation of Mie coefficients, one can treat the scattering problem of double-layered sphere irradiated by a plane wave. The scattering and absorption properties of double-layered sphere containing left-handed media can thus be discussed in detail, which are now under consideration and will be submitted elsewhere for the publication.

Acknowledgements This project is supported by the National Natural Science Foundation of China under the project No. 90101024.

-
- [1] Smith, D. R., Padilla, W. J., Vier, D. C. et al., Phys. Rev. Lett. 84, 4184 (2000).
 - [2] Klimov, V. V., Opt. Comm. 211, 183 (2002).
 - [3] Pendry, J. B., Holden, A. J., Robbins, D. J. and Stewart, W. J., IEEE Trans. Microwave Theory Tech. 47, 2075 (1999).
 - [4] Shelby, R. A., Smith, D. R. and Schultz, S., Science 292, 77 (2001).
 - [5] Ziolkowski, R. W., Phys. Rev. E 64, 056625 (2001).
 - [6] Veselago, V. G., Sov. Phys. Usp. 10, 509 (1968).
 - [7] Pendry, J. B., Holden, A. J., Stewart, W. J. and Youngs, I., Phys. Rev. Lett. 76, 4773 (1996).
 - [8] Pendry, J. B., Holden, Robbins, D. J. and Stewart, W. J., J. Phys. Condens. Matter 10, 4785 (1998).
 - [9] Maslovski, S. I., Tretyakov, S. A. and Belov, P. A., Int. J. Microwave Opt. Tech. Lett. 35, 47 (2002).
 - [10] Wang, W. and Asher, S. A., J. Am. Chem. Soc. 123, 12528 (2001).
 - [11] Stratton, J. A., Electromagnetic theory (New York: McGraw-Hill) (1941).