

Energy band structure and intrinsic coherent properties in two weakly linked Bose-Einstein Condensates

Wei-Dong Li¹, Yunbo Zhang^{1,2}, J.-Q. Liang¹

¹ Department of Physics and Institute of Theoretical Physics, Shanxi

University, Taiyuan 030006, China

² Department of Physics, University of Turku, FIN-20014 Turun yliopisto,

Finland

(January 9, 2022)

The energy band structure and energy splitting due to quantum tunneling in two weakly linked Bose-Einstein condensates were calculated by using the instanton method. The intrinsic coherent properties of Bose Josephson junction were investigated in terms of energy splitting. For $E_C = E_J$, the energy splitting is small and the system is globally phase coherent. In the opposite limit, $E_C \ll E_J$, the energy splitting is large and the system becomes a phase dissipation. Our results suggest that one should investigate the coherence phenomena of BJJ in proper condition such as $E_C = E_J$.

PACS numbers: 03.75.Fi, 05.30.Jp, 64.60.My, 67.40.Fd

I. INTRODUCTION

Two weakly linked Bose-Einstein condensates (BECs) behave as the superconductor Josephson junction. This remarkable feature has been investigated both theoretically [1-6] and experimentally [8-12]. The existence of a Josephson current through a potential barrier between two superconductors or between two superfluids is a direct manifestation of macroscopic quantum coherence [13]. The experimental realization of Bose-Einstein condensation (BEC) of weakly interacting alkali atoms [8] has provided a route to study neutral superfluids in a controlled and tunable environment [9]. The possibility of loading a BEC in a one-dimensional periodic potential has allowed the observation of quantum phase effects on

a macroscopic scale such as quantum interference [10], superfluidity on a local scale [11] and an oscillating atomic current in Josephson junction arrays [12].

The analogy of the voltage-current characteristic in superconductor Josephson junction was proposed theoretically in Bose-Josephson junction [25]. The macroscopic BEC's coherence has been demonstrated by interference experiments [14], and the first evidence of the coherent tunneling in an atomic array, related to the "ac" Josephson effect, has been reported [10]. A "dc" current-biased Bose-Josephson junction can be simulated by a tunneling barrier moving with constant velocity across the trap [3]. At a critical velocity of the barrier (proportional to the critical tunneling current), a sharp transition between the "dc" and "ac" Josephson regimes was predicted. Thus two weakly linked condensates exhibit the analog of the resistively shunted superconductor Josephson junctions. The "secondary" quantum macroscopic effects in small capacitance Josephson junction had attracted much attention both from theorists and experimentalists for decades [15-18]. These phenomena manifest the quantum behavior of a Josephson junction as a macroscopic object, in contrast with such "primary" quantum macroscopic phenomena as the Josephson junction itself [15]. In these cases, quantum fluctuations of the phase difference φ across the junctions become important. This necessitates treating the phase as a quantum operator $\hat{\varphi}$, which is canonically conjugate to the operator \hat{N} . Novel macroscopic quantum phenomena, such as "Bloch" oscillation, had been reported in current-biased Josephson junctions [15]. To understand these effects and present a general picture of the low temperature dynamics of Josephson junctions, a simple theory [16-18] has been suggested based on the extended coordinates, $\varphi^2 \in [-1; +1]$.

More recently, the first experimental of BJJ have been realized in a purely magnetic double-well potential [24]. It will stimulate the further study on the BJJ. Then, can we investigate the "secondary" quantum macroscopic effects [15] in two weakly linked BECs? It was argued in [2,3,6] that the dynamics of the phase of this system can be mapped onto the sine-Gordon quantum mechanical Hamiltonian (1). In this paper, we calculated the energy band structure, "Bloch" wave and the energy splitting of the Bose-Josephson junction due

to the quantum tunneling using the instanton method [21]. As a simply application, we investigate the coherent properties of the Bose-Josephson junction. The results from this method agree exactly with the prediction of Stringari [6].

II. MODEL AND ITS INSTANTONS SOLUTION

All quantities describing the junction should be considered as operators rather than the classical variables. The operators corresponding to them main variables, the phase difference ϕ and the particle number N of the Bose-Josephson junction, satisfy the commutation relation: $[\phi, N] = i$, so that ϕ and N are related by the Heisenberg uncertainty relation which was disregarded by the "classical" theory of the Bose-Josephson junction [2].

The idealization of two weakly linked BECs can be described as a two-mode bosonic system [2]. In the "phase" representation the relevant quantum observables are the difference of phases and number of atoms between the two condensates in each trap. The Hamiltonian can be written in terms of a "quantum pendulum" (Mathieu) equation

$$\hat{H} = \frac{E_C}{2} \frac{\partial^2}{\partial \phi^2} + E_J \cos \phi \quad (1)$$

The "charging energy" E_C and the "Josephson coupling energy" E_J can be calculated as overlap integrals [2,3],

$$\begin{aligned} E_C &= 2g \int_{-Z}^Z |\psi_1(r)|^4 dr = 2g \int_{-Z}^Z |\psi_2(r)|^4 dr; \\ E_J &= N \int_{-Z}^Z |\psi_1(r)|^2 \left[\frac{\hbar^2}{2m} |\psi_2(r)|^2 + V_{\text{ext}} + \frac{gN}{2} (|\psi_1(r)|^2 + |\psi_2(r)|^2) \right] |\psi_2(r)|^2 dr \end{aligned} \quad (2)$$

with the one-body wave functions $\psi_1(r)$, $\psi_2(r)$ localized in the trap 1, 2; and $\int_{-R}^R |\psi_1(r)|^2 |\psi_2(r)|^2 dr = 0$, $\int_{-R}^R |\psi_{1,2}(r)|^2 dr = 1$, $g = \frac{4\hbar^2 a}{m}$; a is the scattering length and m the atomic mass; $N = \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2$ is the total number of atoms.

For convenience, we modify the potential as $U_J(\phi) = E_J(1 + \cos \phi)$. Then, the Hamiltonian \hat{H} in Eq. (1) can be considered as one-dimensional quantum particles with mass $m = \frac{\hbar^2}{E_C}$ moving along the ϕ axis in sine-Gordon potential $U_J(\phi)$. Due to the translational

symmetry ($\phi \rightarrow \phi + 2\pi$) of the Hamiltonian (1), the set of the eigenfunctions should include the Bloch wave functions

$$\begin{aligned} \psi_n(\phi) &= u_{n,q}(\phi) \exp(iq\phi); \quad n = 0, 1, 2, \dots; \\ u_{n,q}(\phi) &= u_{n,q}(\phi + 2\pi); \quad 1 > q > -1 \end{aligned} \quad (3)$$

where q is an arbitrary (real) constant vector. Substitution of the wave function (3) in the Schrodinger equation leads immediately to the picture of band energy spectrum and relative well known effects in solid state theory. From Ref. [17,18], this is nothing but Mathieu equation, so that these functions can be readily calculated. Some of their asymptotic properties (in tight-binding limit and weak-binding limit) can be expressed analytically.

The energy band structure and energy splitting for the sine-Gordon potential can be calculated alternatively by instanton method [19,20]. The advantage of this nonperturbative method, as presented here, is that it gives not only a more accurate description of the tunneling phenomena but also a comprehensive physical understanding in the context of quantum field theory.

The effective Lagrangian is

$$L = \frac{1}{2}m \left(\frac{d\phi}{dt} \right)^2 - U_J(\phi); \quad (4)$$

The classical solution which extremizes the action is seen to satisfy the equation of motion

$$\frac{1}{2}m \left(\frac{d\phi_c}{d\tau} \right)^2 + U_J(\phi_c) = E_{cl}; \quad (5)$$

where the Wick rotation $t \rightarrow i\tau$ has taken the system into Euclidean time. Equation (5) can be regarded as the equation of motion of a pseudo-particle with the classical energy $E_{cl} = 0$, which is a constant of integration. With E_{cl} being confined to a region $0 \leq E_{cl} \leq E_J$, the configuration ϕ_c becomes periodic such that $\phi_c(\tau + T) = \phi_c(\tau)$, which now corresponds to the periodic boundary condition in the space coordinate [19]. The classical solution is

$$\phi_c = 2 \arcsin [k \operatorname{sn}(\sqrt{E_{cl}}(\tau - \tau_0))]; \quad (6)$$

where $\omega_0 = \sqrt{E_J E_c} / \hbar$ is the classical plasma frequency [6], sn is the Jacobian elliptic function with modulus $k = \sqrt{1 - E_{cl}/2E_J}$. The elliptic function $\text{sn}(\omega_0(\tau + \tau_0))$ has a period $4K(k)$ with $K(k)$ the complete elliptic integral of first kind. For zero energy $E_{cl} = 0$ ($k = 1$), the periodic solution reduces to the vacuum instanton configuration $\phi_c = 2 \arcsin[\tanh(\omega_0(\tau + \tau_0))]$.

III. ENERGY BAND STRUCTURE AND THE TRANSITION AMPLITUDE FOR QUANTUM TUNNELING

The sine-Gordon potential has an infinite number of degenerate vacua. Quantum tunneling between neighboring vacua leads to the level splitting, while the levels extend to bands due to the translational symmetry expressed by $U_J(\tau + 2\pi) = U_J(\tau)$. In the narrow-band approximation one finds, for the energy, the expression

$$E = E_i + \sum_n J(R_n - R_0) e^{iq(R_n - R_0)}; \quad (7)$$

where E_i denotes the i th eigenvalue of the energy in each well for the harmonic oscillator potential $U(\phi) = \frac{1}{2}(\phi - R_n)^2$, $R_n = 2n\pi$ is the position of the n th minimum, $J(R_n - R_0) = \int_{R_0}^{R_n} \phi_i(\phi - R_0) [U_J(\phi) - U(\phi)] \phi_i(\phi - R_n) d\phi$ is the overlap integral, ϕ_i is the eigenfunction corresponding to eigenvalue E_i , and q is the Floquet parameter associated with the Bloch wave function. If only the contribution from the nearest neighbors is taken into account, i.e., $J(R_n - R_0)$ for $|n| > 1$ is taken to be zero, the energy band formula reduces to

$$E = E_i + 2J \cos(2q\phi); \quad (8)$$

The parameter J is just the level splitting resulting from quantum tunneling (the wave functions are periodic for $q = 0$). We will consider the case of potential wells surrounded by very high potential barriers with correspondingly small tunneling contribution to the eigenvalues. They are almost those of degenerate harmonic oscillators, and in this asymptotic case we are not concerned with the entire bands but only with their edges which correspond

to alternately even and odd states. Then, we suppose ψ_{iR}, ψ_{iL} are degenerate eigenstates in neighboring wells, respectively with the same energy eigenvalue E_i such that $H^0 \psi_{iR,L} = E_i \psi_{iR,L}$ where H^0 is the Hamiltonian of the harmonic oscillator as the zero order approximation of the system. The degeneracy will be removed by the small tunneling effect which leads to the level splitting. The eigenstates of the Hamiltonian H become

$$\psi_{io} = \frac{1}{\sqrt{2}} (\psi_{iR} + \psi_{iL}); \quad \psi_{ie} = \frac{1}{\sqrt{2}} (\psi_{iR} - \psi_{iL}) \quad (9)$$

with eigenvalues $E_i \pm E_i$, respectively. E_i denotes the shift of one oscillator level. It is obvious that $\langle \psi_{iR} | H | \psi_{iL} \rangle = H^0 \psi_{iL} = 2J = -E_i$. In the following we calculate this energy shift E_i as resulting from periodic instantons and instanton-anti-instanton pairs.

The amplitude for a transition from one well to its neighboring well at the energy E_i due to instanton tunneling can be written as

$$A_{+,-} = \pm \hbar E_i \int_0^T dt \langle \psi_{iL} | \psi_{iR} \rangle e^{2 \frac{E_i T}{\hbar}} \sinh(2 \frac{E_i T}{\hbar}); \quad (10)$$

where we neglect overlap of the wave functions which dominate over either well. The amplitude (10) can also be calculated with the help of the path-integral method, $A_{+,-} = R \int_{\psi_{i+}}^{\psi_{i-}} K(\psi_f; \psi_f; \psi_i; \psi_i) e^{-E_i T} d\psi_f d\psi_i$, where the Feynman kernel is defined as usual by

$$K(\psi_f; \psi_f; \psi_i; \psi_i) = \int_{\psi_i}^{\psi_f} D[\psi] e^{-S[\psi]} \quad (11)$$

with $\psi_f = (\psi_f)$, $\psi_i = (\psi_i)$ and $\tau_f - \tau_i = 2T$. What we are interested in is this expression in the limits $\tau_i \rightarrow a$, $\tau_f \rightarrow a$ (a are the turning points), namely, the tunneling propagator through one of the barriers. $S = \int_{\tau_i}^{\tau_f} dt [\frac{1}{2} m (\frac{d}{dt})^2 + U_J(\psi)]$ is the Euclidean action of the pseudo-particle, and $\psi_{i+}(\psi_i) = (\psi_{i-}(\psi_i))$ is the wave function of the right- (left-) hand wells.

The functional integral $K(\psi_f; \psi_f; \psi_i; \psi_i)$ can be evaluated with the stationary method by expanding $\psi(\tau)$ about the classical trajectory $\psi_c(\tau)$ and thus we set $\psi(\tau) = \psi_c(\tau) + \eta(\tau)$, where $\eta(\tau)$ is the small fluctuation with boundary conditions $\eta(a) = \eta(b) = 0$. Substitution of $\psi(\tau)$ into Eq.(11) and keeping only terms containing $\eta(\tau)$ up to the one-loop approximation yields $K(\psi_f; \psi_f; \psi_i; \psi_i) = \exp[\frac{S_c(\psi)}{\hbar}] I$, where $I = \int_{\psi_i}^{\psi_f} D[\eta] e^{-S[\eta]} = \int_{\psi_i}^{\psi_f} D[\eta] e^{-S[\eta]}$

is the fluctuation function integral with the fluctuation action $S = \int_0^T M \dot{c}^2 dt$; where $M = \frac{m}{2} \frac{d^2}{dt^2} + V''(c(t))$ is the second variational operator of the action. The classical action $S_c(t)$ is evaluated along the trajectory $c(t)$ so that

$$\begin{aligned} S_c(t) &= \int_0^T \left[\frac{m}{2} \dot{c}^2 + V(c(t)) \right] dt \\ &= 2E_T + \frac{8E_J}{\hbar} [E(k) - K(k)]; \end{aligned} \quad (12)$$

where $E(k)$ denotes the complete elliptic integral of the second kind and $k^2 = 1 - \frac{E_C}{E_J}$. Following the standard procedure of the periodic instanton calculation in Refs. [19] and [20], the functional integral I can be written as $I = \frac{1}{2} \frac{\hbar}{N(c)} N(c_f) \int_0^T \frac{d}{dt} \frac{1}{N^2(c)} dt$, where $N(c) = \frac{d}{dt} c(t)$, is the zero eigenmode of M . To obtain the desired results, the contributions from the infinite number of the instantons and anti-instanton pairs have to be taken into account. The total amplitude is found to be

$$A_{+,-} = e^{2 \frac{E_J}{\hbar} T} \sinh \left(\frac{\hbar}{2K(k^0)} e^{\frac{W}{\hbar}} \right); \quad (13)$$

with $W = \frac{8E_J}{\hbar} [E(k) - K(k)]$. Comparing (13) and (10) leads to $E = \frac{\hbar \hbar_0}{4K(k^0)} \exp \left(\frac{W}{\hbar} \right)$. Rescaling this formula in the unit of E_J , we arrive at our final result.

$$\frac{E}{E_J} = \frac{E_C}{E_J} \frac{1}{4K(k^0)} \exp \left[8 \frac{E_J}{E_C} (E(k) - K(k)) \right]; \quad (14)$$

The formula (14) shows that the widths of the energy bands are very sensitive to the dimensionless parameter of E_C/E_J , which is a critical parameter characterizing the dynamical properties of the system such as the coherence [6] and the transition [15]. A direct application of this energy splitting is to investigate the coherence properties of BJJ. Quantum coherence requires that the relative phase of the order parameter should be preserved over times of the order τ^{-1} , which is just the tunneling amplitude [23]. From the above calculation, we know that the energy splitting E is proportional to E_C . Therefore the energy splitting describes the coherent properties of the system.

Fig. 1 shows the energy splitting E/E_J as a function of the ratio E_C/E_J in the ground state ($E_{cl} = 0$). The figure shows that for values of E_J smaller than E_C the energy

splitting is significantly increased, indicating the occurrence of a continuous transition to the phase dissipation (the number squeezed regime) [6,7]. In the limit $E_C = E_J \rightarrow 1$ the system undergoes small oscillations around the equilibrium. In this limit the correlation between the neighboring wells is small. One can regard the systems as a globally coherent object described by a unique order parameter. On the contrary $E_C = E_J \rightarrow 1$ the behavior of the system is very different. The quantum fluctuation is enhanced due to the increasing of the tunneling between the neighboring wells in phase space, showing that the relative phase between two condensates are distributed in a random way. At the same time, the fluctuation of the relative number of atoms in two traps becomes smaller and smaller.

It is interesting to investigate the properties of the systems under the excited states ($E_{cl} \neq 0$) which may be excited by the thermal fluctuations or other reasons. We plotted the energy splitting as a function of the dimensionless parameter $E_{cl} = E_J$ for two different values of $E_C = E_J$ in Fig. 2. One clearly observes that even if quantum effects are small the decoherence due to higher excited states may become important. It is interesting that the response of the system to the decoherence perform a total different behavior. States with small quantum effects are more sensitive to the decoherence fluctuations than those with relatively large values of $E_C = E_J$. The system with little higher $E_C = E_J$ state should be a good candidate to investigate the quantum coherence phenomena, in $E_C = E_J \rightarrow 1$. Because in this case the system with a good coherence will preserve its coherence for a relative large range of the energy perturbation.

Energy spectrum and energy splitting due to quantum tunneling in BJJ have been calculated by means of instanton method. Based on this energy splitting formula, we also investigated the coherence property of BJJ. Our results agree exactly with that in Ref. [6]. This analysis makes it possible to investigate the "secondary" quantum phenomena in BJJ (see, for instance, [18]) and presents a general picture of low temperature dynamics of Josephson junctions in BEC.

It is great pleasure to thank A. Smirzi and L.P. Pitaevskii for any fruitful discussions. This work is supported by NSF of China (Grants 10175039 and 90203007), Science Funding

(Grant 20001003) and Youth Funding (Grant 20011002) from Shanxi Province of China.

- [1] J.M. Martinis et al, Phys. Rev. Lett. 55, 1543 (1985).
- [2] A. Smerzi et al, Phys. Rev. Lett. 79, 4950 (1997); S. Raghavan et al, Phys. Rev. A 59, 620 (1999); I. Marino et al, Phys. Rev. A 60, 487 (1999).
- [3] S. Giovanazzi et al, Phys. Rev. Lett. 84, 3521 (2000).
- [4] D. Choi and Q. Niu, Phys. Rev. Lett. 82, 2022 (1999), O. Morsch et al, Phys. Rev. Lett. 87 140402 (2001).
- [5] J. R. Anglin et al, Phys. Rev. A 64, 635605 (2001).
- [6] L. Pitaevskii et al, Phys. Rev. Lett. 87, 180402 (2001).
- [7] M. Krammer, L. Pitaevskii and S. Stringari, Phys. Rev. Lett. 88, 180404 (2002).
- [8] M. Inguscio et al, Eds, Bose-Einstein Condensates in Atomic Lasers (IOS Press, Amsterdam, 1999).
- [9] F. Dalfovo et al, Rev. Mod. Phys. 71, 463 (1999); A. J. Leggett, Rev. Mod. Phys. 73, 307 (2001).
- [10] B. P. Anderson et al, Science 282, 1686 (1998); W. M. Liu et al, Phys. Rev. Lett. 84, 2294 (2000); W. D. Li et al, Phys. Rev. A 64, 015602 (2001).
- [11] S. Burger et al, Phys. Rev. Lett. 86, 4447 (2001).
- [12] F. S. Cataliotti et al, Science 293, 843 (2001).
- [13] P. W. Anderson, Basic Notions of Condensed Matter Physics (Benjamin-Cummings, Menlo Park, 1984).
- [14] M. R. Andrews et al, Science 275, 637 (1997); D. S. Hall et al, Phys. Rev. Lett. 81, 1539

- (1998); Phys. Rev. Lett. 81, 1543 (1998).
- [15] G. Schon and A. D. Zaikin, Phys. Rep. 198, 237 (1990).
- [16] W. Zwerger, A. T. Dorsey and Matthew P. A. Fisher, Phys. Rev. B 34, 6518 (1986), S. M. Apenko, Phys. Lett. A 142, 277 (1989).
- [17] K. K. Likharev et al, J. Low. Temp. 59, 347 (1985).
- [18] D. V. Averin and K. K. Likharev, Mesoscopic Phenomena in Solids, edited by B. L. Altshuler et al. (Elsevier Science Publishers B. V., 1991).
- [19] J. Q. Liang et al, Phys. Lett. B 282, 105 (1992).
- [20] J. Q. Liang et al, Phys. Rev. D 51, 718 (1995).
- [21] S. Coleman, Phys. Rev. D 15, 2929 (1977); *ibid* D 16, 1962 (1977); Nucl. Phys. B 298, 178 (1988).
- [22] C. J. Milburn et al, Phys. Rev. A 55, 4318 (1997); I. Zapata et al, Phys. Rev. A 57, R28 (1998); A. Zmerzi et al, Phys. Rev. A 61, 63601 (2000).
- [23] A. J. Leggett, Supp. Pro. Theore. Phys. 69 80 (1980).
- [24] T. G. Tiecke, M. Kemmann, Ch. Buggle, I. Shvarchuk, W. von Klitzing and J. T. M. Walraven, cond-mat/0211604.

Figure Captions

Fig. 1. Ground state energy splitting as a function of the ratio $E_C = E_J$.

Fig. 2. Energy splitting as a function of the ratio $E = E_J$ for $E_C = E_J = 1$ (solid line), $E_C = E_J = 3$ (dashed line).




Fig. 1




Fig.2