## D in ensional crossover of the therm odynam ic properties near the phase transition to superconducting state in type I superconductors

D.V.Shopova, T.P.Todorov<sup>y</sup>

CPCM Laboratory, Institute of Solid State Physics, Bulgarian Academ y of Sciences, BG-1784 So a, Bulgaria.

Corresponding author. E-m ail address: sho@ issp basbg

<sup>y</sup> Perm anent address: Joint Technical College at the Technical University of So a.

K ey words: uctuations, latent heat, order parameter, equation of state.

PACS:05.70 Jk, 64.60 Fr, 74.20 De, 74.78 Db,

## A bstract

A new approach to the treatment of magnetic uctuations in thin lms of type I superconductors is introduced. Results for the dependence of free energy, specic c heat and order parameter pro le on the lm thickness near the equilibrium phase transition point are obtained and discussed.

Recent studies [1, 2, 3] have shown that the Halperin-Lubensky-Ma e ect (HLM) [4], namely, the uctuation-driven change of the order of superconducting phase transition in a zero magnetic eld is much stronger in quasi-two-dimensional (quasi-2D) type I superconducting lm s than in bulk (3D) samples. This fact gives new opportunities for an experimental veri cation of the e ect in suitably chosen samples of thin superconducting lm s. Numerical values of thermodynamic quantities as latent heat, speci cheat capacity

and order parameter jump, have been calculated theoretically with the aim to support future experiments  $\beta$ ].

HLM e ect occurs as a result of the interaction between the superconducting order parameter and the vector potential  $A(\mathbf{x}) = A'(\mathbf{x})$  corresponding to the uctuation part of the magnetic induction B' = B'; note, that the mean value  $B_0$  of the magnetic induction is assumed to be zero. When this interaction is neglected the superconducting phase transition in a zero external magnetic eld is of second order. Taking into account the magnetic uctuations  $A'(\mathbf{x})$  and their gauge-invariant interaction with leads to an e ective G inzburg-Landau (GL) free energy that describes a rst order phase transition

at the phase transition point  $T_{c0} = T_c (B_0 = 0)$ . In type I superconductors the e ect is stronger than in type II superconductors and can be investigated theoretically with the help of a self-consistent (m ean-eld-like) approximation, in which the scalar eld can be considered uniform, i.e., independent of the spatial vector **x**. Here we shall use this approximation; a more detailed justication of this approach can be found in Refs. [1, 4]. The coupling between a scalar eld, like in superconductors, and a vector gauge eld, like A, is present also in models that describe other physical systems, for example, in scalar electrodynamics, certain liquid crystals and early universe phase transitions; see, e.g., [5] for a recent review. The problem is relevant also to quantum phase transitions in superconductors [7, 8] and other system s [9].

In this letter we investigate the dimensional (2D -3D) crossover of phase transition properties of type I superconducting  $\ln s$ . Our treatment is based on preceding investigations of HLM e ect [1, 3], where we have derived an e ective energy of the superconducting  $\ln$  from the G inzburg-Landau model. We consider the elective free energy [3] and related therm odynamic quantities in the framework of a reliable and relatively simple variant of the crossover theory, used in the study of disordered  $\ln s$  [10]. Because of considerable dimensions (11]. Here we outline a new general way of calculation of the therm odynamic functions dependence on the thickness of superconducting  $\ln s$ . The present results, as sum marized and discussed at the end of this paper, con rm and extend the recent investigations [1, 2, 3] of the phase transition properties in thin  $\ln s$  of type I superconductors. Our notations follow those in Ref. [12].

We consider the following e ective free energy density f() = F()=V of a thin superconducting lm of thickness  $L_0$  and volume  $V = (L_1 L_2 L_0)$  [1, 4]:

$$f() = aj j^{2} + \frac{b}{2}j j^{4} + k_{B}TJ[()];$$
(1)

where

$$J() = \int_{0}^{Z} \frac{dk}{2} kS(k;);; \qquad (2)$$

$$S = \frac{1}{L_0} \sum_{k_0 = 0}^{X_0} \ln 1 + \frac{()}{k^2 + k_0^2} ;$$
 (3)

In Eqs. (1){(3), () =  $_0j f$ , where  $_0 = (8 e^2 = m c^2); a = _0 (T = T_{c0})$  and b > 0 are the usual Landau parameters. They are related to the zero temperature coherence length  $_0 = (-2 = 4m = _0T_{c0})^{1=2}$ , the zero-temperature critical magnetic eld  $H_{c0} = _0T_{c0} (4 = b)^{1=2}$ , and the initial (unrenormalized) critical temperature  $T_{c0}$  [12]. The third term in the rhs. of Eq. (1) describes the e ect of the magnetic uctuations [1, 3]. In Eqs. (2){(3), the integral J() and the sum S(k; ) over the wave vector  $\mathbf{q} = (\tilde{\kappa}; k_0)$  are truncated by the upper cuto s and  $_{0}$ . The nite cuto is introduced for the wave number  $k = \frac{1}{2}$  j of the wave vector  $k = (k_{1};k_{2})$  and  $_{0}$  stands for  $k_{0}$ . The free energy (1) { (3) has been derived [1, 3] under the condition = [(T) = (T)]  $1 = \frac{1}{2}$ , i.e. for type I superconductors. Here, is the G inzburg-Landau number and the coherence length is given by  $(T) = (_{0} = \frac{p}{t_{0}})$ , where  $t_{0} = [(T - T_{c0}) = T_{c0}]$  1. The London penetration depth (T) is  $(T) = (_{0} = \frac{p}{t_{0}}); _{0} = (b = _{0} - T_{c0})^{1-2}$  corresponds to T = 0.

In preceding investigations [1, 2, 3] of thin (quasi-2D) Im, i.e., Im sobeying the condition  $a_0 L_0 1$ , where  $a_0$  is the lattice constant, only the  $(k_0 = 0)$  {term in the sum S has been taken into account. This approximation leads to a simple dependence of the therm odynamic quantities on the thickness  $L_0$ , but is not convenient when  $L_0$  varies in broad limits. Here we shall apply a more general treatment which gives the opportunity for the description of 2D -3D crossover from quasi-2D to 3D -superconductors  $L_0 L$ ; ( = 1;2). The choice of the cuto = ( =  $_0$ ) is consistent with the general requirement  $_0 < (T)$  for the validity of GL free energy for type I superconductors (see, e.g., Ref. [12]) and speci c requirements [1, 3] for the validity of the Landau expansion of the free energy (1) { (3) for extremely thin Ims (L<sub>0</sub> 0).

A sour study is based on the quasimacroscopic GL approach the second cuto  $_0$  should be again related to  $_0$  rather than to the lattice constant  $a_0$ , i.e.  $_0$  (1= $_0$ ), which means that phenomena at distances shorter than  $_0$  are excluded from our consideration. We shall assume that the lowest possible value of  $_0$  is (= $_0$ ), as is for , but we shall keep in m ind that both  $_0$  and can be extended to in nity provided the main contribution to the integral J () and the sum S come from the long wavelength limit (q  $_0$  1).

In a cbse vicinity of the phase transition point  $T_{c0}$  from norm al ( = 0) to M eissner state (j j > 0) the parameter <sup>2</sup> is small and the main contribution to the free energy f will be given by the terms in S with small wave vectors k . This allows an approximate but reliable treatment of the 2D -3D crossover by expanding the summation over  $k_0$  in (3) to in nity - <sub>0</sub> 1; for a justi cation, see Ref. [10, 11]. A variant of the theory when <sub>0</sub> is kept nite ( = <sub>0</sub> = = <sub>0</sub>) can also be developed but the results are too complicated [13]. Perform ing the summation and the integration in Eqs. (2) { (3) we obtain J ( ) = ( <sup>2</sup>=2 L<sub>0</sub>)I( ), where

$$I() = \int_{0}^{Z_{1}} dy \ln \left( \frac{\sinh \frac{1}{2} L_{0}}{\sinh \frac{1}{2} L_{0}} \frac{p + y}{\overline{y}} \right)^{\#}; \qquad (4)$$

The integral (4) has a logarithm ic divergence that corresponds to the in nite contribution of magnetic uctuations to the free energy of norm alphase ( $T_{c0} > 0$ ;' = 0). Such type of divergence is a common property of a lot of phase transition models. In the present case, as is in other systems, this divergence is irrelevant, because the divergent term does not depend on the order parameter and the free energy f() is de ned as the di erence between the total free energies of the superconducting and norm al phases: f() = ( $f_S = f_N$ ).

Introducing a dimensionless order parameter  $' = (=_0)$ , where  $_0 = (_0T_{c0}=b)^{1=2}$  is the value of at T = 0, we obtain the free energy (1) in the form

$$f(') = \frac{H_{c0}^2}{8} 2t_0'^2 + \frac{b}{2}j' \frac{4}{3} + 2(1+t_0)CI('^2) ; \qquad (5)$$

with I ( $\prime$ <sup>2</sup>) given by Eq. (4), = (1=)<sup>2</sup>, = =  $_0$ , and

$$C = \frac{2^{-2} k_{\rm B} T_{\rm c0}}{L_0 {}^{2}_{0} {\rm H}^{2}_{\rm c0}} :$$
 (6)

From the equation of state (@f=@'=0) we nd two possible phases:  $'_0 = 0$  and the superconducting phase ( $'_0 > 0$ ) de ned by the equation

$$t_0 + \prime^2 + \frac{(1 + t_0)C L_0 }{4 \ _0^2} K \ (\prime^2) = 0;$$
 (7)

where

$$K(z) = \int_{0}^{Z_{1}} dy \frac{\coth \frac{1}{2} L_{0}}{p \frac{y + z}{y + z}} :$$
(8)

The analysis of the stability condition  $(0^2 f = 0'^2 0)$  shows that the norm alphase is a minimum of f(') for t<sub>0</sub> 0, whereas the superconducting phase is a minimum of f(') provided

$$1 > \frac{1}{4} (1 + t_0) C L_0 \qquad {}^2K (\prime_0^2);$$
(9)

where

$$K^{*}(z) = \int_{0}^{Z_{1}} \frac{dy}{y+z} \frac{\cosh \frac{1}{2} L_{0}}{p + z} + \frac{L_{0}}{2 \sinh^{2} \frac{1}{2} L_{0}} \frac{\#}{y+z} + \frac{L_{0}}{2 \sinh^{2} \frac{1}{2} L_{0}} \frac{\#}{y+z}$$
(10)

The entropy jump  $s = (S=V) = [df('_0)=dT]$  per unit volume at the equilibrium point of the phase transition  $T_c \in T_{c0}$  is obtained in the form

$$s(T_{c}) = \frac{H_{c0}^{2} \prime_{c0}^{2}}{4 T_{c0}} 1 + \frac{C I(\prime_{c0})}{\prime_{c0}^{2}} ; \qquad (11)$$

where  $'_{c0}$   $'_0$  ( $T_c$ ) is the jump of the dimensionless order parameter at  $T_c$ . The second term in s can be neglected. In fact, taking into account the equation  $f['_0(T_c)] = 0$  for the equilibrium phase transition point  $T_c$  we obtain that  $jC I ('_0) = '_0^2 j$  is approximately equal to  $j_{c0} + '_{c0}^2 = 2j$  where  $'_{c0}^2$  and the dimensionless shift of the transition temperature  $t_{c0} = t_0 (T_c)$  are expected to be much smaller than unity. The latent heat  $Q = T_c s(T_c)$  and the jump of the speci c heat capacity at  $T_c$ ,  $C = T_c (@ S=@T)$  can be easily calculated with the help of Eq. (11). For this purpose we need the function  $'_0(T)$ , which cannot be obtained analytically from Eq. (7).

Even the relatively simple 2D-3D crossover scheme outlined in this paper cannot be used for an analytical calculation of the superconducting order parameter ', T) and related therm odynam ic quantities. But the sim plicity of the scheme makes it very convenient for a num erical calculation of experim entally observable quantities in thin superconducting In s. Our num erical treatment of Eqs. (5) and (7) shows a remarkably good con imation of the results for 3D [2] and quasi-2D [1,3] superconductors. Here we want to mention that the reliability of present crossover scheme can be tested also analytically by considering 1, which corresponds to the  $\3D$  lim it", and L<sub>0</sub> the cases:  $L_0$ 1, which gives the quasi-2D lim it". A fiter taking the quasi-2D lim it in Eqs. (5) and (7), we obtain the preceding results for  $\prime_0$  (T) and the free energy f (') reported in Refs. [1, 3]. The results in the 3D lim it of Eqs. (5) and (7) should be compared with the results for the free energy and the equation of state obtained when the continuum limit  $(k_k + k_k)$  in the sum (3) is taken, which means to extend the integration to  $_0 = 1$  . In fact, we have done the calculations in the geometry of an \in nite cylinder" that corresponds to our nite-size treatment. A Itematively, the 2D-3D crossover can be described in the usual \spherical" geometry, when the cuto s and  $_0$  are equal both in the sum (3) and in the respective integral, calculated in the continuum limit ( $L_0$  ! 1) along the smalldimension  $L_0$ . The equations for the free energy and the equation of state in this variant of the theory are much more complicated and less convenient for a num erical evaluation of experim entally observable therm odynam ic quantities.

A cknow ledgem ents. The authors thank Prof. D. I. U zunov for useful discussions. A funding by the European Superconductivity Network (Scenet) is acknow ledged.

## References

- [1] R.Folk, D.V. Shopova and D.I.U zunov, Phys. Lett. A 281 (2001) 197.
- [2] D.V. Shopova, T.P. Todorov, T.E. Tsvetkov and D.I.Uzunov, Mod. Phys. Lett. B 16 (2002) 829.
- [3] D.V.Shopova, T.P.Todorov, D.I.Uzunov, Mod. Phys. Lett. B 17 (2002) 1.
- [4] B.I.Halperin, T.C.Lubensky, and S.K.Ma, Phys. Rev. Lett. 32 (1974) 292.
- [5] R. Folk and Yu. Holovatch, in: Correlations, Coherence, and Order, ed. by D.V. Shopova and D.I.Uzunov (Kluwer Academic/Plenum Publishers, New York-London, 1999), p.83.
- [6] D.I.Uzunov, Theory of Critical Phenom ena (World Scientic, Singapore, 1993).
- [7] M.P.A.Fisher, G.Grinstein, Phys. Rev. Lett. 60 (1988) 208.
- [8] D.V.Shopova, D.I.Uzunov, Phys. Rep. C 279 (2003) 1.

- [9] D.Belitz, T.R.Kirkpatrick, Phys. Rev. Lett. 89 (2002) 247202.
- [10] L.Craco, L.DeCesare, I.Rabu o, I.P.Takov, D.I.Uzunov, Physica A 270 (1999) 486.
- [11] J.C.Rahola, J.Phys. Studies 5 (2001) 304.
- [12] E.M. Lifshitz and L.P.Pitaevskii, Statistical Physics, Part 2, [Landau and Lifshitz Course of Theoretical Physics, vol. 9] (Pergam on Press, Oxford, 1980).
- [13] T.P.Todorov, PhD Thesis, BAS, So a, 2003; unpublished.