LOCAL QUASI-EQUILIBRIUM DESCRIPTION OF SLOW RELAXATION

SY STEM S

I. Santam ar a-Holek, A. Perez-Madrid and J. M. Rub Department de F sica Fonamental Facultat de F sica Universitat de Barcelona Diagonal 647, 08028 Barcelona, Spain

W e present a dynam ical description of slow relaxation processes based on the extension of Onsager's uctuation theory to system s in local quasi-equilibrium. A non-M arkovian Fokker-P lanck equation for the conditional probability density is derived and from it, we obtain the relaxation equation for the moments. We show that the uctuation-dissipation theorem can be formulated in term s of the temperature of the system at local quasi-equilibrium which is related to that of the bath by m eans of a scaling factor revealing lack of therm al equilibrium . O ur theory m ay be applied to a wide variety of system s undergoing slow relaxation. W e discuss in particular slow dynam ics in glassy system s and B rownian m otion in a granular gas.

I. IN TRODUCTION

System s whose global evolution results from the existence of a wide variety of time and length scales frequently exhibit slow relaxation dynamics, which is manifested through the aging behaviour of the correlations and the non-exponential decay of the response function [1]-[3]. To characterize the dynamics of these systems is one of the most challenging problems of nonequilibrium statistical mechanics now adays. M any e orts have been m ade in this sense by proposing di erent approaches including spin m odels, stochastic dynamics and therm odynam ical approaches (see, for example, Refs. [4]-[13]).

Experiments performed in diverse systems as in an orphous polymers [14] and supercooled liquids and glasses [1], [15]-[17], seem to indicate that these systems undergo, in general, non-Markovian and non-stationary dynamics leading to the peculiar behaviour of the two-time correlation functions and to the dependence of the relaxation times on the initial time.

In this paper we present a dynam ical description of slow relaxation systems based on the form ulation of a Fokker-P lanck equation for non-M arkovian processes. This equation is obtained through the generalization of the Onsager theory, originally proposed to account for the dynam ics of uctuations around equilibrium states [18] and extended to nonequilibrium stationary states [19], to nonequilibrium aging states.

An important feature of the dynam ics of these systems is that at the time scales of experiments, they still relax and never reach thermal equilibrium with the bath. We have assumed the existence of a local quasi-equilibrium state [20] characterized by a non-stationary probability density which can be used to determ ine average quantities as, for example, the local quasi-equilibrium temperature [21]. The entropy of the system has been expressed in terms of the conditional probability density by means of the G ibbs entropy postulate, and used to derive the generalized FokkerP lanck equation by taking into account probability conservation and the rules of nonequilibrium therm odynamics [22]. For stationary processes, this equation was derived in [26] and [27], and has subsequently been treated in the literature of stochastic processes in [28]-[30].

The uni ed description of slow relaxation system s we propose can be applied to, am ong others, glassy system s [9], granularm atter [31] and anom alous di usion problem s [29], [32].

The paper is distributed as follows. In Sec. II, we will analyze the non-stationary dynamics of slow relaxation system s under the fram ework of a generalized O nsager theory. Section III is devoted to the form ulation of relaxation equations and of the generalized uctuation-dissipation theorem for non-stationary systems. An illustration of the theory presented is given in Sec. IV by analyzing the dynamics of systems having two well-separated time scales (glassy systems). In Sec. V, we will discuss the B row nian motion in a granular gas. W e will compare our results with those obtained by means of kinetic theory and simulations. In the discussion section we will summarize our main results.

II. FOKKER-PLANCK EQUATION FOR NON-STATIONARY DYNAM ICS

The non-stationary dynam ics of slow relaxation system s can be characterized at the m esoscopic level by m eans of the conditional probability density $P(_0;t_0;t_1;t)$, which satis es the norm alization condition $P(_0;t_0;t_1;t)d_{=}=1$. This quantity depends in general on a set of uctuating variables $_= (_1; ...; _n)$, or elds $_(\tilde{K})$, [33], necessary to univocally determ ine the state of the system during the relaxation and which m ay represent, for instance, uctuating positions and velocities, order parameters, reaction coordinates or hydrodynamic elds.

$$\frac{\underline{\theta}}{\underline{\theta}t}P(\underline{\}_{0};t_{0}\underline{i};t) = \frac{\underline{\theta}}{\underline{\theta}}P(\underline{\}_{0};t_{0}\underline{i};t)\underline{v}(\underline{\}_{0};\underline{i};t_{0};t);t)$$
(1)

expressing probability conservation during its natural evolution in _-space. The probability current $P(_0;t_0;t_0;t) \leq (_0;:t_0;t)$, with $v_{(_0;:t_0;t)}$; a stream velocity in _-space, has to be determined in order to render the dynam ical description complete.

This task can be accomplished by assuming that the scheme of nonequilibrium thermodynamics [22] remains valid in _-space. Thus, a linear relationship can be established between the di usion current P (t_0 <u>t</u>)<u>v</u> (t_0 <u>t</u>) and its conjugate force in _-space, which can be obtained from the entropy production calculated in the fram ework of 0 nsager's theory [22]. To derive the entropy production we will assume that, for su ciently long times t, the system reaches a local quasi-equilibrium state, characterized by the entropy per unit mass s_{qe} (t) and the probability density P_{qe} (<u>;</u>t), related form ally to the conditional probability through $\lim_{t \le 1} P(\underline{0}; t_0; \underline{1}; t) = P_{qe}$ (<u>;</u>t).

In accordance with [23], the variation of the entropy s_{qe} (t) satis es the G ibbs equation

$$\begin{array}{c} & Z \\ T (t) s_{qe} (t) = e_{qe} (t) & {}_{qe} (t) P (t)d_{}; \end{array}$$

where the tem perature T (t) of the system at local quasi-equilibrium will be assumed to be only a function of time, e_{qe} (t) is the mean internal energy, and $_{qe}$ (t) the chem ical potential at local quasi-equilibrium, which do not depend on the _'s. For simplicity we have om itted the explicit dependence of the probability density on phase variables. The entropy production corresponding to the di usion process in _-space can be computed by using the G ibbs entropy postulate [24]

$$s(t) = \frac{k_{\rm B}}{m} P(t_0 t) \ln \frac{P(t_0 t)}{P_{\rm qe}(t)} d_{\rm resc} + s_{\rm qe}(t);$$
(3)

where s(t) is the entropy of the system, k_B the Boltzm ann's constant and m the molecular mass. Notice that, according with mesoscopic nonequilibrium thermodynamics (MNET), the entropy s_{qe} (t) appearing in Eq. (3), constitutes a macroscopic quantity whose variations are governed by Eq. (2). The evolution in time of s_{qe} (t) can also be described with the MNET scheme by using the expression of Eq. (2), and the corresponding Eq. (3) referred to the equilibrium state (see, for example, [23] and [25]). In that case, by taking the variation $s_{qe} = s_{qe}$ (fP_{qe} + Pg) s_{qe} (fP_{qe}g) at rst order in P and combining the resulting expression with Eq. (2), we obtain

$$P_{qe}(t) = P_{e} e^{\frac{m}{k_{B} T(t)} \left[qe - e_{qe}\right]};$$
(4)

which incorporates the topology of the energy landscape through e_{qe} , and where P_e is the equilibrium probability distribution.

By combining the rate of change of s_{qe} (t) obtained from Eq. (2) with the time derivative of Eq. (3), after using Eq. (1) and integrating by parts assuming that P (t_0 ; t) v_ (t_0 ; t) vanishes at the boundary, we obtain [34]

$$\frac{\theta}{\theta t}s(t) = \frac{1}{T(t)}\frac{\theta}{\theta t}e(t) - \frac{1}{T(t)}^{Z}P(t_{0}t)\underline{v}(t_{0};t) - \frac{\theta}{\theta} - \frac{k_{B}T(t)}{m}\ln\frac{P(t_{0}t)}{P_{qe}(t)} + q_{e}(t) d_{;}$$
(5)

where $e(t) = e_{q_e}(t) + \frac{k_B T(t)}{m}^R P(t_0 t) \ln P_{q_e}(t)d_$ is the variation of the nonequilibrium internal energy e(t). If no external therm odynam ic forces as, for example, velocity or tem perature gradients act on the system, it may be shown by using the G ibbs-D uhem relation that e(t) = 0. As a consequence of this, the entropy production is only due to the underlying di usion process in _-space. M oreover, in the last term of Eq. (5) one may identify the nonequilibrium \chem icalpotential" $(t_0; t)$ as

$$(t_{0};t) = \frac{k_{B}T(t)}{m} \ln \frac{P(t_{0}t)}{P_{qe}(t)} + qe(t):$$
(6)

In portant to notice is the fact that $(t_0;t)$ m ay depend, in general, on interaction potentials or entropic barriers [35] through the corresponding dependence of $P_{qe}(t)$.

A coording to the rules of nonequilibrium therm odynam ics expressing currents as linear functions of the conjugate therm odynam ic forces [22], from the entropy production (5) the velocity $\underline{v}_{(t_0;t)}$ can now be expressed in terms of $\frac{\theta}{\theta_{(t_0)}}$ (t), obtaining

$$\underline{\mathbf{v}}_{(t_0;t)} = \underline{\mathbf{B}}_{(t_0;t)} (t_0;t) + \frac{\mathbf{k}_{\mathbf{B}} \mathbf{T}(t)}{\mathbf{m}} \frac{\mathbf{0}}{\mathbf{0}} \ln \mathbf{P}(t_0;t) ; \qquad (7)$$

where we have identi ed the generalized force [30], [22]

$$\underline{X}(\underline{;}t) = \frac{\theta}{\theta} \frac{k_{\rm B} T(t)}{m} \ln P_{\rm qe}(t) ; \qquad (8)$$

and introduced the coe cients $\underline{B}_{(t_0;t)}$ which may, in general, depend on the state variables _. These coe cients, incorporating memory elects through their time dependence, can be determined in terms of the conditional relaxation functions of the corresponding variables _, [22]. Notice that, in general, the \underline{B}_{ij} 's do not satisfy reciprocal relations since for non-stationary systems and non-conservative interactions, the time reversal symmetry and the principle of detailed balance are not necessarily full led. The time dependence of the \underline{B}_{ij} 's is the origin of anom alous di usion, which is characterized by the power law behaviour in time of the correlation function [29], [32], and the nonexponential time decay of the relaxation functions [1].

By substituting the expression of the probability current obtained from Eq. (7) into Eq. (1), we obtain the Fokker-P lanck equation

$$\frac{(e_{\rm P}(t_0 \pm))}{(e_{\rm T})} = \frac{(e_{\rm T})}{(e_{\rm T})} = \frac{(e_{\rm T})}{(e_{\rm T})} \frac{X_{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm B})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm B})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} + \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})}} \frac{(e_{\rm T})^{\rm T}(t_{\rm T})}{(e_{\rm T})} \frac{(e_{\rm T})^{\rm T}($$

Sim ilar equations describing stationary non-M arkovian dynam ics were obtained in Refs. [26]-[30], to study linear and nonlinear transport. From Eq. (9), it follows that the characteristic relaxation times of the system $B_{ij}^{1}(t_0;t)$, are functions of the waiting time t_0 as a consequence of the non-M arkovian nature of the dynam ics.

At the time scales considered, the system relaxes towards the time-dependent local quasi-equilibrium state and then them all equilibrium between system and bath can not be established. The complete description of the dynamics can then be performed by giving the relation between the system and bath temperatures. In order to derive it, the in unce of the bath is incorporated through the second law $s_T = s_{qe} + s_b = 0$, with so the total entropy of the system and s_B that of the bath. From this expression, by assuming that the variation s_T takes place at constant volume (see Eq. (2)), it can be expressed as $s_T = [1 \quad c_T^{\frac{1}{1}}(t) [e_{qe}(t) = g_e(t)]$, where we have taken into account the G ibbs equation (2), used the G ibbs free energy per mass unit $g_{qe}(t) = \frac{1}{q_e(t)} P(t)d_{-}$ and dened $c = -\frac{p}{-} = 1$, with $_B$ the entropy production of the bath. The equality occurs when the process is reversible, i.e., $_B = -$. In a similar way, by using the G ibbs equation of the bath we obtain $s_T = [1 \quad c_T^{-1}]_{T_B}(t) = e_B(t)$, with $e_B(t)$ and $T_B(t)$ the energy and tem perature of the bath. A simple comparison of the expressions for s_T leads to the relation

$$T(t) = AT_B(t);$$
 (10)

where we have de ned $A = c \frac{h_{q_{q_e}(t)} e_{q_e}(t)}{e_{b}}^{i}$. Since $g_{q_e}(t) = s_{q_e} T(t)_{q_e}$, we conclude that the factor A also depends on the density and the temperature of the system. When the process is reversible, A = 1 since therm al equilibrium between system and bath is attained. In this case, $P_{q_e}(t)$ reduces to a local equilibrium probability density in which the temperature is that of the bath [35], [36].

III. RELAXATION EQUATIONS AND THE FLUCTUATION-DISSIPATION THEOREM

Once the Fokker-P lanck equation has been derived, we will proceed to establish the relaxation equations for the moments of the probability density. The analysis of the general case can be accomplished by considering that the

coe cients are functions of the state variables [30]. For the sake of simplicity, we will focus on the case in which the coe cients B $_{ij}$ are only functions of time.

F irst, we will consider the two-time correlation function

$$Z C_{i \circ j}(t_0;t) = i \circ_j P(_0;t_0) P(_0;t_0;t_0;t_0) d_{j}; t) d_{0} d_{j}; t$$
(11)

whose evolution equation m ay be found by taking the time derivative of Eq. (11), substituting Eq. (9) and integrating by parts, obtaining

$$\frac{d}{dt}C_{i \ 0j}(t_0;t) = B_{i1}(t_0;t) hX_1(t)_{0j}(t_0)i;$$
(12)

where we have de ned hX_i(t) _{0j}(t₀)i $R_{X_i(j;t) _{0j}P(_0;t_0)P(_0;t_0j;t)d_0}d_0$. This equation in plicitly provides an operational de nition of the transport coe cients B_{ij}, showing aging e ects through their dependence on the waiting time. The relation between the dissipation of the energy and the response of the system to an external timedependent force_(t) = _(t t_0), with (x) being the step function, m ay be determined by m eans of the evolution equation for the rst m om ent of the probability density, h_i(t) i = $R_{iP}(_0;t_0)P(_0;t_0j;t)d_0d_0$. In the presence of the external force, the system is described by m eans of the corresponding Fokker-P lanck equation

$$\frac{\underline{\theta}P(t_0 \underline{t})}{\underline{\theta}t} = \frac{\underline{\theta}}{\underline{\theta}} = \frac{B}{\underline{\theta}} d\underline{t} t) \quad \underline{X}(\underline{t};t)P(t_0 \underline{t}) + \frac{k_B T_B(t)}{m} A \frac{\underline{\theta}P(t_0 \underline{t})}{\underline{\theta}} \\ \frac{\underline{\theta}}{\underline{\theta}} = [(t)P_0(\underline{t})]; \tag{13}$$

where the external force has introduced the additional drift _ (t). A coording to linear response theory β 7], the response R _ i _ i (t_0;t) of the system is then given through the functional derivative

$$R_{i}(t_{0};t) = \frac{h_{i}(t)i}{j(t_{0})};$$
(14)

Notice that, due to the time dependence of the coe cients B $_{ij}$ (t₀;t), the response function will contain modi cations to the M arkovian response.

The response function can be calculated from the equation for the $rst m om ent h_i(t)i which follows from the Fokker-P lanck equation (13). We obtain$

$$R_{i_{0}}(t_{0};t) = \frac{m}{k_{B}} \frac{1}{T(t)} \frac{\theta}{\theta t_{0}} C_{i_{0}}(t_{0};t); \qquad (15)$$

which constitutes the formulation of the uctuation-dissipation theorem. This expression indicates that the uctuation-dissipation theorem remains valid when the system is in local quasi-equilibrium; the temperature entering that expression is that of the system at local quasi-equilibrium. If one uses Eqs. (10) and (15), it can also be expressed as

$$R_{i0j}(t_0;t) = \frac{m}{k_B} \frac{A^{1}}{T_B(t)} \frac{Q}{Qt_0} C_{i0j}(t_0;t):$$
(16)

The result (16) can in general be interpreted as a violation of the uctuation-dissipation theorem [8] and [9]. When analyzing its equivalent form (15) it expresses validity of the uctuation-dissipation theorem in terms of the local quasi-equilibrium temperature.

The quantity A has been derived in Section II from them odynam ic grounds. An expression in term s of correlation functions can be obtained from the equal-time correlation function (t) $\frac{1}{2}^{R} - P_{(i,t_0)}P_{(0,i,t_0)}(t_0) = \frac{1}{2} + \frac{1$

$$\frac{d}{dt} \qquad (t) = B (t_0;t) h \underline{X} (t) - (t) i + \frac{k_B T (t)}{m} B (t_0;t); \qquad (17)$$

where for simplicity, we have considered $\underline{B}(t_0;t) = B(t_0;t)\underline{1}$, with $\underline{1}$ the unit tensor. For times t $(t_0;t)$, the time derivative of Eq. (17) may be neglected and the system relaxes to the reference state, leading to the expression

$$A = \frac{m}{k_{\rm B} T_{\rm B} (t)} \frac{h \underline{X} (t)}{L} (t) (t)_{\rm qe}^{\rm i};$$
(18)

where $h:::i_{qe}$ represents the average in respect to the local quasi-equilibrium probability density P_{qe} (t). This equation indicates that A m ay in general be expressed by m eans of the m on ents of the probability density. A coording to Eqs. (10) and (18), it follows that the tem perature T (t) entering the G ibbs equation (2) results from the contribution of the energies related to the corresponding degrees of freedom _ by

T (t)
$$\frac{m}{k_{\rm B}}$$
 hX (t) _ (t) i : (19)

An important fact to be noticed from this expression is that the temperature is not robust since it implicitly depends on the number of relevant variables or observables used to describe the system, as has been evidenced in [38]. Given that the probability density of the reference state depends on the initial condition, it follows that T (t) is also a function of t_0 . In the case when the force is linear, Eq. (19) reduces to T (t) = $\frac{m}{k_B}$ ² (t) _{qe}, which coincides with the corresponding result obtained when local therm odynam ic equilibrium between system and bath exists. In that case, when _= u with u the velocity of a particle, Eq. (19) becomes the kinetic de nition of the temperature.

IV. SLOW DYNAM ICS IN GLASSY SYSTEMS

As a rst application of the theory presented, we will analyze the dynam ics of systems characterized by two sets of variables: fast -variables and slow -variables. The -variables relax to a local equilibrium state characterized by the tem perature of the bath T_B , whereas the -variables relax to a quasi-equilibrium state characterized by the tem perature T (t).

The probability density is now a function of the state vector $_ = (_; _)$. Since -variables equilibrate with the bath, we may assume that the probability density characterizing the reference state can be factorized in the form $P_{qe}(t) = P_{le}(_)P_{qe}(_;t)$, with $P_{le}(_)$ and $P_{qe}(_;t)$ stationary and non-stationary distributions. Consequently, $s_{qe}(t) = s_{le} + s_{qe}(t)$, where $T_B = s_{le}$ and $T(t) = s_{qe}$ enter G ibbs equations similar to Eq. (2). Consistent with the analysis of Sec. II, the probability densities of the reference states are given by

$$P_{le}(\underline{}) = e^{\frac{m}{k_{B}T_{B}}\left[l_{e} e_{le}(\underline{})\right]} \text{ and } P_{qe}(\underline{};t) = P_{e}e^{\frac{m}{k_{B}T(t)}\left[q_{e} e_{qe}(\underline{})\right]};$$
(20)

By taking the time derivative of Eq. (3) and using Eq. (20) and the corresponding continuity equation, the entropy production can be written as

$$= P(t_0 \ddagger) \underline{v}(\underline{t}) \frac{e}{e_{\perp}} (t_0; t) d_{\underline{t}}$$
(21)

where we have de ned the nonequilibrium Massieu function

$$(t_{0};t) \quad \frac{k_{B}}{m} \ln P (t_{0};t) + \frac{e_{le}(\underline{})}{T_{B}} + \frac{e_{qe}(\underline{})}{T (t)}; \qquad (22)$$

which is the appropriate function in the entropy representation [39], [40].

Since $\underline{v}(\underline{;}t) \quad \underline{v}; \underline{v}$ and $\frac{\underline{\theta}}{\underline{\theta}} = \frac{\underline{\theta}}{\underline{\theta}}; \frac{\underline{\theta}}{\underline{\theta}}$, from the entropy production (21) we can obtain the expressions for \underline{v} and \underline{v} (see Eq. (7)). In general, the rules of nonequilibrium therm odynamics introduce couplings between the velocities \underline{v} and \underline{v} and the corresponding forces through the mobility \underline{B} which is in general a function of $t_0, t, _$ and $_$.

In this case, since no couplings appear due to the separation of time scales, the mobility matrix \underline{B} can be written as the direct sum of two submatrices \underline{B} and \underline{B} . For simplicity, we will assume that the B_{ij} 's are constant since the -variables decay to a local equilibrium state [22]. The B_{ij} 's will be functions of two times, $B_{ij} = B_{ij}$ (t₀; t).

Now, by taking into account that -variables them alize at T_B and -variables at T (t), [11]-[13], after substituting the expressions for \underline{v} and \underline{v} into the continuity equation one obtains the generalized Fokker-P lanck equation

$$\frac{(e_{P} (t_{0} \pm))}{(e_{t})} = \frac{(e_{t})}{(e_{t})} = \frac{B(t_{0};t) \times (c_{0})P(t_{0}\pm) + \frac{k_{B}T(t)}{m} \frac{(e_{P} (t_{0}\pm))}{(e_{t})} + \frac{k_{B}T_{B}}{(e_{t})} \frac{(e_{P} (t_{0}\pm))}{(e_{t})} + \frac{k_{B}T_{B}}{m} \frac{(e_{P} (t_{0}\pm))}{(e_{t})} ;$$

$$(23)$$

where, according with Eqs. (8) and (20), \underline{X} (_) = $\frac{@e_{le}(\underline{)}}{@}$ and \underline{X} (_) = $\frac{@e_{qe}(\underline{)}}{@}$.

O ne may obtain the Fokker-Planck equation for the slow -variables by multiplying Eq. (23) by P $(__0;t_0)$ and integrating over $__0$ and $_$ taking into account that the current P $(t_0;t)\underline{v}$ vanishes at the boundaries. Our result is Eq. (9). In similar form, the evolution equation for the -variables can be derived by integrating over $__0$ and $_$. The resulting Fokker-P lanck equation is

$$\frac{@P(\underline{;t)}}{@t} = \frac{@}{@} = B \quad \underline{X} \quad (\underline{)}P(\underline{;t}) + \frac{k_B T_B}{m} \frac{@}{@}P(\underline{;t}) ; \qquad (24)$$

where P (;t) is the single-time distribution function.

The computation of the moments of the probability density following the procedure indicated in Sec. III, allows one to nd the statistical expression of the temperatures

$$T_{B} = \frac{m}{k_{B}} \frac{D}{X} (\underline{)} = \frac{E}{k_{B}}; \qquad (25)$$

and

$$T(t) = \frac{m}{k_{\rm B}} h \underline{X} \quad (\underline{t}; t) = \frac{i}{q_{\rm e}} t$$
(26)

Eq. (25) in plies that the fast degrees of freedom satisfy the equipartition law and its energy is a quadratic function in 's, whereas (26) is a de nition of the tem perature of the system in terms of the correlations at the quasi-equilibrium state.

>From Eq. (23), we may compute the response of the system to an external perturbation and derive the uctuationdissipation theorem related to both sets of variables. Following the procedure indicated in Sec. III, the response functions of the fast variables satisfy the relation

$$R_{i_{0}}(t_{0};t) = \frac{m}{k_{B}} \frac{1}{T_{B}} \frac{\theta}{\theta t_{0}} C_{i_{0}}(t_{0};t); \qquad (27)$$

where $R_{i_{0}j}(t_{0};t) = \frac{h_{i}(t)i}{j(t_{0})}$. The corresponding expression for the response of the -variables is given in Eq. (15). From Eq. (27) it follows that the fast degrees of freedom satisfy the uctuation-dissipation theorem in terms of the tem perature of the bath.

V. BROWNIAN MOTION IN A GRANULAR GAS

As a second application of our form alism, we will discuss in this section the Brownian motion of a grain in a granular gas, which has been previously addressed in [31] by means of kinetic theory for systems with inelastic collisions. The phase-space vector is now $_ = (u; r)$, where u and r are the velocity and position of the Brownian grain. The corresponding continuity equation for P (t₀ t) is given by

$$\frac{\partial \mathbb{P}(t_0 \pm)}{\partial t} + r \qquad \forall \mathbb{P}(t_0) = \frac{\partial}{\partial u} \qquad \mathbb{P}(t_0) \mathsf{v}_u(t_0; t)]; \tag{28}$$

$$P_{qe}(t) = e^{\frac{m}{k_B T(t)} \left[qe - \frac{1}{2}u^2\right]};$$
(29)

where $_{qe}$ normalizes P_{qe} (t).

Once the P_{qe} (t) has been specified, the Fokker-P lanck equation for the B row nian gas m ay be derived by using Eqs. (3), (28), (29). Following the steps indicated in Sec. II, we nally obtain the generalized Fokker-P lanck equation

$$\frac{(e_{\rm P}(t_0)\pm)}{(e_{\rm T})} + r \qquad e_{\rm P} = \frac{(e_{\rm T})}{(e_{\rm T})} \qquad B_{\rm I}(t_{\rm T}) \qquad e_{\rm P}(t_{\rm I})\pm) + \frac{k_{\rm B}T_{\rm B}(t)}{m} A \frac{(e_{\rm P}(t_{\rm I})\pm)}{(e_{\rm T})}; \qquad (30)$$

where we have assumed locality in u-space for which the velocity v_u (t_0 ;t) couples only to the gradient of the chem ical potential at the same value of u in accordance with Eq. (5). By considering the action of an external force eld ~ (t) which can be an impulsive one (tapping force), we may derive by following the steps indicated in Sec. III, the uctuation-dissipation relation

$$R_{uu_0}(t_0;t) = \frac{m}{k_B} \frac{A^{-1}}{T_B(t)} \frac{\theta}{\theta t_0} C_{uu_0}(t_0;t):$$
(31)

Recent computer simulations of di usion in dilute granular gases con m the validity of Eq. (31), [41]. The corresponding integrated version of the generalized uctuation-dissipation theorem has been also obtained in dense granular systems in [42]. Now, by comparing Eq. (30) with the corresponding one found in [31] by following kinetic theory m ethods, we may identify

$$A = \frac{(1 +)}{2} ; B (t_0; t) = \frac{1}{2} (1 +) e (t_0; t);$$
(32)

where the restitution coe cient (1) of the B row nian grains has been assumed constant, and $_{e}(t_{0};t)$ is the friction coe cient for the case when elastic collisions take place, [31].

>From the Fokker-P lanck equation (30), it follows that the evolution equation of the second moment of the velocity of the B rownian grains can be expressed in the form

$$\frac{d}{dt}T(t) = 2B(t_{B};t)[T(t) A T_{B}(t)]:$$
(33)

which is a particular case of Eq. (17). Eq. (33) establishes that the relation between system and bath temperatures at the local quasi-equilibrium state given by Eq. (10), is valid for times t B^{-1} (t₀;t).

The relation between T (t) and T_B (t) can be generalized in order to describe shorter time scales by incorporating the therm al relaxation of the heat bath through the evolution equation $\frac{d}{dt}T_B$ (t) = (t) T_E (t), with (t) the cooling rate. A ssum ing that 0 < (t) < 2B (t₀;t), the time evolution of $\frac{T}{T_B}$ (t) is then given by [31]

$$\frac{d}{dt}\frac{T(t)}{T_{B}(t)} + 2(1) B(t;t)\frac{T(t)}{T_{B}(t)} = 2B(t_{0};t)A;$$
(34)

where we have de ned $\frac{(t)}{2B(t_0,t)}$. From Eq. (34), we then obtain

$$T(t) = \frac{A}{(1)} T_{B}(t);$$
 (35)

which gives the relation between the system and bath tem peratures for times t $[(1) B(t;t)]^{1}$.

By contracting the description, we enter the di usion regime described by the density $(\mathbf{r}^0; \mathbf{t}_0; \mathbf{r}_0; \mathbf{t}_0; \mathbf{r}_0; \mathbf{t}_0; \mathbf{r}_0; \mathbf{t}_0; \mathbf{r}_0; \mathbf{t}_0; \mathbf{r}_0; \mathbf{t}_0; \mathbf{r}_0; \mathbf{t}_0; \mathbf{t}_$

$$\frac{\theta}{\theta t} (t_0 t) = D_{eff} (t_0; t) r^2 (t_0 t);$$
(36)

where D_{eff} (t₀;t) is the elective di usion coe cient given by

$$D_{eff}(t_{0};t) = \frac{k_{B} T_{B}(t)}{m B(t_{0};t)} \frac{A}{(1 - \frac{2}{3})};$$
(37)

which has been obtained by taking into account Eqs. (34) and (35). Eq. (37) was derived in [31] by m eans of kinetic theory. In particular, when the cooling rate of the bath satisfies (t) 2B, the expression of the diffusion coelected (37) reduces to $D_{eff}(t_0;t) = \frac{k_B T_B}{m B(t_0;t)} A$. In the general case, the dynamics of the system may be described by m eans of the corresponding Fokker-Planck equation in which the probability density of the reference state has the form $P_{qe}(t) = e^{\frac{m}{k_B T_B(t)} A^{-1} (1 -) [qe^{-\frac{1}{2}u^2}]}$, and the coelecters A and B (t₀;t) satisfy m ore general relations than Eq. (32).

The Fokker-Planck and Sm oluchow skiequations derived previously may also describe di usion in a colloidal system. In that case, the di usion coe cient (37) accounts for the violation of the Stokes-Einstein relation reported in dense granular system s [42] and supercooled colloidal uids [17].

VI. DISCUSSION

In this paper, we have proposed an extension of Onsager's theory, originally form ulated to describe nonequilibrium aged systems, in order to analyze slow relaxation dynamics. Our theory is based on the assumption of the existence of a local quasi-equilibrium state in phase space characterized by a non-stationary probability density. This hypothesis entails the existence of a time-dependent temperature of the system T (t) rejecting lack of therm all equilibrium between system and bath, which enables us to form ulate a G ibbs equation. The relation between the local quasi-equilibrium and bath temperatures was given in terms of a scale function obtained by taking into account the in uence of the bath.

The entropy production derived from the G ibbs entropy postulate leads to the explicit expression of the probability current in phase space and, consequently, to the generalized Fokker-P lanck equation (9). Memory elects were incorporated in the description through the time dependence of the coel cients entering into that equation. They are responsible for the dependence of the characteristic relaxation times on the initial (waiting) time. Our form alism could straightforwardly be extended to the case of a description in terms of elds _(k), [33], [44].

We have shown that the uctuation-dissipation relation remains valid in term softhe local quasi-equilibrium tem perature T (t) of the system whose value depends on the time scale considered. The fact that the local quasi-equilibrium tem perature depends on time and on the number of variables relevant to characterize the state, clari es why two system s with slow relaxation dynamics put in therm al contact do not necessarily satisfy a \zero law".

As a rst application, we have analyzed the slow dynamics of glasses. We have shown that the dynamics of the system can be described by a generalized Fokker-P lanck equation governing the time evolution of both the fast and slow degrees of freedom. By using this equation, we have veried that the uctuation-dissipation theorem (27) is satisfied by the fast modes. For the slow modes the theorem (15) is valid in terms of the local quasi-equilibrium temperature.

We have also studied the problem of Brownian di usion in a dilute granular gas reproducing the Fokker-Planck and Sm oluchow skiequations derived previously by means of kinetic theory of granular gases. These equations may also be used to describe the dynamics of supercooled colloidal suspensions leading to a generalization of the Stokes-Einstein relation.

The theory presented o ers a uni ed description of system s undergoing slow relaxation and constitutes a generalization of the O nsager theory describing the evolution of uctuations around nonequilibrium aging states.

VII. ACKNOW LEDGMENTS

W e want to acknow ledge D r. D. Reguera for valious com m ents. IS H. acknow ledges UNAM -DGAPA for econom ic support. This work was partially supported by DGICYT of the Spanish G overnm ent under G rant N o. PB 2002-01267.

- [1] M.D.Ediger, C.A.Angell and S.R.Nagel, J.Phys.Chem. 100, 13200 (1996).
- [2] P.G.Debenedettiand F.H.Stillinger, Nature, 410, 259 (2001).
- [3] F.H.Stillinger, Phys. Rev. B 41, 2409 (1990).
- [4] R.G. Palmer, D.L. Stein, E. Abrahams and P.W. Anderson, Phys. Rev. Lett. 53, 958 (1984).
- [5] A.J.Bray and G.J, Rodgers, Phys. Rev. B 38, 11461 (1988).
- [6] G.Parisi, Phys. Rev. Lett. 79, 3660 (1997).
- [7] H.Som polinski and A.Zippelius, Phys. Rev. Lett. 47, 359 (1981).
- [8] L.Cugliandolo and J.Kurchan, Phys. Rev. Lett. 71, 173 (1993).
- [9] L. Cugliandolo, J. Kurchan and L. Peliti, Phys. Rev. E 55, 3898 (1997).
- [10] W .G otze and L.Sjogren, Rep.Prog.Phys.55, 241 (1992).
- [11] S.Mossa, et al, Phys.Rev.E 65, 041205 (2002).
- [12] E. la N ave, et al, cond-m at/0303040
- [13] Th.M.Nieuwenhuizen, Phys.Rev.E 61, 267 (2000).
- [14] L.C. E. Struick, Physical aging in am orphous polymers and other materials, Elsevier (1978).
- [15] T.S.G rigera and N.E. Israelo, Phys. Rev. Lett. 83, 5038 (1999).
- [16] L.Bellon ans S.Ciliberto, Physica D 168-69, 325 (2002).
- [17] D.Bonn and W.K.Kegel, J.Chem.Phys.118, 2005 (2003).
- [18] L.Onsager, Phys. Rev. 38, 2265 (1931).
- [19] J.R.Dorffnan, T.R.Kirkpatrick and J.V.Sengers, Annu.Rev.Phys.Chem. 45, 215 (1994).
- [20] S.Franz and M.A.Virasoro, J.Phys.A:Math.Gen. 33, 891 (2000).
- [21] R. Zwanzig, Nonequilibrium statistical mechanics (Oxford University Press Inc., Oxford, 2001).
- [22] S.R.de Groot and P.M azur, Non-equilibrium thermodynamics (Dover, New York, 1984).
- [23] J.M. Rub, A. Perez-Madrid, Physica A 264, 492 (1999).
- [24] N.G. van Kampen, Stochastic processes in physics and chemistry (North-Holland, Amsterdam, 1981).
- [25] P.Mazur, Physica A 274, 491 (1999).
- [26] S.A.Adelman, J.Chem. Phys. 64, 124 (1976).
- [27] M. Tokuyam a and H. Mori, Prog. Theor. Phys. 55, 411 (1976).
- [28] S.Okuyam a and D.W. Oxtoby, J.Chem. Phys. 84, 5824 (1986).
- [29] K.G.W ang and M. Tokuyam a, Physica A 265, 341 (1999).
- [30] I.Santam ar a-Holek and J.M.Rub, Physica A 326, 384 (2003).

- [31] J.J.Brey, A.Santos and J.W. Duffy, J.Stat. Phys. 97, 281 (1999).
- [32] R.Morgado, F.A.Oliveira, G.G.Batrouni and A.Hansen, Phys. Rev. Lett. 89 100601 (2002).
- [33] J.M. Rub and P.M azur, Physica A 276, 477 (2000).
- [34] A.Perez-Madrid, J.M.Rub, P.Mazur, Physica A 212, 231 (1994).
- [35] D.Reguera and J.M.Rub, Phys.Rev.E 64,061106-1 (2001).
- [36] L. Landau and E. M. Lifshitz, Course of Theoretical Physics Vol 5 (Statistical Physics Part 1) (Pergam on Press, New York, 1980).
- [37] R.Kubo, M.Toda and N.Hashitsum e, Statistical Physics II.Nonequilibrium Statistical Mechanics (Springer-Verlag, Berlin, 1985).
- [38] A. Perez-Madrid, D. Reguera and J. M. Rub, Physica A, in press.
- [39] H.Callen, Thermodynamics and an introduction to therm ostatistics (W iley, New York, 1985).
- [40] J.M.Rub, P.Mazur, Physica A 250, 253 (1998).
- [41] A. Puglisi, A. Baldassarri and V. Loreto, Phys. Rev. E 66, 061305 (2002).
- [42] H.A.Makse and J.Kurchan, Nature 415, 614 (2002).
- [43] I.Santam ar a-Holek, D.Reguera and J.M.Rub, Phys.Rev.E 63 051106 (2001).
- [44] A.Perez-M adrid, D.Reguera and J.M.Rub, J.Phys.: Condens. Matter 14, 1651 (2002).