Colloidal glass transition: beyond mode-coupling theory

Grzegorz Szam el

Department of Chemistry, Colorado State University, Fort Collins, CO 80525

(D ated: M arch 22, 2024)

A new theory for dynam ics of concentrated colloidal suspensions and the colloidal glass transition is proposed. The starting point is the mem ory function representation of the density correlation function. The mem ory function can be expressed in terms of a time-dependent pair-density correlation function. A n exact, form al equation of motion for this function is derived and a factorization approximation is applied to its evolution operator. In this way a closed set of equations for the density correlation function and the mem ory function is obtained. The theory predicts an ergodicity breaking transition similar to that predicted by the mode-coupling theory, but at a higher density.

PACS num bers: 82.70 D d, 64.70 P f, 61.20 Lc

There has been a bt of interest in recent years in the theoretical description of dynam ics of concentrated suspensions and the colloidal glass transition [1]. It has been stimulated by ingenious experiments which provide detailed inform ation about m icroscopic dynam ics of colloidal particles [2]. Due to the abundance of experimental data the colloidal glass transition has emerged as a favorite, m odel glass transition to be studied [3].

One of the conclusions of these studies is the acceptance of the mode-coupling theory (MCT) as the theory for dynamics of concentrated suspensions and their glass transition [4]. Historically, this is somewhat surprising since MCT was rst formulated for simple uids with Newtonian dynamics [5] and only afterwards was adapted to colloidal system s with stochastic (Brownian) dynamics [6]. On the other hand, basic approximations of MCT are less severe for Brownian systems [7].

MCT is a theory for correlation functions of slow variables, i.e. variables satisfying local conservation laws. For Brownian system s there is only one such variable: localdensity. MCT's starting point is the memory function representation of the density correlation function [8, 9]. The memory function is expressed in terms of a timedependent pair-density (i.e. four-particle) correlation function evolving with so-called projected dynam ics. For Brownian system s this step is exact [10]. The central approximation of MCT is the factorization approximation in which the pair-density correlation function is replaced by a product of two tim e-dependent density correlation functions. As a result one obtains a closed, nonlinear equation of motion for the density correlation function. This equation predicts an ergodicity breaking transition that is identied with the colloidal glass transition. MCT has also been used to describe, e.g., linear viscoelasticity [11], dynam ics of sheared suspensions [12], and colloidal gelation [13]. By and large, its predictions agree with experim ental and sim ulational results [4, 14].

In spite of these successes, M C T's problem s are well known [4]. The most important, fundam ental problem is that once the factorization approximation is made there is no obvious way to extended and/or improve the theory. This is most acute for Brownian systems because there the density is the only slow mode and thus couplings to other modes cannot be invoked! Furthermore, MCT system atically overestim ates so-called dynamic feedback e ect. Thus, e.g., it underestim ates the glass transition volume fraction for a Brownian hard-sphere system (by about 10% [4]) and overestimates the glass transition temperature for a Lennard-Jones mixture (by a factor of 2 [15]). Finally, MCT cannot describe slow dynamics in systems without static correlations [16].

A way to improve upon MCT would be to introduce m any-particle dynam ic variables into the theory. Such an attempt has been made for simple uids [17]; it was argued that these variables (essentially, pair-density uctuations) describe clusters of correlated particles. Unfortunately no quantitative results have been reported based on this interesting approach.

W e propose a di erent way to go beyond M C T.R ather than factorizing the pair-density correlation function, we derive an exact, form alequation of m otion for it [18]. The structure of this equation is very sim ilar to that of the equation of motion for the density correlation function; \pair" analogues of the usual frequency matrix and the irreducible m em ory function can be identi ed. The basic approximation of our theory is a factorization of the evolution operator of the pair-density correlation function. After this approximation we obtain a closed system of equations of motion for the density correlation function and the m em ory function. These equations predict an ergodicity breaking transition; for a Brownian hard sphere system the glass transition volume fraction, q, is equal to :549 (note that $\frac{M}{q}^{CT} = :525$, $\frac{exp}{q}$:58).

O ur theory is similar to M C T in that it relies upon an uncontrollable factorization approximation. In contrast to M C T, it uses this approximation one step later. Thus, e.g., our theory preserves the memory function representation of the pair-density correlation function while M C T approximates the latter by a product of two density correlation functions. However, as usual in the liquid state theory, a priori these features do not guarantee the superiority of our approach as compared to M C T. O ur theory starts from the m em ory function representation of the density correlation function, F(k;t),

$$F(k;t) = \frac{1}{N} \ln(k) \exp(t) n(k)i:$$
 (1)

Here N is the number of particles, n (k) is the Fourier transform of the density, n (k) = ${}_{1}e^{ik r_{1}}$; and is the N -particle evolution operator, i.e. the Sm oluchow skioperator, = $D_{0} {}_{1}\frac{\theta}{\theta r_{1}} {}_{1}\frac{\theta}{\theta r_{1}}$ F₁ [19], with D_{0} being the di usion coe cient of an isolated B row nian particle, = 1= (k_B T), and F₁ a force acting on particle 1. Finally, h:::i denotes the canonical ensemble average; the equilibrium distribution stands to the right of the quantity being averaged, and all operators act on it as well as on everything else. U sually, the m em ory function representation of the Laplace transform of the density correlation function, F (k;z), is written as [20]

$$F(k;z) = \frac{S(k)}{z + \frac{D_0k^2}{S(k)(1+M_0(k;z))}}$$
(2)

where S (k) is the static structure factor and M (k;z) is the Laplace transform of the irreducible memory function. We re-write (2) in a form that will allow us to identify the pair analogues of the frequency matrix and the memory function. We write a memory function expression for the Laplace transform (LT) of $F_{k;t}$)

$$LT (F-(k;t)) = k 1 + M (k;z)O^{1} O k$$
$$\frac{1}{hn (k)n (k)i} F (k;z):$$
(3)

Here 1 denotes a unit 3d tensor, O is de ned through k = m(k) n(k)i (note that $O = \frac{1}{2}N$), and M(k;z) is the Laplace transform of the current correlation function evolving with projected dynamics,

$$M (k;t) = j(k) \exp(irrt)j(k)$$
 : (4)

where j(k) is a projected current density,

$$j(k) = \hat{Q}_n D_0$$
 ($ik + F_1)e^{ik r_1}$: (5)

In Eq. (5) $\hat{Q}_n = 1$ \hat{P}_n , and \hat{P}_n is a projection operator on the density subspace,

$$\dot{P}_{n} = ::: \prod_{q}^{X} n(q) \frac{1}{\ln (q)n(q)i} n(q) :::: (6)$$

Finally, in Eq. (4) irr is the \one-particle irreducible Sm oluchow ski operator" [20],

$$^{\mathrm{irr}} = \hat{Q}_{n} \overset{\mathrm{X}}{\underset{1}{\overset{\mathrm{e}}{\underset{1}{\overset{\mathrm{e}}{\operatorname{er}}}}} \hat{Q}_{1} \qquad \frac{\underline{e}}{\underline{e}r_{1}} \qquad F_{1} \quad \hat{Q}_{n}; \qquad (7)$$

where $\hat{Q}_1 = 1$ \hat{P}_1 , and the projection operator \hat{P}_1 reads

$$\hat{P}_{1} = ::: q^{q} e^{iq r_{1}} ihe^{iq r_{1}} :::: (8)$$

To make connection with the usual form of the memory function representation we note that k O $k=\ln(k)n(k)i = D_0k^2=S(k)$ is the frequency matrix and $\hat{k} = M(k;z)O^1$ $\hat{k} = M(k;z)$; where $\hat{k} = k=k$, is the irreducible memory function.

To obtain a convenient expression for M (k;t) in terms of a pair-density correlation function we use the following exact [21] equality:

$$j(k) = \begin{array}{c} X & X \\ n_2(k_1; k_2) \\ (k_1; k_2; k_3; k_4) \\ g(k_1; k_2; k_3; k_4) hn_2(k_3; k_4) j(k) i: (9) \end{array}$$

Here $n_2 (k_1; k_2)$ is the part of pair-density uctuations orthogonal to the one-particle density uctuations,

$$n_{2}(k_{1};k_{2}) = \hat{Q}_{n} \overset{X}{\underset{l \in m}{=}} e^{ik_{1}r_{1}ik_{2}r_{m}} :$$
(10)

Furthermore, in Eq. (9) the sum s over $k_i < k_{i+1}$ are understood and g denotes the inverse pair-density uctuationsmatrix (it is a pair analogue of 1= hn (k)n (k)i),

$$X g(k_1;k_2;k_3;k_4) m_2(k_3;k_4)n_2(k_5; k_6)i = (k_3;k_4) k_1;k_5 k_2;k_6:$$
(11)

U sing identity (9) we can express mem ory function (4) in terms of the time-dependent pair-density correlation function evolving with one-particle irreducible dynam ics,

$$F_{22} (k_1; k_2; k_3; k_4; t) = n_2 (k_1; k_2) \exp(\frac{irr}{t}) n_2 (k_3; k_4) : (12)$$

Rather than factorizing F_{22} , we use the projection operatorm ethod to derive an exact, form alequation of motion for this function. The derivation will be given elsewhere [22]; here we present the structure of the nal form ula for the Laplace transform of the time-derivative of the pair-density correlation function, F_{22} ,

$$LT F_{22}(k_{1};k_{2};k_{3};k_{4};t) = \begin{pmatrix} 0 & 1 \\ Y^{7} & X \\ 0 & A & (k_{1};k_{2}) & I + M & (k_{5};k_{6};k_{7};k_{8};z)O & (k_{7};k_{8};k_{9};k_{10})^{1} \\ i = 3 & (k_{21} - 1;k_{21}) \\ O & (k_{9};k_{10};k_{11};k_{12}) & k_{11} \\ k_{12} & g & (k_{11};k_{12};k_{13};k_{14})F_{22} & (k_{13};k_{14};k_{3};k_{4};z) \end{pmatrix}$$
(13)

In Eq. (13) I denotes a unit 6d tensor, O and M are block matrices, e.g.

$$O (k_{1};k_{2};k_{3};k_{4}) = O_{11} (k_{1};k_{2};k_{3};k_{4}) O_{12} (k_{1};k_{2};k_{3};k_{4}) O_{21} (k_{1};k_{2};k_{3};k_{4}) O_{22} (k_{1};k_{2};k_{3};k_{4}) ; (14)$$

and the following short-hand notation is used:

$$(k_1;k_2) \circ (k_1;k_2;q_1;q_2) = \begin{cases} q_1 \\ q_2 \end{cases} = \begin{cases} X \\ k_1 & Q_j \\ i;j \end{cases} k_1 = (15)$$

O and M are the pair analogues of O and M (com pare Eqs. (3) and (13)); in particular

$$(k_1;k_2) \circ (k_1;k_2;k_3;k_4)$$

 $k_1 = n_2 (k_1;k_2)$ $irr n_2 (k_3; k_4) ;$ (16)

and M $_{ij}$ are pair-current correlations evolving with a two-particle irreducible evolution operator 2irr , e.g.,

$$M_{11} (k_1; k_2; k_3; k_4; t) = j_2 (k_1; k_2) \exp(^{2irr} t) j_2 (k_3; k_4)$$
(17)

where, e.g.,

$$\dot{\underline{j}}(k_3; k_4)\dot{\underline{i}} = \hat{Q}_{n_2} D_0 \frac{X}{\underset{l \in m}{\underline{0}}} \frac{\underline{\theta}}{\underline{\theta} r_1} \hat{Q}_1 e^{ik_3 r_1 + ik_4 r_m}$$
 :
(18)

Explicit form ulae for O and M (including de nitions of 2irr and \hat{Q}_{n_2}) will be given elsewhere [22].

The main approximation of our theory is factorization of the evolution operator for F_{22} . W ithin this approximation the diagonal blocks of O and M are given by

$$O_{11} (k_1; k_2; k_3; k_4) = O_{22} (k_2; k_1; k_4; k_3) =$$

$$N OS (k_2)_{k_1; k_3} k_{2; k_4};$$

$$M_{11} (k_1; k_2; k_3; k_4; t) = M_{22} (k_2; k_1; k_4; k_3; t) =$$

$$N M (k_1; t) F (k_2; t)_{k_1; k_3} k_{2; k_4};$$
(20)

and the o -diagonal ones vanish. Consistently, we also factorize g and F_{22} (t = 0).

U sing (19(20) we can express F_{22} in terms of the density correlation function and the memory function (note that F_{22} does not factorize for t > 0). Substituting F_{22} into the form ula for the memory function and using convolution approximation for static vertices [5, 6] we get

$$M(k;z) = \frac{nD_0}{2}^{Z} \frac{dk_1 dk_2}{(2)^3} (k + k_1 + k_2)^{L} \hat{k} + (c(k_1)k_1 + c(k_2)k_2)^{L} \frac{h}{z + \frac{D_0(k_1^2 = S(k_1))}{1 + LT(M(k_1;t)F(k_2;t) = S(k_2))} + (1 \leq 2)^{L}} \hat{k}$$
(21)

where n is the density and c(k) is the direct correlation function. Eqs. (2) and (21) determ ine time dependence of density correlations and the memory function.

Eqs. (2) and (21) predict an ergodicity breaking transition. In the non-ergodic regime F (k;t) has a nonzero long-time limit, $\lim_{t \ge 1} F(k;t) = f(k)S(k)$, where f(k) is called a non-ergodicity parameter. It follows from Eq. (2) that in this regime also the memory function has a non-zero long-time limit, $\lim_{t \ge 1} M(k;t) = m(k)D_0k^2=S(k)$, and that f(k) and m(k) are related by

$$\frac{f(k)}{1 - f(k)} = m(k):$$
 (22)

U sing (21-22) we get a self-consistent equation for f (k):

$$\frac{f(k)}{1 f(k)} = \frac{n}{2k^2} \left[\frac{dk_1 dk_2}{(2)^3} (k + k_1 + k_2) + k + c(k_2)k_2 \right]^{\frac{1}{2}} \frac{S(k)S(k_1)S(k_2)f(k_1)f(k_2)}{1 + (1 + f(k_1))(1 + f(k_2))} \right]$$
(23)

O ne should note that the right-hand-side of an analogous

self-consistent equation derived from MCT has a similar




FIG. 1: Non-ergodicity parameter. Lines: theoretical predictions at the ergodicity breaking transition; solid line: our theory, $_{g}$ = :549; dashed line: MCT, $_{g}^{MCT}$ = :525. Symbols: experimental data taken at = :563 [14].

form; the di erence is that within MCT the right-handside is a quadratic functional of f(k) [5] whereas in the present approach it includes term s of all orders in f(k).

For low enough densities Eq. (23) has only trivial solutions (i.e. f(k) = 0). For the hard-sphere interaction a non-trivial solution appears at $n_g = 3=6 = g = 549$. Qualitatively, the ergodicity breaking transition is similar to that predicted by MCT: f(k) has a jump at the transition. A lso, f(k) at the transition is similar to that of MCT at the MCT transition, $\prod_{m=1}^{MCT} m = 525$ (Fig. 1).

The factorization approximation proposed here is the simplest possible one. There are two ways to improve upon it. First, one could try to include in an approximate way the o -diagonal blocks of M. To this end one could express them in term sofa triple-density correlation function and then factorize this function into a product of three density correlation functions. Second, since the frequency matrix involves only static correlations, one could try to include it in a more sophisticated way. For exam - ple, one could include two-particle dynam ics exactly [18]. The second extension could describe glassy dynam ics in system s without static correlations [16].

To sum marize, we proposed a new theory for dynamics of concentrated suspensions and the colloidal glass transition. The theory goes beyond MCT in that includes, in an approximate way, time-dependent pair-density uctuations. In contrast to an earlier approach [17], the present one uses pair-density correlation function evolving with one-particle irreducible dynamics. The new theory predicts an ergodicity breaking transition similar to that of MCT, but at a higher density.

The author bene ted from inspiring discussions with RolfSchilling and A run Yethira j support by NSF G rant No.CHE-0111152 is gratefully acknow ledged.

- [2] See, e.g., W K.Kegeland A.van Blaaderen, Science 287, 290 (2000); E R.W eeks and D A.W eitz, Phys.Rev.Lett. 89, 095704 (2002).
- [3] See, e.g., W . Hartl, Curr. Opin. Colloid Interface Sci. 6, 479 (2001), K A . Dawson, ibid 7, 218 (2002).
- [4] M E.Cates, cond-m at/0211066.
- [5] W .G otze, in Liquids, Freezing and G lass Transition, J.P. Hansen, D. Levesque, and J. Zinn-Justin, eds. (North-Holland, Am sterdam, 1991).
- [6] G.Szam eland H.Lowen, Phys. Rev. A 44, 8215 (1991).
- [7] This applies to system s in which hydrodynam ic interactions can be neglected: strongly charged suspensions or simulated m any-particle system s with B row nian dynam – ics. For real hard-sphere-like suspensions neglecting hydrodynam ic interactions (or including them via a rescaling procedure (M . M edina-N oyola, Phys. Rev. Lett. 60, 2705 (1988))) constitutes an additional approximation; the importance of this approximation is largely unknown.
- [8] There are several ways to derive M C T. Here we discuss the original projection operator technique [5] since it is the one used to derive the new theory.
- [9] Note that for Brownian systems one has to use inreducible memory function (S J. Pitts and H C. Andersen, J. Chem. Phys. 113, 3945 (2000); see also Ref. [20]).
- [10] For Newtonian system s expressing the mem ory function in term s of pair-density correlation function neglects couplings to current modes. W ithin extended MCT currents restore ergodicity and cut o the ideal glass transition (W .G otze and L.Sjogren, Z.Phys.B 65, 415 (1987)).
- [11] G.Nagele and J.Bergenholtz, J.Chem. Phys. 108, 9893 (1998).
- [12] M. Fuchs and M E. Cates, Phys. Rev. Lett. 89, 248304 (2002).
- [13] K A. Daw son, G. Fo , M. Fuchs, et al., Phys. Rev. E 63, 011401 (2001).
- [14] W .van M egen, S M .Underwood, and P N .Pusey, Phys. Rev.Lett. 67, 1586 (1991).
- [15] M. Nauroth and W. Kob, Phys. Rev. E 55, 657 (1997); this work discusses a Newtonian system; within MCT the location of the transition does not depend on the microscopic dynamics [6].
- [16] R. Schilling and G. Szam el, Europhys. Lett.61, 207 (2003).
- [17] C Z .-W . Liu and I. Oppenheim, Physica A 247, 183 (1997).
- [18] For a sim ilar idea in a di erent context see K.M iyazaki and A.Yethiraj J.Chem. Phys. 117, 10448 (2002).
- [19] Following prior works on the colloidal glass transition, hydrodynamic interactions are neglected.
- [20] B.Cichockiand W .Hess, Physica A 141, 475 (1987).
- [21] Eq. (9) is exact for system s with pairw ise-additive interactions.
- [22] G. Szam el, to be published.