

Comment on Electron Spectral Function and Algebraic Spin Liquid for the Normal State of Underdoped High T_c Superconductors
 In a recent Letter [1], W. Rantner and X.-G. Wen made a theoretical prediction of the power-law behavior of the electron spectral function in the pseudogap phase of underdoped cuprates, reminiscent of that in the one-dimensional Luttinger liquid. This conclusion was drawn on the basis of the following (somewhat heuristic) propositions: 1) the properties of the pseudogap phase are described by the slave-boson QED₃-like effective theory formulated in terms of neutral spinons, charged holons, and a gauge field A ; 2) the propagator of physical electrons G_e can be computed as a simple product of the spinon (G_s) and the holon (G_h) ones; 3) provided that the holons are (nearly) condensed ($G_h \sim \text{const}$), G_e becomes proportional to the gauge-invariant spinon amplitude (here $x = (t; r)$ is a position vector in the three-dimensional space-time)

$$G_s(x) = \langle \psi(x) \exp(i \oint A(z) dz) \psi(0) \rangle \quad (1)$$

with the contour chosen as a straight line between the end points; 4) the amplitude (1) decays algebraically, $G_s(x) \sim |x|^{-\alpha}$, and the anomalous exponent α is positive.

It turns out, however, that the value of α quoted in Ref. [1] and later derived in the original (the only one available at the time of submitting this Comment) version of Ref. [2] had a wrong sign, as was pointed out in Ref. [3] where instead a negative value, $\alpha = -3/2N$, was obtained (the number N of fermion species in cuprates is $N = 2$).

In fact, thus far no physically motivated gauge-invariant alternative to Eq. (1) that would exhibit a power-law decay with a positive α has ever been constructed, and it remains unknown if such a function can exist at all in the pure massless QED₃. In turn, the negative value of α disqualifies Eq. (1) introduced in Ref. [1] from being a viable candidate to the role of the gauge-invariant spinon propagator (let alone the physical electron one), because instead of the anticipated suppression (as in other examples of doped Mott insulators governed by strong electron correlations), the amplitude (1) manifests enhancement as compared to the mean-field ($N = 1$) result.

The argument appears to be particularly compelling in the limit of zero doping, in which case, while preventing the electrons' spatial motion by making $G_e(x)$ vanish for any $r \neq 0$, the holon factor $G_h(x)$ does not affect the amplitude $G_e(t; 0)$ which is directly related to the physical electron density of states (DOS) proportional to $\text{Im} \int dt e^{i\epsilon t} G_e(t; 0) \sim |\epsilon|^{-\alpha}$. Therefore, a negative α would have given rise to a sub-linear DOS which is increased with respect to the mean-field "V-shaped" one.

Thus, taken at its face value, the negative α invalidates the main prediction of Ref. [1] regarding the Luttinger-like behavior of the electron spectral function $A(\epsilon; p)$

$\text{Im} G_e(\epsilon; p)$ which, under the above assumptions, was identified in Ref. [1] with the Fourier transform of Eq. (1). Moreover, once the holon factor $G_h(x)$ becomes non-trivial as well, the electron spectral function (now given by a convolution of the Fourier transforms of $G_s(x)$ and $G_h(x)$) can no longer feature a simple algebraic behavior, unless both functions decay as power-law. In the absence of any evidence suggesting otherwise, however, the possibility of such a behavior for the holons seems to be even more remote than for the spinons.

Furthermore, unless proven wrong, the absence of a physically sensible alternative to Eq. (1) in massless QED₃ may indicate a need for a revision of some of the above propositions which the work of Ref. [1] was based upon. Indeed, albeit obtained in the framework of a perturbative $1/N$ -expansion, the unphysical behavior of $A(\epsilon; p)$ derived from Eq. (1) would have manifested itself at all energies/temperatures above a characteristic scale associated with such non-perturbative effects as spinon chiral symmetry breaking, holon condensation, and/or gauge field instantons. At still lower energies/temperatures, however, any of the above mechanisms may generate a finite spinon and/or gauge field gap, thereby drastically altering the power-law decay of Eq. (1), consistent with the anticipated onset of such intrinsic instabilities of the pseudogap phase as antiferromagnetism, superconductivity and/or stripe order.

To conclude, despite its strong intellectual appeal, the QED₃-theory of underdoped cuprates has not yet provided a firm theoretical support for the Letter's prediction of the Luttinger-like or algebraic (which must be distinguished from both a generic non-Fermi liquid, characterized by a mere absence of the quasiparticle peak, and a virtually spin-charge separated Fermi liquid which has a small coherent peak at low energies) behavior of the electron spectral function, thus still leaving unsubstantiated the claim of its satisfactory agreement with the normal state photoemission data which was made in Ref. [1] on the basis of the original erroneous evaluation of Eq. (1).

This research was supported by the NSF under Grant DMR-0071362.

D. V. Khveshchenko

Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC 27599

¹ W. Rantner and X.-G. Wen, Phys. Rev. Lett. 86, 3871 (2001).

² W. Rantner and X.-G. Wen, cond-mat/0105540 (v1).

³ D. V. Khveshchenko, Phys. Rev. B 65, 53111 (2002); Nucl. Phys. B 642, 515 (2002); V. P. Gusynin, D. V. Khveshchenko, and M. Reenders, Phys. Rev. 67, 115201 (2003).