

## Collective transport and optical absorption near the stripe criticality

C. Di Castro<sup>a</sup>, S. Caprara<sup>a</sup>, and M. Grilli<sup>a</sup>

<sup>a</sup>Dipartimento di Fisica, Università di Roma "La Sapienza", and INFN Center for Statistical Mechanics and Complexity, Piazzale Aldo Moro 2, I-00185 Roma, Italy

Within the stripe quantum critical point scenario for high  $T_c$  superconductors, we point out the possible direct contribution of charge collective fluctuations to the optical absorption and to the d.c. resistivity.

There are by now several experimental evidences that the peculiar properties of the cuprates are controlled by a Quantum Critical Point (QCP), located near (actually slightly above) the optimal doping  $x = x_{opt}$  [1]. Already several years ago some of us proposed [2] that the pseudogap region of the underdoped cuprates is characterized by the pronounced tendency to form (local) spatial ordering (the so-called "stripe") phase below some charge ordering (CO) temperature  $T_{CO}(x)$ . This last temperature can therefore be related to the temperature  $T^*(x)$  at which the pseudogap forms. As a schematic model, we adopted the Hubbard-Holstein model in the presence of long-range Coulombic forces. With reasonable parameters for the cuprates, for infinite Hubbard repulsion and for moderate electron-phonon coupling there occurs a mean-field zero-temperature charge-density wave instability with a finite modulating wavevector  $q_c$  and below a rather large critical doping ( $x_c^{(0)} \approx 0.2 - 0.3$ ) (see Ref. [3] and references therein). This mean-field CO instability extends at finite temperatures below a doping dependent temperature  $T_{CO}^{(0)}(x)$ , which rapidly increases towards electronic energy scales upon underdoping. We then considered how  $x_c^{(0)}$  and  $T_{CO}^{(0)}(x)$  are modified by dynamical charge fluctuations. The critical doping  $x_c$  is reduced towards optimal doping and  $T_{CO}^{(0)}(x)$  is reduced to lower values  $T_{CO}(x)$  (typically a few hundreds of Kelvin and vanishing at the CO-QCP at  $x_c$ ). Owing to the dynamical character of the charge fluctuations,  $T_{CO}(x)$  (and the related

$T^*(x)$ ) were shown to display a marked dependence on the typical time-resolution of the experimental probes (taken as an infrared cutoff of the theory) and peculiar isotopic dependencies [3].

Recently [4] we considered the possibility that nearly critical charge fluctuations could provide an independent absorption channel in parallel with the usual Drude quasiparticle (QP) absorption. We explicitly calculated the current-current response function related to the excitation of two collective modes as represented in Fig. 1(a) in the form of an Aslamazov-Larkin-type of diagram. An additional peak appears at the finite frequency needed to excite two charge collective modes at  $q = q_c$ . This allows for a simple interpretation of low energy absorption peaks observed in several overdoped compounds [5]. These broad absorption peaks occur at low energy (below  $300 \text{ cm}^{-1}$ ), and soften upon decreasing  $T$ , thereby suggesting a collective character of the absorbing excitations. Our theoretical analysis showed that the critical character of the charge excitations entails a scaling form for the finite-frequency absorption. This expectation is well verified experimentally, see Fig. 3 of Ref. [4], and is accompanied by the temperature behavior of the absorption peak frequency (see inset of Fig. 3 in Ref. [4]), which is proportional to the expected "mass"  $m(T; x) = T + (x - x_c) + O(T^2)$  for a critical mode at the crossover from a quantum critical ( $m/T$ ) to a quantum disordered region ( $m/(x - x_c)$ ). This point of view corresponds to the idea that optimally and overdoped cuprates, when approaching the pseudogap region, are a "melting pot" of fermionic QP and collective critical excitations, both contributing to the optical

We acknowledge financial support from MIUR COFIN 2001, prot. 2001023848, and INFN/GPA-G0-4.




Figure 1. (a) CM contribution to the current-current response function. (b) CM resistivity at  $\mu = 0.22$  (dotted),  $\mu = \mu_c = 0.19$  (dashed) and  $\mu = 0.16$  (solid). The parameters of the model calculations are adjusted to yield typical values for the cuprates.

properties. We here adopt this point of view considering the charge CM contribution to transport. According to the diagram of Fig. 1(a), the result for the resistivity due to collective transport is

$$\rho_{CM}(T) / T^2(T; \mu) [a + b(T; \mu)] \quad (1)$$

Here  $a = 1:1$  and  $b = 0:4$  are numerical coefficient, while  $\mu(T; \mu) = m(T; \mu) = T$ . In Fig. 1(b) we plot three typical curves referring to the overdoped, critical, and underdoped regimes. The most apparent feature is that  $\rho_{CM}(T)$  is linear in the quantum critical region down to  $T = 0$  for  $\mu = 0:19$  and stops at finite temperature in the other two cases. This is suggestive of the linear behavior of the resistivity in nearly optimally doped cuprates. On the other hand, on the overdoped region, when the temperature decreases below the finite mass ( $m_c > 0$ ), the QP's in the fermionic loops of Fig. 1(a) no longer can emit and reabsorb these "massive" CM's and this transport channel is suppressed. At this stage  $\rho_{CM}(T)$  diverges and the transport from QP's will eventually take over. On the opposite doping side,  $\rho_{CM}(T)$  seems to vanish at a finite temperature of  $T_{CO}(\mu)$  (near  $T_c(\mu)$ ), where the mass of the CM's also vanishes. Of course, our perturbative theory can not be pushed down to this regime, where other contributions should be taken into account.

We point out that several issues have to be addressed before definite conclusions can be drawn. First of all, here we fully neglected the parallel transport channel arising from QP's. This conventional transport channel would lead to a  $T^2$  resistivity behavior, and whenever the CM resistance is not much lower than the QP one, a curvature in  $\rho(T)$  will surely arise in the total resistivity. On general grounds we expect that moving towards the overdoped region, the QP transport will eventually dominate over the collective one. Secondly, in this preliminary analysis of  $\rho_{CM}$ , we only considered the leading (i.e., critical) temperature and doping dependencies. For instance we neglected the temperature and doping dependence of the fermionic loops of the diagram in Fig. 1(a) as well as the doping dependence of the current vertices, which should be present in a strongly correlated system. While this is acceptable to identify some overall tendencies (like  $\rho_{CM} / T$ ), these dependencies must be considered to (hopefully) account for non-universal aspects like the doping dependence of the slope of the linear resistivities and the downturn in  $\rho(T)$  in underdoped cuprates. All these issues are presently under investigation.

## REFERENCES

1. See, e.g., C. Castellani, et al, Z. für Physik 103, (1997) 137; J. L. Tallon, and J. W. Loram, Physica C 349, (2001) 53.
2. C. Castellani, C. Di Castro, and M. Grilli, Phys. Rev. Lett. 75, (1995) 4650.
3. S. Andergassen, et al, Phys. Rev. Lett. 87, (2001) 056401.
4. S. Caprara, et al, Phys. Rev. Lett. 88, (2002) 147001.
5. S. Lupi, et al, Phys. Rev. B 62, (2000) 12418; E. J. Singley, et al, *ibid.*, 64, (2001) 224503; T. Startseva, et al, Physica C 321, (1999) 135. A. Lucarelli et al, Phys. Rev. Lett. 90, (2003) 037002.