

## Probing many-body states of ultra-cold atoms via noise correlations

Ehud Altman, Eugene Demler, and Mikhail D. Lukin  
 Physics Department, Harvard University, Cambridge, MA 02138  
 (dated: June 9, 2003)

We propose to utilize density-density correlations in the image of an expanding gas cloud to probe complex many-body states of trapped ultra-cold atoms. In particular we show how this technique can be used to detect superfluidity of fermionic gases and reveal broken spin symmetries in Mott-states of atoms in optical lattices. The feasibility of the method is investigated by analysis of the relevant signal to noise ratio including experimental imperfections.

Much of the excitement in the field of Bose-Einstein condensation owes to the clear demonstration it provides of the wave character of matter. The condensed state of bosons involves macroscopic occupation of a delocalized single particle state. Consequently, it is characterized by sharp density peaks in the freely expanding gas cloud after it is released from the trap [1]. Patterns that appear when two or more superfluid clouds interfere [2], are a direct probe of the single particle coherence, amplified by macroscopic occupation.

Recent experiments open intriguing directions for studying many-body phenomena beyond single particle coherence. For example, observation of the superfluid to Mott-insulator transition [3], as well as experiments involving ultra-cold fermions near a Feshbach resonance [4], address strongly correlated states of matter. The most intriguing aspect of such systems is the existence of non-trivial correlations and complex order that defy a description in terms of (single particle) matter waves. Accordingly, they cannot be characterized simply by the density profile of an expanding cloud. For example, the localized atoms in Mott states of the optical lattice display a vanishing interference pattern [3], which can hardly reveal detailed properties of the quantum state. Likewise, superfluidity of paired fermions is not evident as a coherence peak in the density profile [5], and detecting the order parameter, presents a considerable challenge. Observation of some theoretically proposed "exotic" many-atom states [6, 7, 8] may prove even more elusive.

In this Letter, we show that the quantum nature of strongly correlated states can be revealed by spatial noise correlations in the image of the expanding gas. This is similar in spirit to measurements of non-classical correlations of light in optical systems [10] and temporal current noise in mesoscopic conductors [9]. In analogy to quantum optics, this technique allows to study matter waves that lack single particle coherence. Specifically we show that (i) fermionic atoms released from the trap would display a clear signature of superfluidity in their density correlations. Furthermore, detailed properties of the fermionic superfluidity can be determined, such as pairing symmetry, and BCS to BEC crossover [11]; (ii) Atoms released from a Mott-insulating state of the optical lattice display sharp (Bragg) peaks in the density-




FIG. 1: Time of flight imaging. The atoms are released from the confining potential at time  $t = 0$  and the density of the expanding cloud is imaged at a later time. Spatial noise correlations in this image can be used to probe the quantum state at  $t = 0$ .

density correlation function as a consequence of quantum statistics; (iii) These peaks can be used to probe the spin ordered Mott states proposed for two component bosons [7, 12]. Finally we verify the experimental feasibility of the proposed measurements.

Before proceeding we note that earlier proposals to detect fermionic superfluidity relied on dynamical response of the cloud [13], inelastic scattering of light to induce and measure excitations [14, 16], or to microscopically image the pair wave-function in the trap [15]. In contrast, the present technique provides a direct probe of the pair coherence as well as information on the pairing symmetry. Earlier work on probing Mott-insulating phases should also be noted [17].

We proceed by formulating a detection scheme for atoms released from a single macroscopic trap. Suppose, for simplicity, that the system is initially in some pure state  $|j\rangle$ . In a typical experimental setup (Fig. 1), the trapping potential is turned off suddenly, and the atoms evolve independently under the influence of the free propagator  $U_0(t)$ . This is valid provided that the free-atom collision cross-section is not too large. Such conditions can be achieved by switching the magnetic field to values far from the Feshbach resonance when turning off the trap [4].

In such time of flight experiments, the column integrated density of the expanding cloud is measured by light absorption imaging [18]. The images are commonly

analyzed by comparing to theoretical predictions for the density expectation value:

$$\hat{n}(\mathbf{r})i_t = \langle \hat{\psi}^\dagger(\mathbf{r}) \hat{\psi}(\mathbf{r}) \rangle_{i_t} \quad (1)$$

where  $\hat{\psi}$  is the field operator for bosons or fermions and  $i_t$  denotes an internal atomic quantum number (spin). After a long time of flight the density distribution becomes proportional to the momentum distribution in the initial trapped state  $\hat{n}(\mathbf{r})i_t = (\hbar t)^3 \hat{n}_{\mathbf{Q}}(\mathbf{r})i_t$ . The wave-vector  $\mathbf{Q}(\mathbf{r}) = m\mathbf{r}/(\hbar t)$  defines a correspondence between position in the cloud and momentum in the trap.

It is important to realize, that in each experimental image, a single realization of the density is observed, not the expectation value. Eq. (1) is still meaningful, because the density is a self-averaging quantity. Each bin in the image represents a substantial number of atoms  $N$ , while the atomic noise scales as  $O(1/\sqrt{N})$ . But since  $N$  is not macroscopic the density fluctuations are visible. They are characterized by the correlation function:

$$G(\mathbf{r};\mathbf{r}^0) = \langle \hat{n}(\mathbf{r})\hat{n}(\mathbf{r}^0)i_t \rangle = \langle \hat{n}(\mathbf{r})i_t \hat{n}(\mathbf{r}^0)i_t \rangle \quad (2)$$

In analogy with Eq. (1) this can be related to ground state momentum correlations:

$$G(\mathbf{r};\mathbf{r}^0) / \langle \hat{n}_{\mathbf{Q}}(\mathbf{r}) \rangle \langle \hat{n}_{\mathbf{Q}}(\mathbf{r}^0) \rangle = \langle \hat{n}_{\mathbf{Q}}(\mathbf{r}) \hat{n}_{\mathbf{Q}}(\mathbf{r}^0) \rangle \quad (3)$$

The proportionality constant is  $(\hbar t)^6$  and we shall omit it henceforth. We shall be concerned primarily with pure density-density correlations  $G(\mathbf{r};\mathbf{r}^0) = \langle \hat{n}(\mathbf{r})\hat{n}(\mathbf{r}^0) \rangle$ , which does not require state-selective measurement.

In practice it may be more convenient to consider the quantity  $n(\mathbf{r};\mathbf{r}^0) = \langle \hat{n}(\mathbf{r})\hat{n}(\mathbf{r}^0) \rangle$  whose fluctuations are closely related to  $G(\mathbf{r};\mathbf{r}^0)$ . If  $\hat{n}(\mathbf{r})i_t = \hat{n}(\mathbf{r}^0)i_t$ , then

$$\langle \hat{n}(\mathbf{r};\mathbf{r}^0)^2 i_t \rangle = G(\mathbf{r};\mathbf{r}) + G(\mathbf{r}^0;\mathbf{r}^0) - 2G(\mathbf{r};\mathbf{r}^0) \quad (4)$$

Fermionic superfluids. As a specific example, we now consider superfluid states of fermionic atoms. Such superfluids sustain macroscopic coherence and their transport properties are similar to their bosonic counterparts. However, the first order coherence measured by the density profile of the expanding cloud would not reveal their superfluid nature. Consider a system at zero temperature described by a BCS-like ground state

$$|j_{\text{BCS}}\rangle = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} a_{\mathbf{k}}^\dagger a_{-\mathbf{k}}^\dagger) |j_0\rangle \quad (5)$$

Note that this wave function can describe both weak-coupling BCS-like pairing, as well as tightly bound pairs for which  $u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  have a wide momentum distribution [11]. The average density profile of the expanding cloud (1) is proportional to the BCS momentum distribution function  $\hat{n}(\mathbf{r})i_t = 2j_{\mathbf{Q}}(\mathbf{r})j_{-\mathbf{Q}}(\mathbf{r})$ , which is qualitatively indistinguishable from a Fermi distribution at  $T = T_c$  [4].

The essential difference between these states lies in the two particle correlations. For every atom with momentum  $\mathbf{k}$  in the BCS state, there is another one at exactly  $-\mathbf{k}$ . This implies, pronounced correlations between density fluctuations on diametrically opposite points in the expanding cloud. Indeed, a straightforward application of (3) for the BCS state gives:

$$G(\mathbf{r};\mathbf{r}^0) = 2j_{\mathbf{Q}}(\mathbf{r})j_{-\mathbf{Q}}(\mathbf{r}^0) \tilde{\sim}(\mathbf{r}+\mathbf{r}^0); \quad (6)$$

where  $\tilde{\sim}(\mathbf{r}+\mathbf{r}^0)$  is a sharply peaked function of  $\mathbf{r}+\mathbf{r}^0$ . This sharp peak is a direct analogue of the first order coherence peak seen in bosonic condensates. It indicates condensation of zero momentum pairs. As for BEC, it has a width  $\hbar t = mL$ , limited by the finite size  $L$  of the initial cloud. The pair distribution  $n_s(\mathbf{r}) = 2j_{\mathbf{Q}}(\mathbf{r})j_{-\mathbf{Q}}(\mathbf{r})$  can be read off directly from the weight of the peak.




FIG. 2: Fermionic superfluid. (a) Angle integrated weight of the correlation peak at diametrically opposite points. (b) Weight of the correlation peak for  $d_{x^2-y^2}$  pairing. (c) and (d)  $\langle \Delta n(r, r^0)^2 \rangle$  at  $T > 0$  (superfluid fraction 0.7). The width of the narrow dip  $\hbar t = mL$  is limited by the system's size. The dashed line in (d): BEC of tightly bound pairs.

Fig 2(a) depicts the angular averaged  $n_s(r)$  (dashed). The solid line as well and other plots in Fig 2 depict observable column integrated functions. Pairing symmetries other than s-wave, would be detected by higher angular harmonics in the pair distribution  $n_s(r)$  and its column integral. As an example we plot this function for  $d_{x^2-y^2}$  pairing in Fig. 2(b).

In practice it may be more convenient to measure the fluctuations of  $n(\mathbf{r};\mathbf{r}^0)$ . Noting that the state (5) is an eigenstate of  $n(\mathbf{r};-\mathbf{r})$  with eigenvalue zero, we have  $\langle \hat{n}(\mathbf{r};-\mathbf{r})^2 \rangle = 0$  at  $T = 0$ . Well away from  $\mathbf{r}+\mathbf{r}^0 = 0$   $\langle \hat{n}(\mathbf{r};\mathbf{r}^0)^2 \rangle = n_{\mathbf{Q}}(\mathbf{r})(1-n_{\mathbf{Q}}(\mathbf{r})) + n_{\mathbf{Q}}(\mathbf{r}^0)(1-n_{\mathbf{Q}}(\mathbf{r}^0))$ , which is non-zero on the smeared Fermi surface of a fermi liquid (at  $T > 0$ ) and for a BCS state. A sharp dip, whose width  $\hbar t = mL$  is limited by the system size, appears around  $\mathbf{r}^0 = -\mathbf{r}$  in the superfluid state (Fig 2(c,d)). Note that at finite  $T$  the magnitude of the dip is reduced and

it disappears at  $T_c$ . Heating a tightly bound pair condensate will produce thermal pairs at momenta  $k > 0$ , producing a wide background dip in analogy to the bimodal distribution in BEC.

**Optical lattice.** We now consider atoms initially confined to an optical lattice. Calculation of  $\langle \hat{n}(r) \rangle$  and  $\langle \hat{n}(r) \hat{n}(r^0) \rangle$  involves operating on the lattice state  $|j\rangle$  with  $U(t)^\dagger (r) U(t)$ . In normal ordered expectation values this operator can be safely replaced by its projection into the lowest Bloch band,  $A(r;t) = \int_{\mathbf{k}} w_{\mathbf{k}}(r;t) a_{\mathbf{k}}^\dagger$  [19]. Here  $w_{\mathbf{k}}(r;t)$  is the free evolution of a Wannier wave function, initially centered around the lattice site at  $R_{\mathbf{k}}$  and  $a_{\mathbf{k}} = \int d\mathbf{r} w_{\mathbf{k}}(r;0) \hat{\psi}(\mathbf{r})$  is a particle annihilation operator at this site. At long times  $w_{\mathbf{k}}(r;t) \sim e^{iQ(r)R_{\mathbf{k}}}$  times a gaussian envelope, which we shall approximate by a square of the same width  $W = \hbar t / (a_0 m)$ .  $a_0$  is the width of the Wannier state on the lattice. Now  $\langle \hat{n}(r) \rangle_t$  can be related to the off-diagonal correlation function in the ground state [19]:

$$\langle \hat{n}(r) \rangle_t = \langle \hat{n}_Q(r) \rangle = \frac{1}{W^d} \sum_{i,j} e^{i(R_i - R_j) \cdot Q} \langle \hat{a}_i^\dagger \hat{a}_j \rangle; \quad (7)$$

The wave-vector  $Q(r)$  now defines a correspondence between position in the cloud and the quasimomentum on the lattice.  $G$  can also be written in terms of correlations in the lattice state:

$$G(r; r^0) = \frac{1}{W^d} \sum_{i,j} e^{iR_{i1} \cdot Q(r) + iR_{j0} \cdot Q(r^0)} \langle \hat{a}_i^\dagger \hat{a}_j \hat{a}_{j^0}^\dagger \hat{a}_{i^0} \rangle + (r - r^0) \langle \hat{n}(r) \rangle_t \langle \hat{n}(r^0) \rangle_t \quad (8)$$

Here  $(r - r^0)$  is a true  $\delta$ -function originating from normal ordering in the continuum. Note that formulas (7) and (8) hold for bosons and fermions.

In the superfluid state of bosons, where  $\langle \hat{a}_i^\dagger \hat{a}_j \rangle = \langle j | j^2 \rangle$ ,  $n(r;t)$  exhibits Bragg peaks at  $Q(r)$  corresponding to reciprocal lattice vectors  $G$ . In the Mott state on the other hand  $\langle \hat{a}_i^\dagger \hat{a}_j \rangle = \delta_{ij}$ , and there is no interference pattern in  $n(r;t)$ .

The Mott state, however, displays non-trivial correlations in the second-order correlation function (8) associated with atom number fluctuations. To illustrate their origin, and to make contact with the Hanbury-Brown-Twiss effect in quantum optics, we first apply (7,8) to the "toy" model of two atoms localized in two wells. For an initial Fock state  $|j\rangle = |a_1^\dagger a_2^\dagger\rangle |j\rangle$  it gives:

$$G(r; r^0) = \frac{2}{W^{2d}} \cos \frac{m}{\hbar t} (r - r^0) \quad (9)$$

where  $\pm = 1 (-1)$  for bosons (fermions). The oscillations reflect a two body interference effect which amounts to bunching in the case of bosons and antibunching for fermions, as in the textbook example [10, 20].

Addressing the optical lattice we first consider the two

extremes of a superfluid and a Mott state:

$$|j\rangle_{SF} = \prod_i e^{za_i^\dagger} |j\rangle_i; |j\rangle_{Mott} = \prod_i \frac{1}{n!} (a_i^\dagger)^n |j\rangle_i \quad (10)$$

According to Eq. (8),  $G_{SF}(r; r^0) \sim 0$ , while

$$G_{Mott}(r; r^0) \sim \frac{N}{W^d} \frac{2}{l} \frac{a_0}{1} \sim \frac{d}{G} r - r^0 + \frac{\hbar t}{m} G \quad (11)$$

where  $l$  is a lattice spacing and  $a_0$ , the width of a Wannier function. We conclude that the Mott state sustains second order coherence, seen as Bragg peaks in  $G(r; r^0)$ . Note, again, that for a finite lattice of length  $L$ , the peaks are not true  $\delta$ -functions. They have a finite height  $(N/W^d)^2$  and width  $\sim 2\hbar t / mL$ .

Fig. 3 shows how the two-site bunching signature of Eq. (9) develops into coherent Bragg peaks with increasing lattice size. A fermionic insulating state will display similar evolution, only the peaks will be replaced by dips.




FIG. 3: Normalized density density correlations,  $g_2(r; r^0) = \langle \hat{n}(r) \hat{n}(r^0) \rangle_t / \langle \hat{n}(r) \rangle_t \langle \hat{n}(r^0) \rangle_t$  in the Mott state for a chain of 2, 4 and 40 sites.

Note that the Bragg peaks do not reflect reduced number fluctuations in the Mott state, but rather the quantum statistics of bosonic or fermionic atoms. In particular, we note that a thermal Bose gas on a lattice would exhibit similar coherent Bragg peaks, which can be seen as a result of the permutation symmetry imposed on the wave-function. However in a Mott state with non-trivial internal structure, the noise correlations can reveal complex order as we now illustrate.

As a specific example let us consider spin ordered Mott states, that have been predicted in the Mott phase of two component bosons on an optical lattice [7]. The two components can be represented by a pseudospin- $\frac{1}{2}$ :  $a_{i\#}^\dagger |j\rangle_i = |j^\# \rangle_i$ . A fairly general mean-field state of a spin-ordered insulator is given by the wave-function

$$|j\rangle_q = \prod_i (\cos \frac{1}{2} a_{i\#}^\dagger + e^{iq \cdot R_i} \sin \frac{1}{2} a_{i\#}^\dagger) |j\rangle_i \quad (12)$$

This state has a non zero spin component along the  $\hat{z}$  axis and helical component with momentum  $q$  on the  $xy$  plane. Note that the limiting case  $q = 0$  corresponds to a  $xy$  ferromagnet while  $q = \pi$  is a Néel state. An application of (8) to the state (12), yields after summation over spin indices:

$$G(r; r^0) / \left( \cos^4 \frac{q}{2} + \sin^4 \frac{q}{2} \right) \sim r - r^0 + \frac{\hbar t}{m} G + \sin^2 \frac{q}{2} \left( r - r^0 + \frac{\hbar t}{m} (q + G) \right) : \quad (13)$$

We find that the  $q \neq 0$  spin order produces satellite peaks, whose separation from the central peaks at reciprocal lattice vectors is proportional to  $q$ . The intensity ratio of the satellite to the central peaks depends on the  $\hat{z}$  polarization of the spins. The satellites reach their maximal relative intensity, half that of the central peaks, for spins polarized on the  $xy$  plane. Note that in this scheme, an  $xy$  ferromagnet cannot be observed directly. However a time dependent helical twist can be induced to this state by a magnetic field gradient.

**Experimental Issues.** To observe the proposed effects, two criteria must be met: (i) the atomic noise should be observable in an experiment and (ii) correlated fluctuations should rise above the uncorrelated noise in a statistical sense.

In a typical experiment, (6) would be averaged over narrow cylinders around  $r$  and  $r^0$ , whose bases correspond to the spatial resolution. The noise of detected probe photons in a bin of area  $A$  is a sum of contributions from atomic and laser light fluctuations. The latter is fundamentally limited by photon shot noise. The atomic noise exceeds the photon shot noise, provided the number of photons per bin in the incoming pulse,  $p > \exp(2) \hbar \Omega / \omega = (\lambda^2 / 4\pi A)$ , where  $\lambda$  is the optical depth of the cloud, which should be chosen close to unity, and  $\eta$  is the photon detection efficiency. The practical limitation on the number of photons is associated with photon recoil which results in image blurring. Under standard conditions 10–100 photons per atom can be scattered in a measurement time of 10  $\mu$ s without blurring the image on the scale of 10  $\mu$ m [18], indicating that detection of atomic noise is possible. A typical cloud of  $N = 10^7$   $^6\text{Li}$  atoms [21] will reach optical depth  $\lambda \approx 1$  after expansion to about 1 mm. Therefore each 10  $\mu$ m bin will contain about  $10^3$  atoms. The atomic noise contribution will cause bin-to-bin variation in the optical absorption at the level of few percent, which is detectable with current technology.

We now verify that in the case of a fermionic superfluid the correlated fluctuations rise above the background. For fixed  $r$  the value of  $n(r; r^0)^2$  will exhibit fluctuations of order  $n_s(r)$  as  $r^0$  varied. To observe the superfluid dip at  $r + r^0 = 0$  we must average the result over

a number of independent positions  $r$ . Different positions yield independent results if they correspond to distinct momenta in the trap. Momentum quantization thus limits the number of independent positions on the 2d image of the cloud to  $O(N^{2/3})$ , where  $N$  is the total number of atoms. Consequently the background statistical fluctuations can be averaged to  $N^{-1/3}$  of the dip magnitude making it observable in a single image measurement.

In conclusion, we demonstrated that spatial noise correlations in the image of an expanding gas cloud released from a trap, can reveal key properties of strongly correlated states of cold atoms. We anticipate that a similar technique can be used to observe signatures of more exotic states such as spin liquids and valence bond solids.

We acknowledge valuable discussions with I. Bloch, M. Prentiss and W. Ketterle. This work was supported by ARO, NSF (PHY-0134776, DMR-0132874), Sloan Foundation and Packard Foundation.

- 
- [1] M. H. Anderson et al, Science 269, 198 (1995).
  - [2] M. R. Andrews et al, Science, 275 637 (1997).
  - [3] M. Greiner, O. Mandel, T. Esslinger, T. W. Hansch and I. Bloch, Nature, 415, 39 (2002).
  - [4] A. G. Truscott et al, Science 291, 2570 (2001); D. Loffus et al, Phys. Rev. Lett. 88, 173201 (2002); K. M. O'Hara et al, Science 298, 2179 (2002); T. Bourdel et al, cond-mat/0303079; S. Gupta et al, Science, in press, (2003).
  - [5] M. Houbiers et al, Phys. Rev. A 56, 4864 (1997); M. Holland et al, Phys. Rev. Lett. 87, 120406 (2001); E. Timmermans Phys Lett A 285, 228 (2001).
  - [6] E. Demler, F. Zhou, Phys. Rev. Lett. 88, 163001 (2002).
  - [7] L.-M. Duan, E. Demler and M. D. Lukin cond-mat/0210564, E. Altman et al, in preparation.
  - [8] A. Recati et al, J. of Opt. B 5, S55 (2003); B. Peredes and J. I. Cirac, Phys. Rev. Lett. 90, 150402 (2003).
  - [9] Ya. M. Blanter and M. Buttiker, Phys. Rep. 336, 1 (2000).
  - [10] see e.g. M. O. Scully and M. S. Zubairy, "Quantum Optics", Cambridge University Press (1997).
  - [11] M. Randeria in "Bose-Einstein Condensation", ed. A. Griffin, D. W. Snoke and S. Stringari, Cambridge University Press (1995).
  - [12] A. B. Kulikov and B. V. Svistunov, cond-mat/0205069.
  - [13] C. Menotti, P. Pedri, S. Stringari, Phys. Rev. Lett. 89, 250402 (2002)
  - [14] W. Hofstadter et al, Phys. Rev. Lett. 89, 220407 (2002)
  - [15] W. Zhang, C. A. Sackett and R. G. Hulet, Phys. Rev. A, 60, 504 (1999).
  - [16] P. Toma and P. Zoller, Phys. Rev. Lett. 85, 487 (2000).
  - [17] D. C. Roberts and K. Burnett, Phys. Rev. Lett. 90, 150401 (2003).
  - [18] W. Ketterle, D. S. Durfee, D. M. Stamper-Kum, cond-mat/9904034.
  - [19] R. Roth and K. Burnett Phys. Rev. A 67, 031602(R) (2003).
  - [20] W. Oliver et al, Science, 284, 299 (1999).
  - [21] Z. Hadzibabic et al, cond-mat/0306050.