

Cyclotron Resonance Study of the Two-Dimensional Electron Layers and Double-Layers in Tilted Magnetic Fields

Nataliya A. Goncharuk, Ludvík Smrčka¹, and Jan Kučera

Institute of Physics ASCR, Cukrovarnická 10, 162 53 Praha 6, Czech Republic

Abstract

The far-infrared absorption in two-dimensional electron layers subject to magnetic field of general orientation was studied theoretically. The Kubo formula is employed to derive diagonal components of the magneto-conductivity tensor of two-dimensional electron single-layers and double-layers. The parabolic quantum well is used to model a simple single-layer system. Both single-layer and double-layer systems can be realized in a pair of tunnel-coupled, strictly two-dimensional quantum wells. Obtained results are compared to experimental data.

Key words: single-layer, double-layer, two-dimensional electron system, cyclotron resonance

PACS: 73.21.Fg, 73.63.Hs

1. Introduction

The cyclotron resonance in the absorption of far-infrared radiation in two-dimensional electron single-layers and double-layers occurs when its frequency ω coincides with the frequency ω_c of the circular motion of electrons subjected to a magnetic field \mathbf{B} .

In the quasi-classical picture of the cyclotron resonance, the current (and, consequently, the absorbed power) induced in the layer by the electric field reach maximum at resonance conditions, $\omega = \omega_c$.

From the point of view of quantum-mechanics, the cyclotron resonance is described as excitation of electrons from occupied to unoccupied Landau levels by photons with the energy $\hbar\omega$ equal to the energy difference between levels, $\hbar\omega_c = \hbar|e|B/m_c^*$.

The cyclotron mass is defined by $m_c^* = |e|B/\omega_c$. In strictly two-dimensional systems only B_\perp , the field component perpendicular to the electron layers, contributes to the cyclotron motion of electrons and a single cyclotron mass characterizes the cyclotron motion for all magnitudes of the magnetic field B_\perp . Both quasi-classical and quantum-mechanical approaches should yield the same results for small B_\perp , when many Landau levels are occupied.

The situation becomes more complicated if we consider quasi-two-dimensional electron layers and double-layers in magnetic fields of a general orientation. Due to the finite width of quantum wells, the electronic structure is strongly affected by the in-plane component of the field, B_\parallel . The cyclotron mass is not longer a constant and becomes a function of both B_\parallel and B_\perp , $m_c^* = m_c^*(B_\parallel, B_\perp)$, as demonstrated by recent experiments (see e.g. [1]).

We have studied this problem theoretically. Assum-

¹ Corresponding author. E-mail: smrcka@fzu.cz

ing the Landau gauge of the vector potential, $\mathbf{A} = (-B_{\perp}y + B_{\parallel}z, 0, 0)$, and a wave function in the form $\psi \propto \exp ik_x x \varphi(y, z)$, the Hamiltonian can be written as

$$H = \frac{1}{2m^*}(\hbar k_x + eB_{\perp}y - eB_{\parallel}z)^2 + \frac{p_y^2}{2m^*} + \frac{p_z^2}{2m^*} + V_{\text{conf}}(z). \quad (1)$$

First, parabolic quantum wells are employed to model single-layer systems. Then, a pair of tunnel-coupled, strictly two-dimensional quantum wells located at the distance d , is studied. These simple models allow to carry out most calculations analytically and yet are able to capture the essence of physics of single-layer/double-layer systems.

In quantum-mechanical approach, the linear response theory (the Kubo formula) is used to evaluate diagonal components of the magnetoconductivity tensor responsible for determination of the dissipated power. Since our aim is to study mainly the field-dependence of the energy levels $E_{m,n} = E_{m,n}(B_{\parallel}, B_{\perp})$ and of the transitions between them, we completely neglect the broadening of levels by scattering of electrons.

In both single-layers and double-layers we consider only the ground and excited subbands which yield two fans of Landau levels. Each Landau level is denoted by a subband index m , and a level index n . Corresponding wave functions determine probabilities of transitions between levels via the dipole matrix elements in the Kubo formula.

2. Results and discussion

2.1. Parabolic quantum wells

The quantum well with a parabolic confining potential $V_{\text{conf}}(z) = m^* \Omega^2 z^2 / 2$ represents a particularly simple model which can be solved analytically. The frequency Ω defines the separation of two subbands at $\mathbf{B} = 0$. Eigenenergies are given as solutions of two coupled harmonic oscillators with frequencies depending strongly on B_{\parallel} and B_{\perp} . The quasi-classical results are obtained from quantum-mechanical ones simply by taking limit $B_{\perp} \rightarrow 0$.

In tilted magnetic fields Landau-level mixing occurs and, besides intrasubband transitions, $\delta m = 0$, $\delta n =$




Fig. 1. Landau eigenenergies as functions of B_{\parallel} for fixed $B_{\perp} = 0.5$ T. Results of calculation for the parabolic model of 2D electron single-layers. The energy of subband separation is $\hbar\Omega = 4$ meV.

± 1 , also intersubband transitions, $\delta n = 0$, $\delta m = \pm 1$, become bright. For fixed B_{\perp} , the separation of mixed levels decreases with increasing B_{\parallel} . As it is the same for all adjacent levels, the cyclotron mass does not depend on the level index and grows with B_{\parallel} in our simple model.




Fig. 2. The B_{\parallel} -dependence of m_c^* for different fixed B_{\perp} and Ω . The parameter Ω determines the subband separation, $\Delta E = E_{1,n} - E_{0,n} = \hbar\Omega$.

Variation of the electron concentration can be modeled by changing Ω : the higher the concentration, the narrower the well. We have found that the wider

parabolic wells are more sensitive to the magnetic field as demonstrated in Fig. 2. We have also shown (see Fig. 3) that with our model one can reasonably fit the B_{\parallel} - and B_{\perp} -dependence of experimental data [1].




Fig. 3. The B_{\parallel} dependence of m_c^* at $B_{\perp} = 6$ T. Theoretical curves (dashed lines) are shown in comparison with experimental results of S. Takaoka [1] (full lines) on samples with different electron concentrations ($2.0 \times 10^{11} \text{ cm}^{-2}$ in sample A and $3.0 \times 10^{11} \text{ cm}^{-2}$ in sample B).

2.2. Double-well structures

In double wells, the lowest bound states of individual quantum wells form symmetric and antisymmetric pairs. The ground and excited states become separated on energy scale by $2|t|$, where t is the coupling constant (see e.g. [2]). In real space, the bonding and antibonding 2D electron layers overlap and share the full width of a double-well structure.

The Landau levels with high quantum numbers can be described quasi-classically [3,4] if $B_{\perp} \ll B_{\parallel}$. The quantum-mechanical approach must be employed for the case of strong tilted field: we must solve coupled differential equations for the left and right components of the double-layer wave function. The resulting eigenenergies and eigenvectors are used to calculate the cyclotron effective mass and transition matrix elements.

The eigenenergies in the excited (antibonding) subband, shown in Fig. 4, sharply grow with increasing B_{\parallel} for given B_{\perp} . The cyclotron mass corresponding to transition between Landau levels of the antibonding




Fig. 4. Eigenenergies of the 2D electron systems as a function of B_{\parallel} at fixed $B_{\perp} = 0.5$ T and $|t| = 2$ meV. Lower and upper bunch of lines denote Landau levels from the ground (bonding) and excited (antibonding) subband, respectively.

subband decreases with increasing in-plane field and differs for each pair of levels (see Fig. 5).

The behaviour of levels from the ground subband is more complicated. The in-plane component of the magnetic field reduces the tunneling between wells and, at the high magnetic field limit, the single-layer of bonding electrons is divided into two layers, each layer being localized in one of the wells. Landau levels of a double-layer are, therefore, degenerated (each level can be de-




Fig. 5. Cyclotron effective masses m_c^* calculated from energy difference between adjacent Landau levels of the band structure spectrum (figure 4) as a function of the in-plane magnetic field.

noted by a well index). The degeneracy in the well index is removed for lower B_{\parallel} (two layers gradually merge into a single-layer) and for $B_{\parallel} = 0$ each originally degenerated level is transformed into an even/odd pair of Landau levels of the bonding subband.

The cyclotron mass corresponding to transitions between such a pair of levels exhibit a singularity at high B_{\parallel} (the transition becomes dark there).

The energy separation of neighbouring levels from different pairs first decreases with increasing B_{\parallel} , reaches minimum around a critical field, and then returns to an original value. Consequently, the corresponding cyclotron mass has a maximum at the critical field. The transitions between such Landau levels are bright in the high field limit, they corresponds to transitions between Landau levels in individual quantum wells.




Fig. 6. The B_{\parallel} -dependence of the cyclotron effective mass m_c^* calculated quasi-classically. The cyclotron effective mass is normalized by the cyclotron effective mass at zero magnetic field limit, m^* . Full and dashed lines correspond to ground and excited subbands, respectively.

The single-layer/double-layer transformation of electrons in the bonding subband is even more clearly demonstrated in the quasi-classical case. For very small B_{\perp} the number of occupied Landau levels increases and it is more convenient to describe the B_{\parallel} -dependence of the cyclotron mass in terms of the Fermi energy, which separates the empty and occupied states, rather than with a help of quantum numbers of empty and occupied Landau levels.

The results are presented in Fig. 6. The cyclotron mass of the bonding electrons exhibits a logarithmic singularity at the critical field, above which the electrons cannot tunnel through the barrier and form two insulated parallel electron sheets. The growth of the electron concentration shifts the singularity position to the side of larger in-plane fields.

3. Conclusions

We have performed a theoretical study of the cyclotron resonance in 2D single-layers and double-layers subject to magnetic fields of general orientation.

For strong B_{\parallel} and B_{\perp} , the quantum-mechanical approach applies: we have calculated the field-dependence of the energy levels and transitions matrix elements between them, and found that m_c^* becomes a function of both field components.

In strictly perpendicular magnetic fields only the intrasubband transitions ($\delta m = 0, \delta n = \pm 1$) are allowed. In tilted magnetic fields also the intersubband transitions ($\delta m = \pm 1, \delta n = 0$) become possible. In general, the intrasubband transitions become dark and the intersubband transitions bright at high B_{\parallel} .

The field dependence of the cyclotron mass in double well structures is more complex than in the single wells.

4. Acknowledgements

This work was supported by AVOZ1-010-914 and by the Grand Agency of the Czech Republic under Grant No. 202/01/0754.

References

- [1] Aikawa H, Takaoka S, Oto K, Murase K, Saku T, Hirayama Y, Shimomura S and Hiyamizu S 2002 *Physica E* **12** 578-580
- [2] Hu J and MacDonald A H 1992 *Phys. Rev. B* **46** 12554-12559
- [3] Smrčka L, Jungwirth T 1995 *J. Phys.: Condens. Matter* **7** 3721-3732
- [4] Takaoka S, Kuriyama A, Oto K, Murase K, Shimomura S, Hiyamizu S, Cukr M, Jungwirth T, Smrčka L 2000 *Physica E* **6** 623-626