

Kirkwood Phase Transition for Boson and Fermion Hard-Sphere Systems

M.A. Solís^{1,2}, M. de Llano³ and J.W. Clark¹

¹ Department of Physics, Washington University
St. Louis, Missouri 63130, USA

² Instituto de Física, UNAM, Apdo. Postal 20-364,
01000 México, D.F., Mexico

³ Instituto de Investigaciones en Materiales, UNAM,
Apdo. Postal 70-360, 04510 México, DF, Mexico

March 22, 2022

Abstract

The London ground-state energy formula as a function of number density ρ for a system of boson hard spheres of diameter c at zero temperature (corrected for the reduced mass of a pair of particles in a “sphere-of-influence” picture) generalized to describe fermion hard-sphere systems with four and two intrinsic degrees of freedom such as ^3He or neutron matter and symmetric nuclear matter, respectively, is proposed as the crystalline energy branch for hard-sphere systems. For the fluid branch we use the well-known, exact, low-density equation-of-state expansions for many-boson and many-fermion systems, appropriately extrapolated to physical densities. Here, via a double-tangent construction the crystallization and melting densities for boson and fermion hard spheres are determined. They agree well with variational Monte Carlo, density-functional, and Green Function Monte Carlo calculations.

PACS: 05.30.-d; 21.65.+f; 67.90.+z

Key words: boson and fermion hard-sphere systems; nuclear and neutron matter; freezing transition.

1 Introduction

As in the classical hard-sphere system where a “freezing” transition from a fluid to a crystalline phase was found [1] to occur as one increased density, one expects the same phenomenon to occur in quantum systems, whether of bosons or fermions.

An analytical formula for the ground state energy E of a boson hard-sphere system for all densities ρ was proposed by London [2] as

$$E/N = \frac{2\pi\hbar^2 c}{m} \frac{1}{(\rho^{-1/3} - \rho_0^{-1/3})^2} \frac{1}{(\rho^{-1/3} + b\rho_0^{-1/3})}, \quad (1)$$

where m is the particle mass, c is the hard-sphere diameter, with $\rho_0 \equiv \sqrt{2}/c^3$ the ultimate density [3] at which a system of identical classical hard spheres close packs in some primitive-hexagonal arrangement, e.g., face-centered-cubic, and $b = (2^{5/2}/\pi) - 1$. The justification given for Eq. (1) is that it reduces smoothly to well-known limiting expressions at both low and high densities, namely

$$E/N \xrightarrow{\rho \rightarrow 0} (2\pi\hbar^2/m)\rho c, \quad (2)$$

$$E/N \xrightarrow{\rho \rightarrow \rho_0} A (\hbar^2/2m)(\rho^{-1/3} - \rho_0^{-1/3})^{-2}. \quad (3)$$

Here $A = \pi^2/2^{1/3} \simeq 7.834$ is a constant called the *residue* of the pole at close packing. The low-density leading term (2) is the celebrated Lenz [4] term, calculated by him as the leading correction to the energy due to an “excluded volume” effect. The limiting case Eq. (3) comes from the lowest Schrödinger equation eigenvalue of a particle in the spherical cavity. It is just the kinetic energy of a point particle of mass m inside a spherical cavity of radius $r - c$, where r is the average separation between two neighboring hard spheres and $r = (\sqrt{2}/\rho)^{1/3}$ by assuming a primitive-hexagonal arrangement for the cavities.

It was found [5] that the derivation of the high-density extreme of the original [2] (boson) London equation (1) contains *one fundamental error*: the spherical cavity of radius $r - c$ alluded to above in reality refers to a “sphere of influence” of *two* particles. Thus, the particle mass used in obtaining (3) *should refer to the reduced mass $m/2$* . This yields the constant

$$b \equiv 2^{3/2}/\pi - 1 \quad (4)$$

instead of $2^{5/2}/\pi - 1$ given by London (1). The result (1) with (4) is designated the *modified London (ML) equation*, which continues to satisfy (2) as this is independent of the constant b . The residue A in (3) then becomes $2^{2/3}\pi^2 \simeq 15.7$ and fully agrees with the empirical residue of 15.7 ± 0.6 , extracted by Cole [6] from high-pressure crystalline-branch data in ^3He , ^4He , H_2 and D_2 systems. Moreover, this ML equation agrees dramatically better than the original London

(L) equation with Green Function Monte Carlo (GFMC) [7] computer-simulation datapoints for both fluid and crystalline branches of the boson hard-sphere system.

A generalized London equation has also been proposed [8] for N -fermion hard-sphere systems, with ν intrinsic degrees of freedom for each fermion. Two differences appear with respect to the boson London formula: a) unlike the boson case, the ground state kinetic energy is nonzero and is added as a well-known [9] ν -dependent leading term, and b) the constant b is allowed to be ν -dependent, namely $b(\nu) = [(\nu - 1)/\nu](b + 1) - 1$. This form ensure a ν -independent energy at close-packing where, since the spheres can be labelled indistinguishability and thus particle statistics are lost as expected in this classical limit. Replacing the value of the constant b by the new one (4) gives a *generalized modified London equation* (ML $_{\nu}$)

$$E/N = C_{\nu} \rho^{2/3} + \left(\frac{\nu - 1}{\nu} \right) \frac{2\pi\hbar^2 c}{m} \frac{1}{(\rho^{-1/3} - \rho_0^{-1/3})^2} \frac{1}{[\rho^{-1/3} + b(\nu) \rho_0^{-1/3}]} \quad (5)$$

with

$$C_{\nu} \equiv \frac{3\hbar^2}{10m} \left(\frac{6\pi^2}{\nu} \right)^{2/3} \xrightarrow[\nu \rightarrow \infty]{} 0 \quad (6)$$

where in the limit $\nu \rightarrow \infty$ the new constant $b(\nu) \rightarrow b$ and (5), i.e., goes over into the boson case stated above. The low-density limit of Eq. (5) becomes

$$E/N \xrightarrow[\rho \rightarrow 0]{} C_{\nu} \rho^{2/3} + \left(\frac{\nu - 1}{\nu} \right) \frac{2\pi\hbar^2}{m} \rho c, \quad (7)$$

where the second term on the rhs is the Lenz term for ν -component fermions. On the other hand, for $\rho \rightarrow \rho_0 \equiv \sqrt{2}/c^3$ one sees that (5) reduces to (3) as it should. In other words, hard-sphere fermions, bosons or “boltzons” *must close pack at the same density*. From this it follows that the residue for bosons or fermions is the same and equal to $2^{2/3} \pi^2 \simeq 15.7$, in line with Ref. [6].

In addition to the Lenz term (2) for the low-density fluid branch, several higher-order corrections to the ground state energy per particle have been derived using quantum field-theoretic many-boson perturbation theory [10], namely

$$E/N = \frac{2\pi\hbar^2 \rho c}{m} \left\{ 1 + C_1 (\rho c^3)^{1/2} + C_2 \rho c^3 \ln(\rho c^3) + C_3 \rho c^3 + o(\rho c^3) \right\}, \quad (8)$$

where $C_1 = 128/15\sqrt{\pi}$, $C_2 = 8(4\pi/3 - \sqrt{3})$, but C_3 is an as yet unknown constant. Here c is the S-wave scattering length in general, and becomes the sphere diameter for a hard-core potential. The series is clearly not a power series expansion, and is at best an asymptotic series.

Similarly, for an N -fermion hard-sphere system the corresponding series is [11]

$$\begin{aligned} E/N &= \frac{3}{5} \frac{\hbar^2 k_F^2}{2m} \left\{ 1 + C_1(k_F c) + C_2(k_F c)^2 \right. \\ &+ [C_3 r_0/2c + C_4 A_1(0)/c^3 + C_5](k_F c)^3 + C_6(k_F c)^4 \ln(k_F c) \\ &+ [C_7 r_0/2c + C_8 A_0''(0)/c^3 + C_9](k_F c)^4 + o(k_F c) \left. \right\}, \end{aligned} \quad (9)$$

where the C_j ($j = 1, 2, \dots, 9$) are dimensionless coefficients depending on ν and are given in Ref. [12] for $\nu = 2$ and $\nu = 4$. The Fermi momentum $\hbar k_F$ is defined through the particle number density $\rho \equiv N/\Omega = \nu k_F^3/6\pi^2$, with Ω the system volume.

Unfortunately, both low-density expansions (8) and (9) lack accuracy at moderate to high densities, including the saturation (or equilibrium, zero-pressure) densities of liquid ^4He ($\nu = \infty$) and liquid ^3He ($\nu = 2$) or nuclear matter ($\nu = 4$). However, one can extrapolate the series for hard-sphere systems to physical and even to close-packing densities through the use of Padé and so-called “tailing” [13] approximants. This method, called Quantum Thermodynamic (or Van der Waals) Perturbation Theory (QTPT) [14], has provided fairly accurate representations of the *fluid* branch of the equation of state [15] beyond the presumable phase transition densities but without enough credibility as one approaches close packing. This is clear since one does not possess a ground-state energy function with implicit information of *both* fluid and crystalline branches. Thus, for the *crystalline* branch we employ the generalized modified London equations (1) which reproduce the correct closest packing density value $\rho = \sqrt{2}/c^3$ as well as good behavior near closest packing as suggested by a correct residue value. For the *fluid* branch we use the Padé extrapolations based on the series (8) and (9).

This paper unfolds as follows. In Sections 2 and 3 we construct the fluid branches for hard-sphere bosons and fermions, respectively. In Sec. 4 we use them together with their respective modified London equation crystalline branches to determine, via a double-tangent construction ensuring equality of pressure in both phases, the melting and crystallization densities as well as energy and density changes. In Sec. 5 we state our conclusions.

2 Boson hard-sphere fluid branch

In order to extrapolate to higher densities the low-density series (8), we define

$$E/N = \frac{2\pi\hbar^2}{m}\rho c e_0(x), \quad x \equiv (\rho c^3)^{1/2}, \quad (10)$$

with

$$e_0(x) \equiv 1 + C_1 x + C_2 x^2 \ln x^2 + C_3 x^2 + O(x^3 \ln x^2). \quad (11)$$

Alternatively, one can analyze the series

$$e_0^{-1/2}(x) = 1 + F_1 x + F_2 x^2 \ln x^2 + F_3 x^2 + O(x^3 \ln x^2) \quad (12)$$

for $x \ll 1$, where the F 's are expressible in terms of the C 's, with C_3 and consequently F_3 unknown. Values of the C 's and F 's are given in Table 1. We analyze $e_0^{-1/2}(x)$ instead of $e_0(x)$ to ensure that zeros in its extrapolants $\epsilon_0^{-1/2}(x)$ are second-order poles in energy as we

expect at close packing (CP). Then we examine the twelve extrapolants [16] from the series (12) with three terms. Fitting extrapolants to go through the four GFMC data points [7] allows estimating a good value for the last coefficient F_3 . Forms VII and XI both satisfied these conditions but the mean square deviation with respect to the four GFMC data points was least with form XI. Thus we use it as our best extrapolant. So the ground state energy for boson hard spheres is represented (symbol \doteq) by

$$E/N \doteq \frac{2\pi\hbar^2}{m}\rho c\epsilon_0(x), \quad (13)$$

with

$$\epsilon_0^{-1/2} = \text{XI}(x) = \frac{1 + F_2 x^2 \ln x^2}{1 - F_1 x - (F_3 - F_1^2) x^2} \quad (14)$$

and $F_3 = -27.956$. In Fig. 1 this is plotted as a full curve and labeled XI for the fluid branch. Open circles and squares are GFMC data for fluid and crystalline branches, respectively. The dashed curve is the Modified London formula (1) representing the crystalline branch. Dots are Diffusion Monte Carlo (DMC) [17] calculations spanning a wider range of densities in the fluid region than the GFMC data. From DMC data we see that although our expression for the fluid branch could be better for intermediate densities, it agrees well with DMC and GFMC calculations around the freezing transition. This $\epsilon_0^{-1/2}$ predicts a random close packing (RCP) density $\rho/\rho_0 = 0.776$ which is only ten percent below classical RCP $\rho/\rho_0 \simeq 0.86$ [3] expected to be the ultimate CP density also for quantum hard sphere fluids just from the fact that particles at CP are localized so that indistinguishability does not hold and thus statistics are irrelevant.

3 Fermion hard-sphere fluid branch

For fermion hard-sphere systems we write (9) as

$$E/N = \frac{3}{5} \frac{\hbar^2 k_F^2}{2m} e_0(x); \quad x \equiv k_F c, \quad \rho \equiv N/\Omega = \nu k_F^3/6\pi^2, \quad (15)$$

Table 1. For bosons, C_i and F_i coefficients appearing in Eqs. (11) and (12), respectively. Numbers in quotation marks are adjusted as indicated in text.

Bosons ($\nu = \infty$)	$i = 1$	2	3
C_i	4.81441778	19.65391518	“73.296”
F_i	-2.40720889	-9.826957589	“-27.956”




Figure 1: The quantity $\varepsilon_0^{-1/2} = \sqrt{2\pi\hbar^2 \rho c N / m E} = [1 - (\rho/\rho_0)^{1/3}] \sqrt{1 + b (\rho/\rho_0)^{1/3}}$ as a function of ρ/ρ_0 for boson hard sphere systems: XI is the fluid branch (14), ML is the Modified London formula (1), open circles and squares are GPMC data for the fluid and crystalline branches respectively, and dots are DMC calculations.

with

$$e_0(x) = 1 + C_1 x + C_2 x^2 + (C_3/3 + C_4/3 + C_5) x^3 + C_6 x^4 \ln x + (C_7/3 - C_8/3 + C_9) x^4 + o(x^4). \quad (16)$$

For $\nu = 2$, $C_6 = 0$ [11] so that (16) becomes

$$e_0(x) \simeq 1 + D_1 x + D_2 x^2 + D_3 x^3 + D_4 x^4 + o(x^4) \quad (17)$$

for $x \ll 1$. With D 's in terms of C 's and $\rho = k_F^3/3\pi^2$. As in the boson case, instead of e_0 we write

$$e_0^{-1/2} \simeq 1 + F_1 x + F_2 x^2 + F_3 x^3 + F_4 x^4 + F_5 x^5 + o(x^5), \quad (18)$$

where the F_i 's depend algebraically on the D_i 's in a simple manner, F_5 being unknown. Values of D_i and F_i are given in Table 2. This series is a simple power series and we use it to construct the usual Padé extrapolants. The approximants to (18) with four terms beside the trivial one were analyzed in Ref. [18] concluding that the best was the Padé $[4/0](x)$. However, this function does not have a zero in the region of physical interest, i.e. $0 \leq \rho/\rho_0 \leq 1$ or $0 \leq x \leq 3.47$, and so the energy does not manifest a CP as it should. It was necessary to introduce the fifth term in (18). Its five *two-point* Padé approximants $[3//2](x)$ were analyzed and F_5 adjusted to ensure a zero. The position of the zero and the approximant were chosen in such way that the QTPT applied to ^3He with the Aziz inter-atomic potential reproduces

the corresponding GFMC [20] data. Eventually, the best extrapolant was the two-point Padé approximant [19]

$$\epsilon_0^{-1/2} \doteq [3//2](x). \quad (19)$$

Hence E/N becomes

$$E/N = \frac{3}{5} \frac{\hbar^2 k_F^2}{2m} ([3//2](x))^{-2}, \quad (20)$$

with a CP density $\rho/\rho_0 = 0.732$. The coefficient F_5 is listed Table 2 between quotation marks. In Fig. 2 $\epsilon_0^{-1/2} = (3\hbar^2(6\pi^2\rho/\nu)^{2/3}N/10mE)^{1/2} = 1 + (20\pi(\nu-1)/3\nu)(2^{1/4}\nu/6\pi^2)^{2/3}\{[(\rho/\rho_0)^{-1/3} - 1]^2[(\rho/\rho_0)^{-1/3} - b(\nu)](\rho/\rho_0)^{2/3}\}^{-1}$ as a function of ρ/ρ_0 is plotted for fermion hard sphere. For $\nu = 2$ the fluid branch $[3//2]$ (full curve) is close to the numerically exact Ladder (Ladder) [21] (open squares), Variational Fermion-Hypernetted-Chain (VFHNC) [22] (plus sign marks) and L-expansion data [23] (open triangles). Fig. 3 is a enlargement of Fig. 2 where we show this excellent agreement over the whole range of available data.




Figure 2: The quantity $\epsilon_0^{-1/2} = (3\hbar^2(6\pi^2\rho/\nu)^{2/3}N/10mE)^{1/2} = 1 + 20\pi(\nu-1)/3\nu \times (2^{1/4}\nu/6\pi^2)^{2/3}\{[(\rho/\rho_0)^{-1/3} - 1]^2[(\rho/\rho_0)^{-1/3} - b(\nu)](\rho/\rho_0)^{2/3}\}^{-1}$ as a function of ρ/ρ_0 for fermion hard sphere with $\nu = 4$ (XII) and $\nu = 2$ ($[3//2]$), are full lines. Dashed lines are the corresponding Modified London formulas.

For $\nu = 4$, Eq. (16) becomes

$$e_0(x) = 1 + D_1 x + D_2 x^2 + D_3 x^3 + D_4 x^4 \ln x + D_5 x^4 + o(x^5) \quad (21)$$

for $x \ll 1$. By analogy with bosons and fermions ($\nu = 2$), we analyze

$$e_0^{-1/2} = 1 + F_1 x + F_2 x^2 + F_3 x^3 + F_4 x^4 \ln x + F_5 x^4 + o(x^5) \quad (22)$$

Table 2. For $\nu = 2$, D_i and F_i coefficients appearing in Eqs. (17) and (18), respectively. Numbers in quotation marks are adjusted as is indicated in text.

$\nu = 2$	$i = 1$	2	3	4	5
D_i	0.353678	0.185537	0.384145	-0.024700	“-0.265544”
F_i	-0.176833	-0.045863	-0.156677	0.109672	“0.130830”

with all stated F_i known. Values of D_i and F_i are given in Table 3. Unlike the $\nu = 2$ case, this series is not a pure power series as it contains logarithmic terms. Its tailing approximants are giving in Table III of Ref. [13]. We choose form XII to avoid approximants with spurious unphysical poles within the interval of physical densities, as well as to avoid residues falling outside the *rigorous* interval

$$1.63 \leq A \leq 27. \quad (23)$$

This interval was obtained [24] for regular CP (face-centered-cubic or hexagonal-close-packing) by generalizing the exact calculation for a simple cubic lattice based on three mutually perpendicular linear lattice which gives $A = \pi^2$.




Figure 3: Enlargement of Fig. 2 at low densities.

Hence, E/N can be written

$$E/N = \frac{3\hbar^2 k_F^2}{10m} \epsilon_0(x), \quad (24)$$

with

$$\epsilon_0^{-1/2}(x) \doteq \text{XII}(x) = \frac{1 + (F_1 - F_3/F_2)x + (F_2 - F_1 F_3/F_2)x^2}{1 - (F_3/F_2)x - F_4 x^4 \ln x}. \quad (25)$$

Table 3. For $\nu = 4$, D_i and F_i coefficients appearing in Eqs. (21) and (22), respectively.

$\nu = 4$	$i = 1$	2	3	4
D_i	1.061033	0.556610	1.300620	-1.408598
F_i	-0.530517	0.143867	-0.5806558	-0.704299

This is plotted in Fig. 2 and 3 as a full curve and labeled XII. We also plot the corresponding VFHNC data (plus signs) and L-expansion data (open triangles). In terms of energy, ours are slightly below the VFHNC ones. The agreement gets better for lower densities. However, the XII approximant is just above the L-expansion data over the range of densities where they are available.

4 Kirkwood phase transition

Fluid-to-crystal (or freezing) phase transitions for boson hard-sphere systems have been determined with VMC, GFMC or Density Functional Theory (DFT) [25]. However, for fermion hard-sphere systems such calculations have not been done. The purpose of this paper is to provide very simple but hopefully accurate analytical expressions for the ground-state energy of quantum hard-sphere systems in general so as to reproduce crystallization transitions or to give the values for the parameter involved, as well as other thermodynamic properties such as pressure and compressibility.

Here, via a double-tangent construction we locate the crystallization ρ_l and melting ρ_s number densities for bosons and for fermions (with $\nu = 2$ and 4). This process is schematically sketched in Fig. 4 for fermions with $\nu = 2$, where both fluid and crystalline energies per particle, in \hbar^2/mc^2 units, as a function of ρ/ρ_0 , are shown close to the phase transition. Dots denote crystalline and fluid phase Maxwell-construction endpoints on the pressure-volume diagram, obtained by constructing a common tangent to both energy branches ensuring equality of pressure in both phases.

For boson hard-spheres, in Table 4 we list our calculation (ML) together with the VMC, GFMC and DFT ones. All of them agree reasonably well with each other. Here $\Delta(\rho c^3)$ and $\Delta(E/N)$ signify the change of density and energy, respectively, between both phases. For fermion hard spheres with $\nu = 4$, e.g., symmetric nuclear matter, we find a crystallization density given by $\rho_l c^3 = 0.056$ and a melting density given by $\rho_s c^3 = 0.069$. This crystallization

Table 4. Crystallization and melting densities for boson hard sphere system. NA means not available.

Bosons	$\rho_l c^3$	$\rho_s c^3$	$\Delta(\rho_s c^3)$	$\Delta(E/N)$
VMC [26]	0.23 ± 0.02	0.25 ± 0.02	0.02	1.28
DFT [25]	0.246	0.284	0.38	2.668
GFMC [7]	0.25 ± 0.01	0.27 ± 0.01	0.02	NA
ML	0.21 ± 0.02	0.23 ± 0.02	0.02	1.37

Table 5. Crystallization and melting densities for fermion ($\nu = 4, 2$) hard sphere system.

Fermions	$\rho_l c^3$	$\rho_s c^3$	$\Delta(\rho c^3)$	$\Delta(E/N)$
$\nu = 4$	0.056 ± 0.005	0.069 ± 0.006	0.013	0.29
$\nu = 2$	0.408 ± 0.001	0.422 ± 0.001	0.014	1.589

density is of the order of that of matter nuclear ($0.32 \pm 0.02 \text{ fm}^{-3}$) [27, 28, 29] when we use the hard-sphere radius $c = 0.64 \text{ fm}$ determined through the equation of state of Jaqaman *et al.* [30] for nuclear matter interacting through a Skyrme effective interaction and adjusted to reproduce energy-per-nucleon and saturation density, i.e., -16 Mev and 0.17 fm^{-3} , respectively.

For fermion hard spheres with $\nu = 2$, e.g., helium three or neutron matter, we find $\rho_l c^3 = 0.408$ at crystallization and $\rho_s c^3 = 0.422$ at melting. These density values agree well with the experimental ($\rho_l c^3 = 0.39$, $\rho_s c^3 = 0.41$) and calculated [31, 32] values for helium three. On the other hand, taking for the hard-core diameter a value of $c = 0.4 \text{ fm}$, and assuming that an additional attractive potential would not change the crystallization density of real neutron matter too much, one could predict a crystallization density for neutron matter of about 6.38 fm^{-3} (or mass density $m\rho = 10.85 \times 10^{15} \text{ g-cm}^{-3}$, with m the neutron mass) which not only agrees with a prediction [33] that pure neutron matter crystallizes when its mass density exceeds $1.5 \times 10^{15} \text{ g-cm}^{-3}$ but also justifies Pandharipande's claim [34] p. 695, of no observation of a crystal phase below 4.2 fm^{-3} . In Table 5 we summarize our freezing transition parameters for fermion hard spheres.




Figure 4: Energy E/N (in units of \hbar^2/mc^2) as a function of ρ/ρ_0 for fermion ($\nu = 2$) hard sphere system and Maxwell double tangent construction. Dots mean crystalline and fluid edges.

5 Conclusions

In summary, we have proposed analytical expressions for crystalline and fluid branches of both boson and fermion quantum hard-sphere systems. We have been able to calculate the fluid-crystal transition parameters for bosons and fermions such as symmetrical nuclear matter and neutron matter. The results support the controversial [34] possibility of crystallization in nuclear matter as well as in neutron matter.

ACKNOWLEDGMENTS

MAS thanks Washington University for hospitality during a sabbatical year. We also acknowledge UNAM-DGAPA-PAPIIT (Mexico), grant # IN106401, and CONACyT (Mexico), grant # 27828 E, for partial support.

References

- [1] J.G. Kirkwood, E.K. Maun, and B.J. Adler, J. Chem. Phys. **18**, 1040 (1950).
- [2] F. London, *Superfluids* Vol. II (Dover, New York, 1964) p 30.
- [3] C.A. Rogers, *Packing and Covering* (Cambridge University Press, Cambridge, UK, 1964); N.J.A. Sloan, Sci. Am. **250**, (Jan., 1984) p. 116.
- [4] W. Lenz, Z. Phys. **56**, 778 (1956).
- [5] M.A. Solís, M. de Llano, and R. Guardiola, Phys. Rev. B **49**, 13201 (1994).
- [6] R.K. Cole, Jr., Phys. Rev. **155**, 114 (1967).
- [7] M.H. Kalos, D. Levesque, and L. Verlet, Phys. Rev. A **9**, 2178 (1974).
- [8] S.Z. Ren and M. de Llano, Eur. J. Phys. **10**, 96 (1989).
- [9] A.L. Fetter and J.D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York, 1971).
- [10] T.D. Lee, K. Huang, and C. N. Yang, Phys. Rev. **106**, 1135 (1957); T. T. Wu, *ibid.* **115**, 1390 (1959).
- [11] G.A. Baker, Jr., Rev. Mod. Phys. **43**, 479 (1971).
- [12] G.A. Baker, Jr., L.P. Benofy, M. Fortes, M. de Llano, S.M. Peltier, and A. Plastino, Phys. Rev. A **26**, 3572 (1982).
- [13] V.C. Aguilera-Navarro, R. Guardiola, C. Keller, M. de Llano, and M. Popovic, Phys. Rev. A **35**, 3901 (1987).
- [14] M.A. Solís, V. C. Aguilera-Navarro, M. de Llano, and R. Guardiola, Phys. Rev. Lett. **59**, 2322 (1987); V.C. Aguilera-Navarro, G.A. Baker, Jr., L.P. Benofy, M. Fortes, and M. de Llano, Phys. Rev. A **36**, 4338 (1987); G.A. Baker, Jr., G. Gutiérrez, and M. de Llano, Ann. Phys. (N.Y.) **153**, 283 (1984).
- [15] C. Keller, S. Z. Ren, M. de Llano, M. A. Solís, and G.A. Baker, Jr., Ann. Phys. (N.Y.) **251**, 64 (1996).
- [16] V.C. Aguilera-Navarro, S. Ho, and M. de Llano, Phys. Rev. A **36**, 5742 (1987).
- [17] S. Giorgini, J. Boronat, and J. Casulleras, Phys. Rev. A **60**, 5129 (1999).

- [18] Z. Hu, M. de Llano, E. Buendía, and R. Guardiola, Phys. Rev. B **43**, 12827 (1991).
- [19] M.A. Solís, R. Guardiola, and M. de Llano, Cond. Matt. Theories VIII, (Plenum Press, 1992).
- [20] R.M. Panoff, in *Recent Progress Many-Body Theories*, Vol. 2, edited by Y. Avishai (Plenum, New York, 1990); R.M. Panoff and J. Carlson, Phys. Rev. Lett. **62**, 1130 (1989); R. M. Panoff (priv. comm.).
- [21] G.A. Baker, Jr., and L.P. Benofy, (priv. comm.).
- [22] A. Fabrocini, S. Fantoni, A. Polls, and S. Rosati, N. Cim. A **56**, 33 (1980)
- [23] G.A. Baker, Jr., L.P. Benofy, M. Fortes, and M. de Llano, Phys. Rev. C **34**, 678 (1986); L.P. Benofy, J.L. Gammel, R.M. Delaney, and G.A. Baker, Jr., Phys. Rev. C **48**, 2685 (1993).
- [24] J.B. Hubbard, L. Masinter, and Z.W. Salsburg, J. Chem. Phys. **51**, 4363 (1969).
- [25] A.R. Denton, P. Nielaba, K. J. Runge, and N.W. Ashcroft, Phys. Rev. Lett. **64**, 1529 (1990).
- [26] J.P. Hansen, D. Levesque, and D. Schiff, Phys. Rev. A **3**, 776 (1971).
- [27] P. W. Anderson and R. G. Palmer, Nature, Phys. Sci. **231**, 145 (1971).
- [28] R. G. Palmer and P. W. Anderson, Phys. Rev. D **9**, 3281 (1974).
- [29] J.W. Clark and N.C. Chao, Nature Physical Science **236**, 37 (1972).
- [30] H. Jaqaman, A. Z. Mekjian, and L. Zamick, Phys. Rev. C **27**, 2782 (1983).
- [31] V.R. Pandharipande, Nuclear Physics **A217**, 1 (1973).
- [32] D. Schiff, Nature, Phys. Sci. **243**, 132 (1973)
- [33] V. Canuto and S.M. Chitre, Nature, Physical Science, **243**, 63 (1973).
- [34] J.W. Clark, in *The Nuclear Many-Body Problem*, Eds. F. Calogero and C.C. Degli Atti (Editrice Compositori, Bologna, 1973) Vol. II, pp. 675-704.