

# Finite size scaling and equation of state for Ising lattices

J.G. García\* and J.A. Gonzalo†  
 Departamento de Física de Materiales C-IV,  
 Universidad Autónoma de Madrid.  
 Cantoblanco, 28049 Madrid, Spain.  
 (Dated: May 26, 2020)

Accurate Monte Carlo data from a set of isotherms near the critical point are analyzed using two RG based complementary representations, given respectively in terms of  $\bar{h}=h/|t|^{\beta\delta}$  and  $\bar{\tau}=t/h^{1/\beta\delta}$ . Scaling plots for data on simple cubic Ising lattices are compared with plots of  $ML^{\beta\nu}$  vs.  $|t|L^{1/\nu}$  for increasing  $L$  values and with high quality experimental data on  $CrBr_3$ . Finite size effects and the equation of state are discussed.

PACS numbers: 64.60.-i, 64.60.Cn, 64.90.+b.

Monte Carlo methods [1] using Wolff algorithms [2] have been extensively used to describe  $M(T)$  for  $H=0$  in the whole range of temperatures both below and above the critical temperature (specially at  $T\cong T_c$ ) but few, if any, Monte Carlo simulations of magnetic isotherms  $M(H)$  at  $T\cong T_c$  have been reported in the literature. It is clear, however, that with recent improvements in computing facilities (larger memory, greater speed, better availability) accurate, well thermalized, closely spaced data can provide very substantial contribution to describing the phase transition and better understanding of finite size effects.

Simulations of this type are reported in this work, performed using Metropolis algorithms [3], which are specially convenient to describe the system evolution at constant temperature and for small field increments[4, 5]. Using relatively large lattices with  $70\times 70\times 70$  spins, an accurate characterization of the scaling behavior, and, therefore, the equation of state in the vicinity of the critical point ( $H=0$ ,  $T_c=4.511523785$ ) [6] for simple cubic Ising lattices can be obtained. Details of the Monte Carlo calculations for isotherms taken at  $T$  near  $T_c$  are given in reference [4]. Periodic boundary conditions were used and 140,000 Monte Carlo steps were taken at each field/temperature to ensure equilibrium.

For the scaling representation of the raw  $M(H)$  data at each  $T$  we did use two complementary ways: (a) the usual way [7], involving  $\bar{m}\equiv M(t, h)/|t|^\beta$  and  $\bar{h}\equiv h/|t|^{\beta\delta}$ , where  $t\equiv (T-T_c)/T_c$  involves the temperature gap to  $T_c$ ,  $h$  (reduced field) is proportional to  $H$ ,  $\beta$  is the spontaneous magnetization critical exponent  $\beta\equiv \partial\log M/\partial\log|t|$  at  $T\rightarrow T_c$ , and  $\delta$  is the critical isotherm exponent  $\delta^{-1}\equiv \partial\log M/\partial\log h$  at  $T\rightarrow T_c$ ; and (b) a complementary way involving  $\bar{\mu}\equiv M(h, t)/h^{1/\delta}$  and  $\bar{\tau}\equiv t/h^{1/\beta\delta}$ .

Near the fixed point corresponding to the critical point, the singular part of the reduced free energy per spin may

be written [7] in scaling form as

$$\bar{f}_s(g_1, g_2, g_3, \dots) \sim \bar{b}^d \bar{f}_s(b^{y_1} g_1, b^{y_2} g_2, b^{y_3} g_3, \dots) \quad (1)$$

where  $b$  is an arbitrary scale factor and the  $y_i$  relate to the usual critical exponents. To get (a) Equation (1) is differentiated with respect to the field in the usual way to obtain

$$M(t, h, g_3, \dots) \sim \bar{b}^{-d+y_2} M(b^{y_1} t, b^{y_2} h, b^{y_3} g_3, \dots) \quad (2)$$

Taking  $b^{y_1}|t|=1$  and setting the irrelevant variables equal to zero

$$M(t, h) \sim |t|^{(d-y_2)/y_1} M(\pm 1, h |t|^{-y_2/y_1}) \quad (3)$$

and using the well known scaling relationships [7] giving  $y_1$  and  $y_2$  in terms of  $\beta$  and  $\delta$  one gets

$$M(t, h) \sim |t|^\beta M(\pm 1, h |t|^{-\beta\delta}) \quad (4)$$

which implies scaling using the ordinary scaling variables

$$\bar{m} \equiv M(t, h) / |t|^\beta, \bar{h} \equiv h / |t|^{\beta\delta} \quad (5)$$

To get (b), on the other hand, we can argue in an analogous way. Taking  $b^{y_2}h=1$  in Equation (2) and setting the irrelevant variables equal to zero,

$$M(t, h) \sim h^{(d-y_2)/y_2} M(t \cdot h^{-y_1/y_2}, 1) \quad (6)$$

and using  $y_1$  and  $y_2$  in terms of  $\beta$  and  $\delta$ , one finally gets

$$M(t, h) \sim h^{1/\delta} M(t^{-1/\beta\delta}, 1) \quad (7)$$

which implies as alternative scaling variables

$$\bar{\mu} \equiv M(t, h) / h^{1/\delta}, \bar{\tau} \equiv t / h^{1/\beta\delta} \quad (8)$$

\*Electronic address: jg.garcia@uam.es

†Electronic address: julio.gonzalo@uam.es




FIG. 1: Scaling plots of Monte Carlo data for a s.c. Ising lattice of  $70^3$  spins. (a)  $\bar{m} \equiv M/|t|^\beta$  vs  $\bar{h} \equiv h/|t|^{\beta\delta}$ , and (b)  $\bar{\mu} \equiv M/h^{1/\delta}$  vs  $\bar{\tau} \equiv t/h^{1/\beta\delta}$ . The data include 30 isotherms in the intervals  $4 < T < 5$  and  $0 < h < 0.02$ , the same symbol has been used for all of them. The critical temperature [6] was taken as  $T_c = 4.511523785$  and the critical exponents [4] as  $\beta = 5/16$  and  $\delta = 5$ .

Figure 1(a) gives our Monte Carlo data using  $\bar{m}$  and  $\bar{h}$  as scaling variables as given by Equation (5) with  $T_c = 4.511523785$  and the exponents  $\beta = 5/16 = 0.3125$  and  $\delta = 5$ . The usual behavior is observed. We may note that the data scale extremely well and that only very minor deviations at lower  $h$ , attributable to finite size effects are perceptible. Figure 1(b) gives the same data using the alternative scaling representation. It may be noted immediately that the plot in Figure 1(b) resembles closely [1] typical scaling plots of  $ML^{\beta/\nu}$  vs  $|t|L^{1/\nu}$  for lattices with linear dimension  $L$ , in our case, for Ising systems of  $L^3$  spins, suggesting [8] formal relationships between  $H$  and  $L$ ,  $M$  and  $L$ , and  $|t|$  and  $L$ , which, through the scaling plot branches defining the critical isotherm, the spontaneous magnetization (coexistence curve) below  $T_c$ , and the low field susceptibility above, imply respectively

$$H \sim L^{-\beta\delta/\nu}, M \sim L^{-\beta/\nu} \text{ and } |t| \sim L^{-1/\nu} \quad (9)$$




FIG. 2: Scaling plot of Monte Carlo data  $ML^{\beta/\nu}$  vs  $|t|L^{1/\nu}$  for s.c. Ising lattices with linear size  $L=30, 60, 90, 115$ . Note that finite size effects for  $L=30$  show up closer to  $|t| \rightarrow 0$ .

TABLE I: Critical exponents

	$\beta$	$\delta^{-1}$	$\gamma$
Ising 3D [4]	$5/16=0.3125$	$1/5=0.2$	$5/4=1.25$
Heisenberg 3D [7]	0.340	0.208	1.39
<i>CrBr</i> <sub>3</sub> [9]	0.368	0.233	1.215

Hence

$$ML^{\beta/\nu} \sim \text{const} (t \geq 0) \rightarrow \text{critical isotherm}$$

$$ML^{\beta/\nu} \sim \text{const} \times (|t|L^{1/\nu})^\beta (t < 0) \rightarrow \text{spontaneous magnetization}$$

$$ML^{\beta/\nu} \sim \text{const} \times (|t|L^{1/\nu})^{\beta(\delta-1)/2} (t > 0) \rightarrow \text{susceptibility}$$

Figure 2 shows data of  $ML^{\beta/\nu}$  vs  $|t|L^{1/\nu}$  for  $H=0$  and  $L=30, 60, 90, 115$  which mimic the behavior shown in Figure 1(b) implying that simulations using periodic boundary conditions of phase transitions with finite size show the effects of an effective straining contribution to the magnetic field  $H_{fs} \sim L^{-\beta\delta/\nu}$ , i.e.  $H_{eff} = H + H_{fs}$  such that for  $L \rightarrow \infty$ ,  $H_{eff} \cong H$ .

Figure 3 gives scaling plots of the high quality data of 30 isotherms at the vicinity of the Curie temperature pertaining to the insulating ferromagnet *CrBr*<sub>3</sub> measured by Ho and Litster [9, 10] which are a classical example of experimental scaling data. We show plots of  $\bar{m}$  vs  $\bar{h}$  (Figure 3a) and  $\bar{\mu}$  vs  $\bar{\tau}$  (Figure 3b) with  $T = T_c = 32.844\text{K}$  for two sets of critical exponents, Ising 3d (fractional values) and Ho & Litster (experimental values) summarized in Table I. Both sets of critical exponents produce good scaling plots in (a) as well as in (b), but the critical exponents of Ho and Litster produce somewhat better scaling plots. The accuracy of the magnetization measurements for the set of isotherms was comparable to that of nuclear magnetic resonance data and it was sufficiently precise to establish the form of the scaling function. Our Monte Carlo data, shown in Figures 1(a) and 1(b) are of comparable quality to establish the 3d Ising scaling function.




FIG. 3: Scaling plots of experimental data for  $CrBr_3$  (a)  $\bar{m} \equiv M/|t|^\beta$  vs  $\bar{h} \equiv h/|t|^{\beta\delta}$ , and (b)  $\bar{\mu} \equiv M/h^{1/\delta}$  vs  $\bar{\tau} \equiv t/h^{1/\beta\delta}$ . The data are made up of 30 isotherms in the interval  $T_c - 0.9\text{K} < T < T_c + 6.7\text{K}$ . The critical temperature was  $T_c = 32.844\text{K}$  and the critical exponents used were  $\beta = 5/16$ ,  $\delta = 5$  (Ising 3D) and  $\beta = 0.368$ ,  $\delta = 4.28$ , as given in Reference [9].




FIG. 4: Optimized fits of the 30 Monte Carlo isotherms to the given Ising 3D equation of state.

Finally we address the question of the form equation of state for 3D Ising lattices in the light of the information provided by the set of isotherms in the vicinity of the critical temperature obtained by the Monte Carlo method in our  $70^3$  s.c. lattice. Figure 4 gives the plot of  $M(h, t)$ , Equation (5), rewritten as

$$\frac{\bar{h}}{\bar{m}} = f(\bar{m}) = A(1 \pm B\bar{m}^z)^{(\delta-1)/z} \quad (10)$$

Here  $A$  can be reduced to unity just by choosing properly the units for the field  $H$ .  $B$  is a more meaningful coefficient which, in the particular case of a phase transition describable by means of the mean field approximation (such as the phase transition in a uniaxial ferroelectric) is equal to  $1/\delta = 1/3$ . And  $z$ , as pointed out in Reference [8] is given by  $z \cong \beta\delta/\nu = 2.5$  for  $T < T_c$  and  $z \cong \beta\delta = 1.562$  for  $T > T_c$ . Figure 4 shows the excellent fit obtained by means of Equation (10) with  $(B/A) = 0.102$ . The equation of state put in the form given by Equation (10) is specially good to show directly the most relevant information: (a) the critical isotherm for  $T < T_c$  and  $T > T_c$ , (b) the spontaneous magnetization curve ( $T < T_c$ ) as a vertical line, and (c) the zero field susceptibility ( $T > T_c$ ) as a horizontal line. The quality of the fit is comparable or better than those obtained with traditional expressions of the scaling function [9, 11, 12, 13, 14, 15, 16, 17].

Work is in progress to obtain Monte Carlo data in larger 3D Ising lattices at closer field/temperature intervals, and to extend the investigation of scaling plots in the vicinity of the transition to higher dimensionalities Ising 4D, Ising 5D, etc, in order to monitor closely how the approach to mean field behavior takes place. Of course we will be limited to more reduced sizes (smaller  $L$ 's) as the dimensionality increases, but we have excellent experimental data [18] on a complete set of isotherms in uniaxial ferroelectric TGS to produce excellent scaling plots with  $T_c = 321.470$  and mean field critical exponents  $\beta = 1/2$  and  $\delta = 3$ .

## Acknowledgments

We specially acknowledge helpful comments and software by M.I. Marqués. Support from the Spanish MECyD through Grant Number BFM2000-0032 is gratefully acknowledged.

- 
- [1] K. Binder, in *Phase Transitions and Critical Phenomena*, Vol. 5b (Academic Press, London, 1976).
- [2] U. Wolff, *Phys. Rev. Lett.* **62**, 361 (1989).
- [3] N. Metropolis, A.W. Rosenbluth, M.N. Rosenbluth, A.H. Teller and E. Teller, *J. Chem. Phys.* **21**, 1087 (1953).
- [4] J. García and J.A. Gonzalo *Physica A* (in press, 2003).
- [5] J. García and J.A. Gonzalo [*cond-mat/0304056*] (to be published, 2003).
- [6] H.W.J. Blöte, L.N. Shchur and A.L. Talapov *Int. J. Mod. Phys. C* **10**, 137 (1999).
- [7] See f.i. J.M. Yeomans, *Statistical Mechanics of Phase Transitions*, (Oxford University Press, 1992).
- [8] M.I. Marqués and J.A. Gonzalo *Physica A* **267**, 165-172 (1999).
- [9] J. Ho and J.D. Litster *Phys. Rev. Lett.* **22**, 603 (1969).
- [10] J. Ho and J.D. Litster *Phys. Rev. B* **2**, 4523-4532 (1970).
- [11] A. Arrot and J.E. Noakes *Phys. Rev. Lett.* **19**, 786 (1967).
- [12] M. Vicentini-Missoni, J.M. Levelt Sengers, M.S. Green *Phys. Rev. Lett.* **22**, 390 (1969).
- [13] S. Milosevic and H.E. Stanley *Phys. Rev. Lett.* **22**, 606 (1969).
- [14] D.S. Gaunt and C. Domb *J. Phys. C* **3**, 1442 (1970).
- [15] J.A. Gonzalo *Phys. Rev. B* **1**, 3125 (1970).
- [16] S. Milosevic and H.E. Stanley *Phys. Rev. B* **6**, 986 (1972).
- [17] S. Milosevic and H.E. Stanley *Phys. Rev. B* **6**, 1002 (1972).
- [18] J.R. Fernandez del Castillo, B. Noheda, N. Cereceda, J.A. Gonzalo, T. Iglesias and J. Przeslawski *Phys. Rev. B* **57**, 805 (1998).