

Applications of Random Matrix Ensembles in Nuclear Systems

Maciej M. Duras

Institute of Physics, Cracow University of Technology, ulica
Podchorążych 1, PL-30084 Cracow, Poland

Email: m.duras@riad.usk.pk.edu.pl

"Quantum Field Theory in Particle and Solid State Physics"; June
2, 2003 – June 6, 2003; Max Planck Institute for the Physics of
Complex Systems, Dresden, Germany.

1 Abstract

The random matrix ensembles (RME), especially Gaussian RME and Ginibre RME, are applied to nuclear systems, molecular systems, and two-dimensional electron systems (Wigner-Dyson electrostatic analogy). Measures of quantum chaos and quantum integrability with respect to eigenenergies of quantum systems are defined and calculated.

2 Introduction

Random Matrix Theory (RMT) studies quantum Hamiltonian operators H which are random matrix variables. Their matrix elements H_{ij} are independent random scalar variables [1, 2, 3, 4, 5, 6, 7, 8]. There were studied among others the following Gaussian Random Matrix ensembles (GRME): orthogonal GOE, unitary GUE, symplectic GSE, as well as circular ensembles: orthogonal COE, unitary

CUE, and symplectic CSE. The choice of ensemble is based on quantum symmetries ascribed to the Hamiltonian H . The Hamiltonian H acts on quantum space V of eigenfunctions. It is assumed that V is N -dimensional Hilbert space $V = F^N$, where the real, complex, or quaternion field $F = \mathbb{R}; \mathbb{C}; \mathbb{H}$, corresponds to GOE, GUE, or GSE, respectively. If the Hamiltonian matrix H is hermitean $H = H^Y$, then the probability density function of H reads:

$$f_H(H) = C_H \exp \left[-\frac{1}{2} \text{Tr}(H^2) \right]; \quad (1)$$

$$C_H = \left(\frac{1}{2} \right)^{N_H} ;$$

$$N_H = N + \frac{1}{2} N(N-1) ;$$

$$\int f_H(H) dH = 1 ;$$

$$dH = \prod_{i=1}^N \prod_{j=i+1}^N dH_{ij}^{(D)} ;$$

$$H_{ij} = (H_{ij}^{(0)} ; \dots ; H_{ij}^{(D-1)}) \in F ;$$

where the parameter assume values $D = 1; 2; 4$; for GOE (N), GUE (N), GSE (N), respectively, and N_H is number of independent matrix elements of hermitean Hamiltonian H . The Hamiltonian H belongs to Lie group of hermitean $N \times N$ F -matrices, and the matrix Haar's measure dH is invariant under transformations from the unitary group $U(N, F)$. The eigenenergies $E_i; i = 1; \dots; N$, of H , are real-valued random variables $E_i = E_i^?$. It was Eugene Wigner who firstly dealt with eigenenergy level repulsion phenomenon studying nuclear spectra [1, 2, 3]. RMT is applicable now in many branches of physics: nuclear physics (slow neutron resonances, highly excited complex nuclei), condensed phase physics (nematic particles, ran-

dom Ising model [spin glasses]), quantum chaos (quantum billiards, quantum dots), disordered mesoscopic systems (transport phenomena), quantum chromodynamics, quantum gravity, field theory.

3 The Ginibre ensembles

Jean Ginibre considered another example of GRME dropping the assumption of hermiticity of Hamiltonians thus defining generic F -valued Hamiltonian K [1, 2, 9, 10]. Hence, K belong to general linear Lie group $GL(N, F)$, and the matrix Haar's measure dK is invariant under transformations from that group. The distribution of K is given by:

$$f_K(K) = C_K \exp\left[-\frac{1}{2} \text{Tr}(K^* K)\right]; \quad (2)$$

$$C_K = \left(\frac{1}{2}\right)^{N_K} ;$$

$$N_K = N^2 ;$$

$$\int f_K(K) dK = 1 ;$$

$$dK = \prod_{i=1}^N \prod_{j=1}^N \prod_{\alpha=0}^{D_Y-1} dK_{ij}^{(\alpha)} ;$$

$$K_{ij} = (K_{ij}^{(0)} ; \dots ; K_{ij}^{(D-1)}) \in F ;$$

where $D = 1; 2; 4$, stands for real, complex, and quaternion Ginibre ensembles, respectively. Therefore, the eigenenergies Z_i of quantum system ascribed to Ginibre ensemble are complex-valued random variables. The eigenenergies $Z_i; i = 1; \dots; N$, of nonhermitean Hamiltonian K are not real-valued random variables $Z_i \notin \mathbb{R}$. Jean Ginibre postulated the following joint probability density function of

random vector of complex eigenvalues $Z_1; \dots; Z_N$ for $N \times N$ Hamiltonian matrices K for $\beta = 2$ [1, 2, 9, 10]:

$$P(z_1; \dots; z_N) = \frac{1}{N!} \prod_{i < j}^N |z_i - z_j|^2 \exp\left(-\sum_{j=1}^N |z_j|^2\right); \quad (3)$$

where z_i are complex-valued sample points ($z_i \in \mathbb{C}$).

We emphasize here Wigner and Dyson's electrostatic analogy. A Coulomb gas of N unit charges moving on complex plane (Gauss's plane) \mathbb{C} is considered. The vectors of positions of charges are z_i and potential energy of the system is:

$$U(z_1; \dots; z_N) = \sum_{i < j}^N \ln |z_i - z_j| + \frac{1}{2} \sum_{i=1}^N |z_i|^2; \quad (4)$$

If gas is in thermodynamical equilibrium at temperature $T = \frac{1}{2k_B}$ ($\beta = \frac{1}{k_B T} = 2$, k_B is Boltzmann's constant), then probability density function of vectors of positions is $P(z_1; \dots; z_N)$ Eq. (3). Therefore, complex eigenenergies Z_i of quantum system are analogous to vectors of positions of charges of Coulomb gas. Moreover, complex-valued spacings 1Z_i of complex eigenenergies of quantum system:

$${}^1Z_i = Z_{i+1} - Z_i; i = 1; \dots; (N-1); \quad (5)$$

are analogous to vectors of relative positions of electric charges. Finally, complex-valued second differences 2Z_i of complex eigenenergies:

$${}^2Z_i = Z_{i+2} - 2Z_{i+1} + Z_i; i = 1; \dots; (N-2); \quad (6)$$

are analogous to vectors of relative positions of vectors of relative positions of electric charges.

The eigenenergies $Z_i = Z(i)$ can be treated as values of function Z of discrete parameter $i = 1; \dots; N$. The "Jacobian" of Z_i reads:

$$\text{Jac}Z_i = \frac{\partial Z_i}{\partial i}, \quad \frac{{}^1Z_i}{{}^1i} = {}^1Z_i: \quad (7)$$

We readily have, that the spacing is a discrete analog of Jacobian, since the indexing parameter i belongs to discrete space of indices $i \in I = \{1; \dots; N\}$. Therefore, the first derivative with respect to i reduces to the first differential quotient. The Hessian is a Jacobian applied to Jacobian. We immediately have the formula for discrete "Hessian" for the eigenenergies Z_i :

$$\text{Hess}Z_i = \frac{\partial^2 Z_i}{\partial i^2}, \quad \frac{{}^2Z_i}{{}^2i} = {}^2Z_i: \quad (8)$$

Thus, the second difference of Z is discrete analog of Hessian of Z . One emphasizes that both "Jacobian" and "Hessian" work on discrete index space I of indices i . The spacing is also a discrete analog of energy slope whereas the second difference corresponds to energy curvature with respect to external parameter "describing parametric evolution" of energy levels [11, 12]. The finite differences of order higher than two are discrete analogs of compositions of "Jacobians" with "Hessians" of Z .

The eigenenergies $E_i; i \in I$, of the hermitean Hamiltonian H are ordered increasingly real-valued random variables. They are values of discrete function $E_i = E(i)$. The first difference of adjacent eigenenergies is:

$${}^1E_i = E_{i+1} - E_i; i = 1; \dots; (N - 1); \quad (9)$$

are analogous to vectors of relative positions of electric charges of one-dimensional Coulomb gas. It is simply the spacing of two adjacent

energies. Real-valued second differences 2E_i of eigenenergies:

$${}^2E_i = E_{i+2} - 2E_{i+1} + E_i; i = 1; \dots; (N - 2); \quad (10)$$

are analogous to vectors of relative positions of vectors of relative positions of charges of one-dimensional Coulomb gas. The 2Z_i have their real parts $\text{Re } {}^2Z_i$, and imaginary parts $\text{Im } {}^2Z_i$, as well as radii (moduli) $|{}^2Z_i|$, and main arguments (angles) $\text{Arg } {}^2Z_i$. 2Z_i are extensions of real-valued second differences:

$${}^2E_i = E_{i+2} - 2E_{i+1} + E_i; i = 1; \dots; (N - 2); \quad (11)$$

of adjacent ordered increasingly real-valued eigenenergies E_i of Hamiltonian H defined for GOE, GUE, GSE, and Poisson ensemble PE (where Poisson ensemble is composed of uncorrelated randomly distributed eigenenergies) [13, 14, 15, 16, 17]. The Jacobian and Hessian operators of energy function $E(i) = E_i$ for these ensembles read:

$$\text{Jac}E_i = \frac{\partial E_i}{\partial i}, \quad \frac{{}^1E_i}{{}^1i} = {}^1E_i; \quad (12)$$

and

$$\text{Hess}E_i = \frac{\partial^2 E_i}{\partial i^2}, \quad \frac{{}^2E_i}{{}^1i^2} = {}^2E_i; \quad (13)$$

The treatment of first and second differences of eigenenergies as discrete analogs of Jacobians and Hessians allows one to consider these eigenenergies as a magnitudes with statistical properties studied in discrete space of indices. The labelling index i of the eigenenergies is an additional variable of "motion", hence the space of indices I augments the space of dynamics of random magnitudes.

References

- [1] F. Haake, *Quantum Signatures of Chaos* (Springer-Verlag, Berlin Heidelberg New York 1990), Chapters 1, 3, 4, 8, pp 1–11, 33–77, 202–213.
- [2] T. Guhr, A. Müller-Groeling and H. A. Weidenmüller: *Phys. Rept.* 299, 189–425 (1998).
- [3] M. L. Mehta, *Random matrices* (Academic Press, Boston 1990), Chapters 1, 2, 9, pp 1–54, 182–193.
- [4] L. E. Reichl, *The Transition to Chaos In Conservative Classical Systems: Quantum Manifestations* (Springer-Verlag, New York, 1992), Chapter 6, p. 248.
- [5] O. Bohigas, in *Proceedings of the Les Houches Summer School on Chaos and Quantum Physics*, (North-Holland, Amsterdam, 1991), p. 89.
- [6] C. E. Porter, *Statistical Theories of Spectra: Fluctuations* (Academic Press, New York, 1965).
- [7] T. A. Brody, J. Flores, J. B. French, P. A. Mello, A. Pandey and S. S. M. Wong, *Rev. Mod. Phys.* 53, 385 (1981).
- [8] C. W. J. Beenakker, *Rev. Mod. Phys.* 69, 731 (1997).
- [9] J. Ginibre, *J. Math. Phys.* 6, 440 (1965).
- [10] M. L. Mehta, *Random matrices* (Academic Press, Boston 1990), Chapter 15, pp 294–310.
- [11] J. Zakrzewski and D. Delande, *Phys. Rev. E* 47, 1650 (1993).
- [12] J. Zakrzewski, D. Delande, and M. Kus, *Phys. Rev. E* 47, 1665 (1993).

- [13] M .M .D uras and K .Sokalski, *Phys. Rev. E* 54, 3142 (1996).
- [14] M .M .D uras, *Finite di erence and finite element distributions in statistical theory of energy levels in quantum systems* (PhD thesis, Jagellonian University, Cracow 1996).
- [15] M .M .D uras and K .Sokalski, *Physica D* 125, 260 (1999).
- [16] M .M .D uras, *Description of Quantum Systems by Random Matrix Ensembles of Large Dimensions*, in *Proceedings of the Sixth International Conference on Squeezed States and Uncertainty Relations*, 24 May–29 May 1999, Naples, Italy (NASA, Greenbelt, Maryland, at press 2000).
- [17] M .M .D uras, *J. Opt. B: Quantum Semiclass. Opt.* 2, 287 (2000).