

# The $sl_2$ loop algebra symmetry of the twisted transfer matrix of the six vertex model at roots of unity

Tetsuo Deguchi

Department of Physics, Ochanomizu University,  
2-1-1 Ohtsuka, Bunkyo-ku, Tokyo 112-8610, Japan

## Abstract

We discuss a family of operators which commute or anti-commute with the twisted transfer matrix of the six-vertex model at  $q$  being roots of unity:  $q^{2N} = 1$ . The operators commute with the Hamiltonian of the XXZ spin chain under the twisted boundary conditions, and they are valid also for the inhomogeneous case. For the case of the anti-periodic boundary conditions, we show explicitly that the operators generate the  $sl_2$  loop algebra in the sector of the total spin operator  $S^z = N/2 \pmod{N}$ . The infinite-dimensional symmetry leads to exponentially-large spectral degeneracies, as shown for the periodic boundary conditions [12]. Furthermore, we derive explicitly the  $sl_2$  loop algebra symmetry for the periodic XXZ spin chain with an odd number of sites in the sector  $S^z = N/2 \pmod{N}$  when  $q$  is a primitive  $N$ -th root of unity with  $N$  odd. Interestingly, in the case of  $N = 3$ , various conjectures of combinatorial formulas for the XXZ spin chain with odd-sites have been given by Stroganov and others. We also note a connection to the spectral degeneracies of the eight-vertex model.

# 1 Introduction

Finite-size spectrum of the XXZ spin chain [1, 2] under the twisted boundary conditions has attracted much interest recently and has been studied numerically or analytically such as by the Bethe ansatz [3, 4, 5, 6, 7, 8]. The XXZ Hamiltonian defined on a ring of  $L$  sites is given by

$$H_{XXZ} = J \sum_{j=1}^L \left( \sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y + \sigma_j^z \sigma_{j+1}^z \right) ; \quad (1)$$

Here  $\sigma_j$  for  $= X; Y; Z$  denotes the Pauli matrix defined on the  $j$ th site. Under the periodic boundary conditions we have  $\sigma_{L+1} = \sigma_1$  for any  $\sigma$ . We now introduce the twisted boundary conditions by

$$\sigma_{L+1} = \exp(i\theta) \sigma_1 ; \quad \sigma_{L+1}^z = \sigma_1^z ; \quad (2)$$

We call the parameter  $\theta$  the twist angle.

There are several physical applications of the twisted boundary conditions. The twist angle corresponds to the magnetic flux threaded through the ring [9]. Taking the variation of eigenvalues under changes of the flux [10], the effective mass has been evaluated exactly for the many-body system of interacting fermions or bosons in one-dimension which is equivalent to the XXZ spin chain [4]. The finite-size spectrum of the XXZ spin chain under the twisted boundary conditions has also been studied from the viewpoint of the finite-size analysis of conformal field theories [3]. Associated with the variation of the ground-state energy, the flows of excited states have been numerically investigated with respect to the twist angle [3, 4, 6, 7]. In the spectral flows, we find several level crossings with large degeneracy. However, it has not been discussed explicitly what kind of symmetry operators correspond to the spectral degeneracies. The question should be interesting in particular from the viewpoint of the violation of the level non-crossing rule as discussed by Heilmann and Lieb [11].

Recently, it is found that the symmetry of the XXZ spin chain becomes enhanced at roots of unity. The  $sl_2$  loop algebra commutes with the XXZ Hamiltonian under the periodic boundary conditions, and the infinite-dimensional algebra leads to many spectral degeneracies whose dimensions increase exponentially with respect to the system size [12]. Let us introduce the parameter  $q$  through the XXZ anisotropy as

$$\Delta = \frac{q + q^{-1}}{2} \quad (3)$$

It is shown that the generators of the  $sl_2$  loop algebra commute or anti-commute with the transfer matrix of the six-vertex model when  $q^{2N} = 1$ , and all the defining relations of the  $sl_2$  loop algebra are explicitly derived for the case in the sector  $S^Z = 0 \pmod{N}$ . Here  $S^Z$  denotes the  $Z$  component of the total spin operator. Several aspects of the  $sl_2$  loop algebra symmetry of the XXZ spin chain has been studied [13, 14, 15, 16, 17, 18, 19]. In particular, its connection to the spectral degeneracies of the transfer matrix of the eight-vertex model has been discussed [16, 17]. There are also some relevant papers [20, 21].

The purpose of the paper is to formulate a family of operators commuting with the XXZ Hamiltonian with the twisted boundary conditions. They may explain the level crossings observed in the spectral flows with respect to the twist angle. Here we generalize the approach given in Ref. [12], and give some extended results. As an illustration, we show that when  $\theta = 0$  and  $q$  is a primitive  $2N$ th root of unity, the operators commuting with the twisted XXZ Hamiltonian generate the  $sl_2$  loop algebra for the sector  $S^Z = N/2 \pmod{N}$ . Furthermore, in the sector  $S^Z = N/2 \pmod{N}$ , we explicitly show the defining relations of the  $sl_2$  loop algebra for the case of the periodic boundary conditions where  $L$  is odd and  $q$  is a primitive  $N$ th root of

unity with odd  $N$ . When  $N = 3$ , it is exactly the case in which various combinatorial formulas were discussed recently [22, 23, 24, 25].

The content of the paper consists of the following: In §2 we review the  $sl_2$  loop algebra symmetry of the periodic XXZ spin chain [12]. In §3 we introduce the transfer matrix of the six-vertex model under the twisted boundary conditions. We also discuss some useful techniques such as the gauge transformation and the crossing symmetry. In §4 we show commutation relations of the twisted transfer matrix with some powers of the quantum group generators, which are fundamental in the paper. We also show those of the inhomogeneous case. In §5 we discuss operators commuting with the twisted transfer matrix when  $q$  is a root of unity. We also discuss the special cases where we can explicitly check the defining relations of  $sl_2$  loop algebra. Finally we note a connection to the eight-vertex model.

## 2 The loop algebra symmetry of the periodic XXZ spin chain

Let us review the  $sl_2$  loop algebra symmetry of the XXZ spin chain under the periodic boundary conditions. Let us introduce the quantum group  $U_q(sl_2)$ . The generators  $S^+$  and  $S^z$  satisfy the defining relations

$$[S^+, S^-] = \frac{q^{2S^z} - q^{-2S^z}}{q - q^{-1}}; \quad [S^z, S^\pm] = \pm S^\pm \quad (4)$$

with the multiplication given by

$$(S^\pm)^n = S^\pm \left( q^{S^z} + q^{-S^z} \right)^{n-1} S^\pm; \quad (S^z)^n = S^z \left( I + I \right)^{n-1} S^z \quad (5)$$

Here, the parameter  $q$  is generic. In fact, we may consider  $U_q(L(sl_2))$ , i.e., the  $q$  analogue of the universal enveloping algebra of the  $sl_2$  loop algebra. For simplicity, however, we only consider  $U_q(sl_2)$  in the paper.

Let  $V$  denote a two-dimensional vector space over  $\mathbb{C}$ . On the  $L$ th tensor product space  $V^{\otimes L}$ , the generators  $S^\pm$  and  $S^z$  are given by [12]

$$\begin{aligned} q^{S^z} &= q^{z=2} \otimes \cdots \otimes q^z \\ S^+ &= \sum_{j=1}^L X_j^L q^{z=2} \otimes \cdots \otimes q^z \otimes \cdots \otimes q^z = \sum_{j=1}^L X_j^L S_j^+ : \end{aligned} \quad (6)$$

Here  $S_j^\pm$  denotes the  $j$ th term with  $j$  in the sum (6). Considering the automorphism of  $U_q(L(sl_2))$ , we introduce the following operators:

$$T^\pm = \sum_{j=1}^L X_j^L q^{z=2} \otimes \cdots \otimes q^z \otimes \cdots \otimes q^z = \sum_{j=1}^L X_j^L T_j^\pm \quad (7)$$

Let us denote by  $S^{(\pm)n}$  and  $T^{(\pm)n}$  the following operators:

$$S^{(\pm)n} = (S^\pm)^n = [n]!; \quad T^{(\pm)n} = (T^\pm)^n = [n]! \quad (8)$$

Here  $n$  is a positive integer, and  $[n]$  and  $[n]!$  denote the  $q$ -integer  $[n] = (q^n - q^{-n})/(q - q^{-1})$  and the  $q$ -factorial  $[n]! = [n][n-1]\cdots[1]$ , respectively. Then, we have

$$\begin{aligned} S^{(\pm)n} &= \sum_{1 \leq j_1 < \cdots < j_n \leq L} X_{j_1}^L \cdots X_{j_n}^L q^{\frac{n}{2}z} \otimes \cdots \otimes q^{\frac{n}{2}z} \otimes \cdots \otimes q^{\frac{(n-2)}{2}z} \\ &\quad q^{\frac{(n-2)}{2}z} \otimes \cdots \otimes q^{\frac{(n-4)}{2}z} \otimes \cdots \otimes q^{\frac{n}{2}z} \otimes \cdots \otimes q^{\frac{n}{2}z} \end{aligned} \quad (9)$$

$$\begin{aligned} T^{(\pm)n} &= \sum_{1 \leq j_1 < \cdots < j_n \leq L} X_{j_1}^L \cdots X_{j_n}^L q^{\frac{n}{2}z} \otimes \cdots \otimes q^{\frac{n}{2}z} \otimes \cdots \otimes q^{\frac{(n-2)}{2}z} \\ &\quad q^{\frac{(n-2)}{2}z} \otimes \cdots \otimes q^{\frac{(n-4)}{2}z} \otimes \cdots \otimes q^{\frac{n}{2}z} \otimes \cdots \otimes q^{\frac{n}{2}z} : \end{aligned} \quad (10)$$

Let the symbol  ${}_{6V}(v)$  denote the transfer matrix of the six-vertex model. We now take the parameter  $q$  a root of unity. We consider the limit of sending  $q$  to a root of unity:  $q^{2N} = 1$ . Then we can show the (anti) commutation relations [12] in the sector of  $S^Z = 0 \pmod{N}$

$$S^{-(N)} {}_{6V}(v) = q^N {}_{6V}(v) S^{-(N)}; \quad T^{-(N)} {}_{6V}(v) = q^N {}_{6V}(v) T^{-(N)} \quad (11)$$

Since the XXZ Hamiltonian  $H_{XXZ}$  is given by the logarithmic derivative of the transfer matrix, we have in the sector  $S^Z = 0 \pmod{N}$

$$[S^{-(N)}; H_{XXZ}] = [T^{-(N)}; H_{XXZ}] = 0: \quad (12)$$

Let us now consider the algebra generated by the operators [12]. When  $q$  is a primitive  $2N$ th root of unity with  $N$  even, or a primitive  $N$ th root of unity with  $N$  odd, we consider the following identification [12]:

$$E_0^+ = S^{-(N)}; E_0 = S^{-(N)}; E_1^+ = T^{-(N)}; E_1 = T^{-(N)}; H_0 = H_1 = \frac{2}{N} S^Z: \quad (13)$$

Using the automorphism

$$(E_0) = E_1; \quad (H_0) = H_1 \quad (14)$$

we may take the following identification:

$$E_0^+ = T^{-(N)}; E_0 = T^{-(N)}; E_1^+ = S^{-(N)}; E_1 = S^{-(N)}; H_0 = H_1 = \frac{2}{N} S^Z: \quad (15)$$

When  $q$  is a primitive  $2N$ th root of unity with  $N$  odd, we may put as follows

$$E_0 = iT^{-(N)}; E_1 = iS^{-(N)}; F_0 = iT^{-(N)}; F_1 = iS^{-(N)}; H_0 = H_1 = \frac{2}{N} S^Z: \quad (16)$$

It is shown in Ref. [12] that the operators  $E_j; H_j$  for  $j = 0; 1$ , satisfy the defining relations of the algebra  $U(\mathfrak{sl}_2)$ . Explicitly they are given by the following:

$$H_0 + H_1 = 0; [H_i; E_j] = a_{ij} E_j; [H_i; F_j] = -a_{ij} F_j; \quad (i; j = 0; 1) \quad (17)$$

$$[E_i; F_j] = \delta_{ij} H_j; \quad (i; j = 0; 1) \quad (18)$$

$$[E_i; [E_i; [E_i; E_j]]] = 0; [F_i; [F_i; [F_i; F_j]]] = 0 \quad (i; j = 0; 1; i \neq j) \quad (19)$$

Here, the Cartan matrix  $(a_{ij})$  of  $A_1^{(1)}$  is defined by

$$\begin{pmatrix} a_{00} & a_{01} \\ a_{10} & a_{11} \end{pmatrix} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix} \quad (20)$$

The relations (17) hold for generic  $q$ , while the higher Serre relations (19) hold if  $q$  is a primitive  $2N$ th root of unity or a primitive  $N$ th root of unity with  $N$  odd [12, 26]. The relation (18) holds if  $q$  is a primitive  $2N$ th root of unity with  $N$  even or a primitive  $N$ th root of unity with  $N$  odd. When  $q$  is a primitive  $2N$ th root of unity with  $N$  odd, we take the identification (16) with the imaginary factors.

### 3 Twisted transfer matrix of the six-vertex model

#### 3.1 Boltzmann weights

Let us consider the configuration around the vertex as shown in Fig. 1. Variables  $a; b; c$  and  $d$  are defined on the edges at the vertex, and they take values 1 or 2. The value 1 corresponds

to a polarization vector that is in the upward or rightward directions, and the value 2 to a polarization vector in the downward or leftward direction. We assign the Boltzmann weight  $X_{bd}^{ac}(u)$  to the configuration in Fig. 1. The weight vanishes unless  $a + c = b + d$ , which we call the 'ice rule' or the 'charge conservation'. All the nonzero Boltzmann weights are given by

$$\begin{aligned} X_{11}^{11}(u) &= X_{22}^{22}(u) = \sinh(2 + u) & X_{21}^{12}(u) &= X_{12}^{21}(u) = \sinh u \\ X_{12}^{12}(u) &= X_{21}^{21}(u) = \sinh 2 \end{aligned} \quad (21)$$

Here  $q = \exp 2$ , and  $u$  is the spectral parameter.

We define operators  $X_j(u)$ s for  $j = 0; 1; \dots; L-1$  by

$$X_j(u) = \sum_{a,b,c,d=1,2} X_{bd}^{ac}(u) I_0 \otimes \dots \otimes I_{j-1} \otimes E_j^{ab} \otimes E_{j+1}^{cd} \otimes I_{j+2} \otimes \dots \otimes I_L \quad (22)$$

where  $I$  denotes the identity matrix and  $E^{ab}$  denotes the matrix

$$E_{c;d}^{ab} = \delta_{a;c} \delta_{b;d} \quad \text{for } c, d = 1, 2 \quad (23)$$

It is easy to show that the operators  $X_j(u)$ s constructed from the Boltzmann weights (21) satisfy the Yang-Baxter equation in the following:

$$X_j(u) X_{j+1}(u+v) X_j(v) = X_{j+1}(v) X_j(u+v) X_{j+1}(u) \quad (24)$$

In terms of the Boltzmann weights the operator relation (24) is written as follows

$$\sum_{i,j} X_{a_1 a_2}^{i j}(u) X_{b_3}^{a_3}(u+v) X_{b_1 b_2}^{j i}(v) = \sum_{i,j} X_{a_2 a_3}^{i j}(v) X_{b_1}^{a_1}(u+v) X_{b_2 b_3}^{j i}(u) \quad (25)$$




Figure 1: Vertex configuration for the Boltzmann weight  $X_{bd}^{ac}(u)$ . The spectral parameter  $u$  corresponds to the angle between the two lines  $b$  to  $c$  and  $d$  to  $a$ , which is important to the Yang-Baxter equation (25).

### 3.2 The six-vertex transfer matrix with the twisted boundary conditions

Recall that the operators  $X_j(u)$ s are acting on the tensor product  $V^{(L+1)} = V_0 \otimes V_1 \otimes \dots \otimes V_L$ . We now define the twisted transfer matrix of the six-vertex model by

$$(u; \tau_0) = \text{tr}_0 \left( q^{\sum_{j=0}^L X_{L-1}(u) \otimes \dots \otimes X_0(u)} \right) \quad (26)$$

where the symbol  $\text{tr}_0$  denotes the trace over the 0th space  $V_0$ .

The logarithmic derivative of the twisted transfer matrix leads to the XXZ Hamiltonian  $H_{XXZ}(\eta)$  under the twisted boundary conditions

$$\begin{aligned} \sinh 2\eta \frac{d}{d\eta} \log \langle u; \eta \rangle_{\eta=0} &= \sum_{j=1}^{N-1} 2\sigma_j^+ \sigma_{j+1}^+ + 2\sigma_j^- \sigma_{j+1}^- + \cosh 2\eta \sum_{j=1}^N \sigma_j^z \sigma_{j+1}^z \\ &+ q^2 \sum_{j=1}^N \sigma_j^+ \sigma_{j+1}^+ + q^2 \sum_{j=1}^N \sigma_j^- \sigma_{j+1}^- + \cosh 2\eta \sum_{j=1}^N \sigma_j^z \sigma_{j+1}^z + L \cosh 2\eta \\ &= H_{XXZ}(\eta) = J + L\eta; \end{aligned} \quad (27)$$

where  $\eta = \cosh 2\eta$  and the twist angle  $\eta$  is related to  $\eta$  through the relation

$$q^2 = \exp(i\eta) \quad (28)$$

We should note that the twisted transfer matrix (26) is different from that of Ref. [12]: the logarithmic derivative of the transfer matrix  $T^{DFM}(\eta; \eta)$  of Ref. [12] leads to the XXZ Hamiltonian under the periodic boundary conditions for any  $\eta$ , while that of (26) gives the twisted XXZ Hamiltonian (27).

Let the symbol  $(12)$  denote the shift operator defined by

$$(12) \begin{pmatrix} 1 \\ e_1 \end{pmatrix} = \begin{pmatrix} 1 \\ e_L \end{pmatrix} \otimes e_1 = \begin{pmatrix} 1 \\ e_L \end{pmatrix} \otimes \begin{pmatrix} 1 \\ e \end{pmatrix} \quad (29)$$

Then, the twisted transfer matrix  $\langle u; \eta \rangle$  gives the twisted shift operator:  $\langle 0; \eta \rangle = \sinh^2 2\eta \langle \eta \rangle$ , where  $\langle \eta \rangle$  is defined by

$$\langle \eta \rangle = q^{\sum_{j=1}^N \sigma_j^z} (12) \begin{pmatrix} 1 \\ e_L \end{pmatrix} (12) \begin{pmatrix} 1 \\ e \end{pmatrix} \otimes \begin{pmatrix} 1 \\ e_L \end{pmatrix} \quad (30)$$

### 3.3 Gauge transformations and the crossing symmetry

We introduce the following transformation on the Boltzmann weights of the six-vertex model [27]:

$$X_{bd}^{ac}(u) \rightarrow \tilde{X}_{bd}^{ac}(u) = a^+ b^- \exp(-(a+b-c-d)u/2) X_{bd}^{ac}(u) \quad (31)$$

Here  $\eta$  is arbitrary and  $\eta = 1$ . We can show that the transformed weights  $\tilde{X}_{bd}^{ac}(u)$ s satisfy the Yang-Baxter equation (25) if  $X_{bd}^{ac}(u)$ s satisfy it. We call (31) a gauge transformation. Let  $\tilde{X}_j(u)$  denote the operator defined by (22) with the  $X_{bd}^{ac}(u)$  replaced with  $\tilde{X}_{bd}^{ac}(u)$ . Then,  $\tilde{X}_j(u)$ s also satisfy the Yang-Baxter equation (24).

Let us discuss the gauge transformation (31) with  $\eta = 1$ . It has two important properties. First, the transfer matrix  $\langle u; \eta \rangle$  is invariant under the gauge transformation due to the charge conservation. Second, when  $\eta = 1$ ,  $\tilde{X}_j(u)$  can be expressed in terms of the generator of the Temperley-Lieb algebra [27]. Let  $\tilde{X}_{bd}^{ac}(u)$  denote the transformed weight  $\tilde{X}_{bd}^{ac}(u)$  with  $\eta = 1$ , respectively. In terms of the Boltzmann weights, we have the following decomposition [27]:

$$X_{bd}^{ac}(u) = \sinh(u+2\eta) r_{a,b} r_{c,d} + \sinh u r_a r_b r_{a,c} r_{b,d} \quad (32)$$

Here  $a$  denotes the conjugate of  $a$ , which is defined by  $1 = \bar{2}$  and  $2 = \bar{1}$ . The quantities  $r_j$  ( $j = 1, 2$ ) are defined by [28]

$$r_1 = i \exp(-\eta); \quad r_2 = -i \exp(-\eta) \quad (33)$$

The weights  $\tilde{X}_{bd}^{ac}(u)$ s have the following symmetry [28]:

$$\tilde{X}_{bd}^{ac}(u) = r_b r_c \tilde{X}_{dc}^{ba}(-2\eta - u) = r_a r_d \tilde{X}_{ad}^{cb}(-2\eta - u) \quad (34)$$

We call it the crossing symmetry.

### 3.4 The Temperley-Lieb decomposition of $X_j(u)$ s

Let us introduce the following operators

$$U_j = \sum_{a,b,c,d=1;2}^X r_a r_b r_c r_d I_0 I_1 \dots I_j I L I E_j^{ab} E_{j+1}^{cd} I_{j+2} \dots I L I \quad (35)$$

The  $U_j$ s satisfy the defining relations of the Temperley-Lieb algebra [29, 30]:

$$(U_j)^2 = Q^{1=2} U_j; \quad U_j U_{j+1} U_j = U_j; \quad U_{j+1} U_j U_{j+1} = U_{j+1}; \quad (36)$$

for  $j = 1; 2; \dots; L-1$ . Here  $Q^{1=2}$  is given by  $Q^{1=2} = (q + q^{-1})$ . We call  $U_j$ s the Temperley-Lieb operators.

In terms of the Temperley-Lieb operators, the operator  $X_j(u)$  can be expressed as follows [30]

$$X_j(u) = \sinh(u + 2) I + \sinh u U_j \quad (37)$$

The decomposition (37) corresponds to (32) for the Boltzmann weights.

The Temperley-Lieb operators  $U_j^+$ s commute with the generators of  $U_q(\mathfrak{sl}_2)$ . We can show  $[S^-, U_j^+] = 0$  for  $j = 0; 1; \dots; L-1$ . It can be important, since the XXZ Hamiltonian can be expressed in terms of  $U_j^+$ s and a boundary term [31].

It is known that the periodic XXZ spin chain does not commute with  $U_q(\mathfrak{sl}_2)$ . In fact, the generators  $S^\pm$  do not commute with the periodic XXZ Hamiltonian, since they are not compatible with the periodic boundary conditions. When  $q$  is a root of unity, however, some powers of  $S^\pm$  can commute with it [31]. As a matter of fact, the observation can be developed much further [12]. We first note that  $U_j^+$ s commute with  $S^\pm$  while they do not with  $T^\pm$ , and  $U_j$ s commute with  $T^\pm$  while they do not with  $S^\pm$ . when  $q^{2N} = 1$ , we can show that the transfer matrix commutes with  $S^\pm(N)$  and  $T^\pm(N)$  simultaneously, and they generate the  $\mathfrak{sl}_2$  loop algebra [12].

## 4 Relations of the transfer matrix for $q$ generic

### 4.1 Decomposition of the transfer matrix

Let us consider the gauge transformations (31) with  $\alpha = 1$  and  $\beta = 1$ . Due to the charge conservation, they do not change any of the matrix elements of the transfer matrix (26). Thus, we have

$$(u; ) = \text{tr}_0 q^{-\frac{z}{2}} X_{L-1}(u) \dots X_1(u) X_0(u) \quad (38)$$

We denote by  $(u; )$  the right hand side of (38). Putting the decomposition (37) for  $X_0(u)$  into  $(u; )$ , and making use of the crossing symmetry (34), we can show the following:

$$(u; ) = ( ) X_{LL}(u) + X_{RR}(u) ( )^{-1}; \quad (39)$$

where the symbols  $X_{LL}(u)$  and  $X_{RR}(u)$  are given by

$$\begin{aligned} X_{LL}(u) &= \sinh(u + 2) X_{L-1}(u) \dots X_1(u) \\ X_{RR}(u) &= (1)^L \sinh(u) X_1(2-u) X_2(2-u) \dots X_L(2-u) \end{aligned} \quad (40)$$

We remark that  $X_j^+(u)$ s commute with  $S^\pm$  while  $X_j(u)$ s commute with  $T^\pm$ . When  $\alpha = 0$ , (39) is reduced to that of the periodic one [12].

## 4.2 Transformations of the twisted shift operator

We can show the following relations of  $(S^-)^n$  and  $(T^-)^n$  for generic  $q$ :

$$\begin{aligned}
 (S^-)^n (S^-)^{-1} &= q^{n \frac{z}{1}} (S^-)^n + q^{(n-1)} [n] (S^-)^{n-1} S_1^- q^{2(S^z - n)} 1 \\
 (S^-)^{-1} (S^-)^n &= q^{n \frac{z}{L}} (S^-)^n + q^{(n-1)} [n] (S^-)^{n-1} S_L^- q^{2(S^z - n)} 1 \\
 (T^-)^n (T^-)^{-1} &= q^{n \frac{z}{1}} (T^-)^n + q^{(n-1)} [n] (T^-)^{n-1} T_1^- q^{2(S^z - n)} 1 \\
 (T^-)^{-1} (T^-)^n &= q^{n \frac{z}{L}} (T^-)^n + q^{(n-1)} [n] (T^-)^{n-1} T_L^- q^{2(S^z - n)} 1
 \end{aligned} \tag{41}$$

Let us briefly discuss the derivation. We first consider the following:

$$\begin{aligned}
 (S^-)^n (S^-)^{-1} &= (S^-)^n (S^-)^{-1} \\
 &= (S^- S_1^-) q^{\frac{z}{1}} + q^2 S_1^- q^{2S^z} q^{\frac{z}{1}} 1^n
 \end{aligned} \tag{42}$$

Here we denote by  $A$  and  $q^2 B$  the first and second terms of the right hand side, respectively. Then we can show the following:

$$A + q^2 B^n = A^n + q^2 \sum_{j=0}^{n-1} A^{n-1-j} B A^j \tag{43}$$

Thus, we can derive the expression (41) through the following:

$$\begin{aligned}
 A^n &= q^{n \frac{z}{1}} S^- S_1^- = q^{n \frac{z}{1}} (S^-)^n q^{(n-1)} [n] (S^-)^{n-1} S_1^- \\
 A^{n-1-j} B A^j &= q^{2(j+1)} q^{n \frac{z}{1}} (S^-)^{n-1} S_1^- q^{2S^z}
 \end{aligned} \tag{44}$$

## 4.3 Relations of the twisted transfer matrix

Using the relations (41), we can derive the following relations of the twisted transfer matrix for generic  $q$ :

$$\begin{aligned}
 (S^-)^n (u; ) &= (u; + n) (S^-)^n \\
 &\quad + q^{(n-1)} [n] (u; + n) (S^-)^{n-1} S_L^- q^{2(S^z - n)} 1 X_{LL}^+(u) \\
 &\quad + q^{(n-1)} [n] X_{RR}^+(u) (u; + n)^{-1} (S^-)^{n-1} S_1^- q^{2(S^z - n)} 1 \\
 (T^-)^n (u; ) &= (u; - n) (T^-)^n \\
 &\quad + q^{(n-1)} [n] (u; - n) (T^-)^{n-1} T_L^- q^{2(S^z - n)} 1 X_{LL}(u) \\
 &\quad + q^{(n-1)} [n] X_{RR}(u) (u; - n)^{-1} (T^-)^{n-1} T_1^- q^{2(S^z - n)} 1
 \end{aligned} \tag{45}$$

Here we note that  $S^-$  commute with  $X_{LL}^+(u)$  and  $X_{RR}^+(u)$ , and also that  $T^-$  commute with  $X_{LL}(u)$  and  $X_{RR}(u)$ . Some different forms of the commutation relations are given in Appendix A.

We remark that some relations of  $S^{(N)}$  with the transfer matrix have been investigated [32].



#### 4.4 Inhomogeneous case of the twisted transfer matrix

We define the inhomogeneous transfer matrix of the six-vertex model under the twisted boundary conditions by

$$(u; ;fw_jg) = \text{tr}_0 q^{\frac{z}{2}} X_{L-1}(u-w_{L-1}) \cdots X_1(u-w_1) X_0(u-w_0) \quad (46)$$

Here  $w_j$ s are called inhomogeneous parameters. The twisted transfer matrix (46) is called inhomogeneous. Under the gauge transformation (31) with  $\lambda = 1$ ,  $(u; ;fw_jg)$  is mapped to  $\sim (u; ;fw_jg)$  as follows

$$(u; ;fw_jg) = V(\lambda) \sim (u; ;fw_jg) V(\lambda)^{-1} \quad (47)$$

where  $V(\lambda)$  is given by the diagonal matrix

$$(V(\lambda))_{b_1 a_1; b_L a_L}^{\lambda} = \exp @ \sum_{j=1}^L w_j a_j^A \lambda^{a_1 b_1 a_2 b_2 \cdots a_L b_L} \quad (48)$$

Let  $(u; ;fw_jg)$  denote the transformed transfer matrices  $\sim (u; ;fw_jg)$  with  $\lambda = 1$ , respectively. We write  $V(\lambda)$  by  $V$ . We thus have

$$(u; ;fw_jg) = V(u; ;fw_jg) V \quad (49)$$

We therefore define  $S$  and  $T$  by

$$S = V^+ S V \quad T = V T V^+ \quad (50)$$

The transfer matrices  $(u; ;fw_jg)$  can be decomposed as

$$(u; ;fw_jg) = (\lambda) X_{LL}(u;fw_jg) + X_{RR}(u;fw_jg) (\lambda)^{-1}; \quad (51)$$

where  $X_{LL}(u;fw_jg)$  and  $X_{RR}(u;fw_jg)$  are given by

$$\begin{aligned} X_{LL}(u;fw_jg) &= \sinh(u+2w_0) X_{L-1}(u-w_{L-1}) \cdots X_1(u-w_1) \\ X_{RR}(u;fw_jg) &= (\lambda)^L \sinh(u-w_0) X_1(2u+w_1) \\ &\quad X_2(2u+w_2) \cdots X_L(2u+w_L) \end{aligned} \quad (52)$$

In the same way as (45) we can show

$$\begin{aligned} (S)^n (u; ;fw_jg) &= (u; +n;fw_jg) (S)^n \\ &\quad + q^{(n-1)} [n] (\lambda + n) (S)^{n-1} S_L q^{2(S^z - n)} (\lambda)^{-1} X_{LL}^+(u;fw_jg) \\ &\quad + q^{(n-1)} [n] X_{RR}^+(u;fw_jg) (\lambda + n)^{-1} (S)^{n-1} S_1 q^{2(S^z - n)} (\lambda)^{-1} \\ (T)^n (u; ;fw_jg) &= (u; -n;fw_jg) (T)^n \\ &\quad + q^{(n-1)} [n] (\lambda - n) (T)^{n-1} T_L q^{2(S^z - n)} (\lambda)^{-1} X_{LL}(u;fw_jg) \\ &\quad + q^{(n-1)} [n] X_{RR}(u;fw_jg) (\lambda - n)^{-1} (T)^{n-1} T_1 q^{2(S^z - n)} (\lambda)^{-1} \end{aligned} \quad (53)$$

Through the relation (49) we can derive commutation or anti-commutation relations for the inhomogeneous twisted transfer matrix  $(u; ;fw_jg)$ . Let  $|k\rangle$  denote such a vector with  $S^z |k\rangle = k |k\rangle$ . Then we have

$$\begin{aligned} (S)^n (u; ;fw_jg) |k\rangle &= (u; +n;fw_jg) (S)^n |k\rangle; \text{ when } q^{2(k-n)} = 1 \\ (T)^n (u; ;fw_jg) |k\rangle &= (u; -n;fw_jg) (T)^n |k\rangle; \text{ when } q^{2(k-n)} = 1 \end{aligned} \quad (54)$$

We note that for the case of  $\lambda = 0$ , the inhomogeneous result was addressed at the end of Ref. [12].

## 5 The loop algebra symmetry

### 5.1 Operators commuting with the twisted transfer matrix

Let us assume that  $q$  is a root of unity:  $q^{2N} = 1$ . We denote by  $|j\rangle$  such a vector that has a fixed  $S^Z$  value and it is equivalent to  $k \bmod N : S^Z = k \bmod N$ . Here  $k$  is an integer or a half-integer. Let  $m$  and  $n$  be two non-negative integers such that  $m \equiv n \pmod{N}$ . Then we have

$$\begin{aligned} S^m T^n (u; \lambda) |j\rangle &= (u; \lambda + m - n) S^m T^n |j\rangle \\ T^m S^n (u; \lambda) |j\rangle &= (u; \lambda - m + n) T^m S^n |j\rangle \end{aligned} \quad (55)$$

The relations (55) are derived from (45). We note that when  $m = n$  the expressions (55) are also valid for the case of  $q$  generic.

Let us denote by  $\ell(n)$  the least non-negative integer which is equivalent to  $n \bmod N$ . In the sector  $S^Z = \ell \pmod{N}$  we have commutation relations  $X(u; p) = (u; p)X$ , where  $X$  s are given by

$$(S^{\ell} T^{\ell+p}) (T^{\ell} S^{\ell+p}); \quad (T^{\ell} S^{\ell-p}) (S^{\ell} T^{\ell-p}); \quad (56)$$

and commutation or anti-commutation relations  $X(u; p) = q^{\ell} (u; p)X$ , where  $X$  s are given by

$$\begin{aligned} (S^{\ell} T^{\ell+p+N}) (T^{\ell} S^{\ell+p}); \quad (S^{\ell} T^{\ell-p}) (T^{\ell} S^{\ell+p+N}) \\ (T^{\ell} S^{\ell-p+N}) (S^{\ell} T^{\ell-p}); \quad (T^{\ell} S^{\ell-p}) (S^{\ell} T^{\ell-p+N}) \end{aligned} \quad (57)$$

Here we assume that  $\ell$  and  $p$  are such integers or half-integers that  $\ell \pm p$  are integers.

In the sector  $S^Z = \ell \pmod{N}$ , the operators (56) and (57) commute with the twisted XXZ Hamiltonian  $H_{XXZ}(\lambda)$  with  $\lambda = p \bmod N$ . We recall that  $H_{XXZ}(\lambda)$  is given by the logarithmic derivative of the twisted transfer matrix  $(u; \lambda)$ . For instance, we can show that  $S^{(N)}$  and  $T^{(N)}$  commute with the anti-periodic XXZ Hamiltonian in the sector  $S^Z = N=2 \pmod{N}$ , when  $q$  is a primitive  $2N$ th root of unity. Putting  $\ell = N=2$  and  $p = N=2$  in (57), we have

$$[S^{(N)}; H_{XXZ}(N=2)] N=2i = [T^{(N)}; H_{XXZ}(N=2)] N=2i = 0; \quad (58)$$

### 5.2 Symmetry operators for the inhomogeneous twisted transfer matrix

From the equations (54), in the sector  $S^Z = \ell \pmod{N}$ , we have the commutation relations  $X(u; p; fw_j g) = (u; p; fw_j g)X$  where  $X$  s are given by

$$(S^{\ell} T^{\ell+p}) (T^{\ell} S^{\ell+p}); \quad (T^{\ell} S^{\ell-p}) (S^{\ell} T^{\ell-p}) \quad (59)$$

and commutation or anti-commutation relations  $X(u; p; fw_j g) = q^{\ell} (u; p; fw_j g)X$  where  $X$  s are given by

$$\begin{aligned} (S^{\ell} T^{\ell+p+N}) (T^{\ell} S^{\ell+p}); \quad (S^{\ell} T^{\ell-p}) (T^{\ell} S^{\ell+p+N}) \\ (T^{\ell} S^{\ell-p+N}) (S^{\ell} T^{\ell-p}); \quad (T^{\ell} S^{\ell-p}) (S^{\ell} T^{\ell-p+N}) \end{aligned} \quad (60)$$

Here we recall that  $\ell$  and  $p$  are such integers or half-integers that  $\ell \pm p$  are integers.

### 5.3 The $sl_2$ loop algebra at $q = 1$

Let us discuss the anti-periodic boundary conditions or the twisted boundary conditions with  $q = 1$ . In the sector  $S^Z = N=2 \pmod{N}$ , we can show explicitly the defining relations of the  $sl_2$  loop algebra. We consider two cases: (i)  $q$  is a primitive  $2N$ th root of unity with  $N$  even ( $L$  is even); (ii)  $q$  is a primitive  $2N$ th root of unity with  $N$  odd ( $L$  is odd).

Let us consider the formula for generic  $q$

$$[S^{+(N)}; S^{-(N)}] = \sum_{j=1}^N \frac{S^{-(N-j)} S^{+(N-j)}}{[j]!} \sum_{k=0}^{j-1} \frac{q^{2S^Z+k} q^{-2S^Z+k}}{q - q^{-1}} \quad (61)$$

Taking the limit  $q^{2N} \rightarrow 1$  in the sector  $S^Z = N=2 \pmod{N}$ , we have the following for the cases (i) and (ii):

$$[S^{+(N)}; S^{-(N)}] = [T^{+(N)}; T^{-(N)}] = \frac{2}{N} S^Z \quad (62)$$

We thus take the following identification for the cases (i) and (ii):

$$E_0^+ = T^{-(N)}; E_0^- = T^{+(N)}; E_1^+ = S^{+(N)}; E_1^- = S^{-(N)}; H_0 = H_1 = \frac{2}{N} S^Z \quad (63)$$

Then, we can show that they satisfy the defining relations of the  $sl_2$  loop algebra: (17), (18) and (19).

### 5.4 The $sl_2$ loop algebra symmetry of the periodic XXZ spin chain with $L$ odd

As an application of the (anti-)commutation relations (45), we discuss the case of the periodic boundary conditions with  $L$  odd, where  $q$  is a primitive  $N$ th root of unity with  $N$  odd. Taking the same identification (63) of the generators, we can show explicitly that they satisfy the defining relations of the  $sl_2$  loop algebra in the sector  $S^Z = N=2 \pmod{N}$ .

For sectors other than  $S^Z = N=2 \pmod{N}$ , we have not explicitly shown the defining relations of the  $sl_2$  loop algebra. However, we have a conjecture that some of the operators given in (57) should generate the  $sl_2$  loop algebra. By a different method, we can show that the spectral degeneracies related to the  $sl_2$  loop algebra also exist in the sectors other than  $S^Z = N=2 \pmod{N}$ . We can derive it from the general result on the spectral degeneracy of the eight-vertex model [16] and through the XXZ limit of the XYZ spin chain. Some details should be discussed elsewhere.

#### Acknowledgements

The author would like to thank A.A. Belavin, C. Kor, K. Kudo, B.M. McCoy, and A. Nishino for their helpful comments. He would also like to thank the organizers: D. A. Maudon, J. Avan, L. Frappat, E. Ragoucy and P. Sorba for their kind invitation to the international workshop on Recent Advances in the Theory of Quantum Integrable Systems (RAQIS03), 25-28 March 2003, LAPTH, Annecy-le-Vieux, France. This work is partially supported by the Grant-in-Aid (No. 14702012).

### A Commutation or anti-commutation relations

We can show the following relations for  $q$  generic:

$$(S^+)^n (S^-)^1 = (S^-)^n + q^{(n-1)} [n] (S^-)^{n-1} S_1^- q^{2(S^Z - 1)} q^{n \frac{Z}{2}}$$

$$\begin{aligned}
(S)^{-1} S^n (S) &= (S)^n + q^{-(n-1)} [h](S)^{n-1} S_L q^{2(S^Z)} - 1 \circ q^{n \frac{Z}{L}} \\
(T)^{-1} T^n (T) &= (T)^n + q^{-(n-1)} [h](T)^{n-1} T_1 q^{2(S^Z)} - 1 \circ q^{n \frac{Z}{L}} \\
(T)^{-1} T^n (T) &= (T)^n + q^{-(n-1)} [h](T)^{n-1} T_L q^{2(S^Z)} - 1 \circ q^{n \frac{Z}{L}}
\end{aligned} \tag{64}$$

By using (64) we have

$$\begin{aligned}
(S)^n (u; ) &= (u; + n) (S)^n \\
q^{-(n-1)} [h](S)^{n-1} S_L q^{2(S^Z - n)} - 1 &= (X)_{LL}^+(u) \\
q^{-(n-1)} [h] X_{RR}^+(u) (S)^{n-1} S_L q^{2(S^Z - n)} - 1 &= (X)^{-1} \\
(T)^n (u; ) &= (u; - n) (T)^n \\
q^{-(n-1)} [h](T)^{n-1} T_1 q^{2(S^Z - n)} - 1 &= (X)_{LL}(u) \\
q^{-(n-1)} [h] X_{RR}(u) (T)^{n-1} T_1 q^{2(S^Z - n)} - 1 &= (X)^{-1}
\end{aligned} \tag{65}$$

## References

- [1] H.A. Bethe, *Zeitschrift für Physik* 71, 205 (1931).
- [2] C.N. Yang and C.P. Yang, *Phys. Rev.* 150, 321 (1966); 150, 327 (1966).
- [3] F.C. A. Icaraz, M. Barber and M. Batchelor, *Ann. Phys.*, NY 182, 280 (1988).
- [4] B.S. Shastri and B. Sutherland, *Phys. Rev. Lett.* 65, 243 (1990).
- [5] V.E. Korepin and A.C.T. Wu, *Int. J. Mod. Phys. B* 5, 497 (1991).
- [6] N. Yu and M. Fowler, *Phys. Rev. B* 46, 14583 (1992).
- [7] R.A. Romer, H.-P. Eckle and B. Sutherland, *Phys. Rev. B* 52, 1656 (1995).
- [8] N. Furuta, H. Itoyama and T. Oota, *Int. J. Mod. Phys. A* 12, 801 (1997).
- [9] N. Byers and C.N. Yang, *Phys. Rev. Lett.* 7, 46 (1961).
- [10] W. Kohn, *Phys. Rev.* 133, A171 (1964).
- [11] O.J. Heinmann and E.H. Lieb, *Ann. N. Acad. Sci.* 172, 583 (1971).
- [12] T. Deguchi, K. Fabricius and B.M. McCoy, *J. Stat. Phys.* 102, 701 (2001).
- [13] K. Fabricius and B.M. McCoy, *J. Stat. Phys.* 103, 647 (2001); *J. Stat. Phys.* 104, 575 (2001).
- [14] C. Kor and B.M. McCoy, *Nucl. Phys. B* 618, 551 (2001).
- [15] K. Fabricius and B.M. McCoy, in *MathPhys Odyssey 2001* edited by M. Kashiwara and T. Miwa, (Birkhauser, Boston), 119 (2002).
- [16] T. Deguchi, *J. Phys. A: Math. Gen.* 35, 879 (2002); *Int. J. Mod. Phys. B* 16, 1899 (2002).
- [17] K. Fabricius and B.M. McCoy, *J. Stat. Phys.* 111, 323 (2003). (cond-mat/0207177.)

- [18] C . K or , J. Phys. A : M ath. G en. 36, 5229 (2003); m ath-ph/0305035.
- [19] T . D eguchi, cond-m at/0212217.
- [20] A A . Belavin and M . J im bo, hep-th/0208224.
- [21] A A . Belavin, hep-th/0305209.
- [22] Yu. Stroganov, J. Phys. A : M ath. G en. 34, L179 (2001).
- [23] A V . Razum ov and Yu. Stroganov, J. Phys. A : M ath. G en. 34, 3185 (2001).
- [24] M . T . Batchelor, J. de G ier and B . N ienhuis, J. Phys. A : M ath. G en. 34, L265 (2001).
- [25] J. de G ier, M . T . Batchelor, B . N ienhuis and S. M itra, cond-m at/0110011.
- [26] G . Lusztig, Introduction to Q uantum G r oups (Birkhauser, 1993).
- [27] Y . A kutsu and M . W adati, J. Phys. Soc. Jpn. 56, 3037 (1987).
- [28] T . D eguchi, M . W adati and Y . A kutsu, J. Phys. Soc. Jpn. 57, 1905 (1988).
- [29] H N V . Tem perly and E H . Lieb, P roc. R . Soc. London Ser. A 322 (1971) 251.
- [30] R J . B axter, J. Stat. Phys. 28 (1982) 1.
- [31] V . Pasquier and H . Saleur, Nucl. Phys. B 330, 523 (1990).
- [32] A A . Belavin, S. Yu. G ubanov, Theor. M ath. Phys. 129, 1484 (2001).