QUANTUM MODEL OF PHONON TRANSPORT AND HEAT CONDUCTIVITY IN CARBON NANOCLUSTERS AND NANOTUBES

April 14, 2024

E Ya G lushko (1 and 2), V N Evteev (3), M N M oiseenko (3), N A Slusarenko (3) and A A Zakhidov (4)

((1) Institute for Sem iconductors Physics NAS Ukraine, PrNauki. 21, Kiev, 03047, Ukraine; (2) Slavonic University, Anry Barbusse Str. 9, Kiev 03150, Ukraine; (3) Pedagogical University, Gagarin Avenue 54, Krivoy Rog, 50086, Ukraine; (4) Texas University at Dallas, Richardson, TX 75083-0688, USA)

(E-m ail: eugene.glushko@ scientist.com)

Abstract

A complex approach phonon quantum discrete model (PQDM) was developed to describe dynamics, kinetics and statistics of phonons in carbon nanostructures with zero-chirality of both zig-zag and arm chair geometry. The model allows include into the pure phonon problem existing interaction with others subsystems: electrons, photons, in purities and defects. We predict that planar C-structures are geometrically stable and may bridge interelectrode space in strong external electric

eld. The exact solution of generalized therm al conductivity (TC) equation was obtained for nanotubes. Tem perature distribution along the tube was derived analytically. The diagonalization procedure for the case ofstrong ph-ph interaction was proposed. It was shown the quadratic increasing of heat conductivity with the grow th of the phonon m ean free path (PM FP). Heat capacitance and the entropy of carbon linear tubes were calculated as the function of tem perature. Our theoretical approach explains the nature of good TC in carbon and carbon-like materials by existing of the soft vibration branch (low frequency radial breathing m ode phonons with high density of states at them al energies) accompanied by structure hardness (high frequency ' - and z-branches) providing big PM FP. TC coe cient for high conducting channel in surrounding medium was calculated. The mechanism of heat conductivity tem perature dam ping was analyzed. Two com petitive tendencies produce TC maximum at interm ediate tem peratures (100-300)K. It was shown the strongly non-linear increasing of e ective heat conductivity with the growth nanotubes concentration. It was shown that insertion of arm chair nanotube inside a m edium or its coating by polyacetilene m olecule considerably changes the structure of radial breathing phonons.

I.M otivation

II. Introduction

III. System s under consideration

 ${\rm IV}\,.\,{\rm P\,honon}$ quantum discrete model (PQDM)

V.V ibrational dynam ics. Linear approximation.

VI.Frequency spectrum and DOS.Zig-zag geometry.

VII.Frequency spectrum and DOS.Am chair geometry.

V III. Am plitude distribution and knot's theorem IX. Heat transfer X. Phonon mean free path and size dependences X I. Thermal coe cient temperature dependence X II. G eneralized phonon transport equation

X III. Sum m ary and discussion

1 M otivation

Phonon dynamics in open and closed carbon and carbon-like nets. How many phonon branches are present? W hether exists a simple correlation between di ered phonon types for di erent NT geometries?

V ibrational eigenm ode and its am plitudes distribution in the net.

V ibrational eigenm odes, phonons, m icro-sound and heat propagation. Phonon m ean free path. Sound waves boundary conditions.

The mechanism of temperature damping at high temperatures and concrete law of increasing at low temperatures.

M axim altherm al conductivity for solitary carbon tube is not measured reliably now and changes in wide interval from 200 W / m K to 3000 W / m K for dierent authors.

The tem perature of maxim altherm alconductivity for carbon tubes is not surely measured also and may change in wide interval from 150 K to 300 K for di erent authors.

The question is whether exist di erences in them al conductivity of solitary carbon tubes depending on type (arm chair or zig-zag), chirality and diam eter. W hich physical factors are essential in electron transport through the m olecule?

D o phonons (vibrons) participate essentially in electron transport through them olecule? A re di erences in therm al conductivity of solitary carbon tubes, at carbon structures like graphene and graphite (in plane) essential?

2 IN TRODUCTION

Fig 2.1 G eneral picture of connected subjects for the problem of phonon propagation in molecular nets.

Fig 3.1 A polymerm olecule absorbed by an chair nanotube. Absorption bonds are shown by red lines, C-C bonds of the polymer are shown by violet color.

Fig.3.2 Heat transistor (3-polar contact) on a $\;$ at carbon structure connecting three baths at di erent temperatures T_1 , T_u and T_r .

4 Phonon quantum discrete model (PQDM)

$$\hat{H}_{0} = \sum_{s;}^{X} h!_{s} (\hat{b}_{s}^{+} \hat{b}_{s} + 1=2)$$
(1)

$$\hat{\mathbf{H}}_{H} = {}^{P} \mathbf{G}_{i_{1}} (\hat{\mathbf{b}}_{i_{1}}^{+} \hat{\mathbf{b}}_{1} + \hat{\mathbf{b}}_{1}^{+} \hat{\mathbf{b}}_{i_{1}}) + + {}^{i_{1}} \mathbf{G}_{i_{r}} (\hat{\mathbf{b}}_{i_{r}}^{+} \hat{\mathbf{b}}_{r} + \hat{\mathbf{b}}_{r}^{+} \hat{\mathbf{b}}_{r})$$

$$(2)$$

5 Vibrational dynam ics. Linear approximation. Zig-zag NT

$$m_{i} = k^{0} (3_{i} \quad i1_{i2} \quad i3)$$
 (3)

$$m z_i = k (z_i z_{i1}) \quad 0.25k (2z_i z_{i2} z_{i3})$$
 (4)

$$m x_{i} = 0.75k (2x_{i} x_{i1} x_{i2}) k^{0} (x_{i} x_{i3})$$
(5)

A m chair N T

$$m z_i = 0.75k (2z_i z_{i1} z_{i2}) k^0 (z_i z_{i3})$$
 (6)

$$m x_i = k (x_i x_{i1}) \quad 0.25k (2x_i x_{i2} x_{i3})$$
 (7)

$$\hat{D}\tilde{A} = !^{2}\tilde{A}$$
(8)

$$!_{s}; \mathcal{A}_{s}^{+} = (C_{s1}; C_{s2}; C_{s3} ::: C_{sr}); r = 2n (m + 1)$$
(9)

Fig.5.1 The problem 's geometry. Z-axes is perpendicular to the gure plane. It's shown j and r shifts of an atom .

Fig.5.2. The problem 's geometry. F is z-axes projection of the force acting between a and b atoms. dz marks the atom a shift along the axes z, $dz\cos(p/3)$ is the shift projection.

Fig.5.3. Sketch of the dynam icalm atrix for zig-zag tubulene created from a graphene sheet f5,5g containing 60 atoms by its rolling up. Rose-colored spots are diagonal elements, green and blue show two types of bonds, rest elements equal to zero, left and upper red bands contain atom ic numbers.

Fig.5.4. Sketch of the dynam icalm atrix for arm chair tubulene created from a graphene sheet f5,5g containing 60 atoms by its rolling up. Rose-colored spots are diagonal elements, green and blue show two types of bonds, rest elements equal to zero, left and upper red bands contain atom ic num bers.

6 Frequency spectrum and DOS.Zig-zag geometry.

$$g(!_{s}) = \frac{1}{!_{s} !_{s}!_{s}}$$
 (10)

Fig.6.1. Radial mode spectrum of zig-zag tubulene created from a graphene sheet $f_{5,15g}$ by rolling up around z-axes. The unity of frequency is $!_0$

Fig.6.2 Radialm ode DOS function for Fig.6.1 system . h = Int[35r/50].

Fig.6.3 A zim utb m ode spectrum of zig-zag tubulene created from a graphene sheet f5,15g by rolling up around z-axes. The unity of frequency is $!_0$.

Fig.6.4 A zim uthal mode DOS function (arbitrary units) for graphene sheet f5,15g h= Int[35r/50].

•

Fig.6.5 A xialm ode spectrum of zig-zag tubulene created from a graphene sheet f5,15g by rolling up around z-axes. The unity of frequency is $!_{\,0}\,$.

Fig.6.6 A xialm ode DOS function (arbitrary units) for tubulene fragment created from a graphene sheet f5,15g by rolling around z-axes. Horizontal axes, frequency in $!_0$

7 Frequency spectrum and DOS.Arm chair geometry.

Fig.7.1. Radial mode density of states for an chair tubulene created from a graphene sheet f8,15g by rolling up around z-axes. Insertion the same for zig-zag tubulene.

Fig.7.2 Tangential jm ode density of states for an chair tubulene created from a graphene sheet f8,15g by rolling up around z-axes. Insertion is zig-zag equivalent for this that is z-m ode DOS.h= Int[4r/5].

•

Fig.7.3 A xialz-branch density of states for an chair tubulene created from a graphene sheet f8,15g by rolling up around z-axes. Insertion is zig-zag equivalent for this that is '-m ode DOS.

Fig.7.4 Low erpart. Three branches of phonon spectrum for am chair tubulene created from a graphene sheet f8,15g by rolling up around z-axes. Upper part. Radial phonon band for zig-zag f8,15g NT.

8 Amplitude distribution and knot's theorem

•

$$A_{j} = \sum_{s;}^{X} j c_{sj} j n_{s};$$
 (11)

Fig.8.1 Calculated distribution of vibrational amplitudes of f8,8g planar carbon net for the 4-th m ode.

Fig.8.2. Calculated r-branch amplitude distribution s= 5 along arm chair tubulene created from a graphene sheet f6,5g by rolling up around marked direction. Transversal knot's lines are degenerated.

F ig 8.3 r-branch amplitude distribution along zig-zag tubulene created from a graphene sheet f6,5g by rolling up around m arked direction. 5th state. Longitudinal knot's lines are degenerated.

F ig 8.4 C alculated m can square am plitude distribution along zig-zag tubulene created from a graphene sheet 15,5 by rolling up around m arked direction. $T = 0.03 \text{ eV} \cdot \text{C}$ incular arrows show rolling up of the structure.

9 Heat transfer

•

$$W_{-1! s} = \frac{2}{h} \ \mathbf{j} \mathbf{G}_{1s} \ \mathbf{j}^{2} \ \mathbf{g}_{1} \ (! s) \ \mathbf{N}_{1} \ (! s) \ (1 + n \ (! s))$$
(12)

$$n_{s} = \frac{j_{i_{1}}^{P} G_{i_{1},s} j cdotg_{1}(!_{s}) N_{1}(!_{s}) + j_{i_{r}}^{P} G_{i_{r},s} j g(!_{s}) N_{1}(!_{s})}{j_{i_{1}}^{P} G_{i_{1},s} j g(!_{s}) + j_{i_{r}}^{P} G_{i_{r},s} j g(!_{s})}$$
(13)

for 3-polar contact

$$n_{s} = (j_{i_{1}}^{X} G_{i_{1},s} j_{1}^{2} g(!_{s}) N(!_{s}) + j_{1}^{X} G_{i_{u},s} j_{1}^{2} g(!_{s}) N_{u}(!_{s}) + j_{1}^{i_{u}} G_{i_{d},s} j_{1}^{2} g(!_{s}) N_{u}(!_{s}) + j_{1}^{X} G_{i_{d},s} j_{1}^{2} g(!_{s}) N_{u}(!_{s})) = (j_{i_{1}}^{X} G_{i_{1},s} j_{1}^{2} g(!_{s}) + j_{1}^{X} G_{i_{u},s} j_{1}^{2} g(!_{s}) + j_{1}^{X} G_{i_{d},s} j_{1}^{2} g(!_{s}))$$
(14)

And for 4-polar contact

$$Q = Qt = 2 \sum_{s;}^{X} j_{g_{1s}}^{2} j_{g_{rs}}^{2} j_{s}^{2} g_{1}(!_{s}) g_{r}(!_{s})$$

$$\frac{N_{1}(!_{s}) N_{r}(!_{s})}{j_{g_{1s}}^{2} g_{1}(!_{s}) + j_{g_{rs}}^{2} g_{r}(!_{s})}$$
(16)

$$^{\circ} = L \frac{@Q = @t}{T_r T_1}$$
(17)

Approximation

$$N_{1}(!_{s}) = N_{r}(!_{s}) = \frac{\Theta N(!_{s})}{\Theta T} (T_{r} = T_{1})$$
 (18)

$${}^{0}(L) = 2 L \sum_{s;}^{X} j_{g_{1s}} j_{g_{rs}}^{2} j_{g_{rs}}^{2} j_{s}^{2} g_{1}(!_{s}) g_{r}(!_{s})$$

$$\frac{0N (!_{s}) = 0T}{j_{g_{1s}} j_{g_{1}}^{2}(!_{s}) + j_{g_{rs}} j_{g_{r}}^{2}(!_{s})}$$
(19)

10 Phonon m ean free path and size dependences

$${}^{0}(\mathbf{l}_{ph}) = \frac{2 \, \mathbf{l}_{ph}}{T^{2}} \frac{X}{s} \, \mathbf{j}_{B} \, \mathbf{j}_{S}^{2} \, \mathbf{j}_{G} \, \mathbf{j}_{S}^{2} \, \mathbf{j}_{S}^{2} \, \mathbf{g}(\mathbf{l}_{S}) \frac{N \, (\mathbf{l}_{S}) (N \, (\mathbf{l}_{S}) + 1)}{\mathbf{j}_{B} \, \mathbf{j}_{S}^{2} + \mathbf{j}_{G} \, \mathbf{r}_{S}^{2}}$$

$$(20)$$

$$G_{rs} = \sum_{i_r}^{X} G_{i_r} C_{i_r s}$$
 (21)

The mean free path along the tube lph.

$$l_{ph} \qquad \frac{s}{N} = \frac{s}{N} \frac{l_{d}}{l_{d}} \frac{ph}{ph}; \qquad l_{ph} \qquad \frac{l_{d}}{< N >}$$
(22)

$$< N > = \frac{1}{2} n (!)g(!)d!; \qquad l_{ph} = \frac{1}{T}$$
 (23)

Fig.10.1. One dimensional phonon dynamics. Calculated total therm al conductivity length dependence that includes all vibration branches of arm chair NT. Lph is phonon mean free path, a is C-bond length.

•

11 Therm alcoe cient tem perature dependence

$$_{0} = \frac{2 \ \text{aG}_{0}^{2}}{h!_{0}} k \tag{24}$$

Fig.11.1 . Them al conductivity temperature dependence. A m chair NT.Curve 1 corresponds to z-branch contribution, curve 2 to jbranch and r-branch contribution is presented by curve 3.

Fig.112 Therm al conductivity temperature dependence. Zig-zag NT.Curve 1 corresponds to z-branch contribution, curve 2 to jbranch and r-branch contribution is presented by curve 3.

12 X II. G eneralized phonon transport equation

$$W_{-1! s} = \frac{2}{h} \mathcal{G}_{1s} \mathcal{J}^{2} g_{1}(!_{s}) N_{1}(!_{s}) (1 + n (!_{s}))$$
(25)

$$\frac{\partial n_{s}}{\partial t} = D_{z}^{s} \frac{\partial^{2} n_{s}}{\partial z^{2}}$$
(26)

$$D_{z}^{s} = \frac{2 l_{ph}}{h} jG_{ls} j^{2}g(!_{s})$$
(27)

$$G_{ls} = G_{rs} = {\stackrel{X}{\underset{i_{l}}{G}}} G_{i_{l}}C_{i_{l}s}$$
 (28)

Tem perature distribution along the tube

•

$$T(z) = !_{s} = Ln 1 + \frac{L}{n_{1}(!_{s}) + [n_{r}(!_{s}) - n_{1}(!_{s})]z}$$
(29)

Fig.12.1 Phonon jumps over a carbon and chair structure (arrows). Nanotubes are ideal one-dimensional systems as to the phonon dynamics.

•

Fig.12.2. One dimensional phonon dynamics. Exact temperature distribution along the tube given by (4.4). Tl, Tr are temperatures of baths connecting with tube ends, L is the tube length. End points come together all modal (partial) temperatures for each ss.

Fig.13.1. A veraging procedures and speed of propagation. a is interatom ic distance, \ll is mean atom ic shift, x is extended coordinate of the freedom 's degree, V is excitation velocity, m is atom mass, k is elasticity coe cient.

Fig.13.2. Tem perature dependence of therm alconductivity $\cos c$ ient. Two $\cos p$ etitive tendencies.

The boundary conditions for m icrosound spreading through periodic NT com plexes

$$0:75ka\frac{\theta < z>}{\theta z} + (1)^{l;r}m!^2 < z > = const$$

$$\frac{\theta < z>}{\theta z} = const$$
(30)

$$\langle z \rangle = A e^{ifz} + B e^{-ifz}$$
 (31)

A complex approach PQDM was applied to describe phonon dynamics and heat spreading processes in carbon NT-solid medium composite material and in nanotubes embraced by a polymer molecule.

A tom ic vibration dynam ics was considered for carbon zero-hirality nanotubes connected with solid matrix. V ibrational eigenmodes, density of states and amplitude distribution for tubes inserted into a cylindrical well inside the medium and bounded with its hard inner wallwere calculated. P honon band parameters were analyzed in linear approximation for three types of vibration: athim uthal or tangential jbranch, radial breathing mode (r-branch) and longitudinal z-branch.

It's shown that phonon propagation in actual nanotubes is characterized by a kind of "com pacti cation" of circular freedom 's degree due to the big phonon m ean free path. N anotubes of actual diam eters are ideal one-dim ensional phonon qnd heat conductors. Phonon band structure was investigated for an chair nanotubes on the base of hierarchical law and system symmetry.

Therm all uxes and therm alconductivity were considered in PQDM. Tem perature dependences were obtained. The mechanism of heat conductivity high tem perature dam ping is cleared. Two competitive tendencies produce therm alconductivity maximum at intermediate tem peratures (100-300)K.

The exact solution of generalized them alconductivity equation was obtained for nanotubes. Tem perature distribution along the tube was derived analytically.

Size dependences were considered for therm al conductivity. It was shown the linear increasing of heat conductivity with the growth of the phonon mean free path.

The explanation of the nature of good therm al conductivity in carbon and carbonlike materials by existing of the soft vibration branch (low frequency RBM phonons with high DOS at therm all energies) accompanied by structure hardness (high frequency j-and z-branches) providing big mean free path for phonons (Fig.5.2).

Adding of new layers or new walls to single-walled NT makes breathing r-branch of vibrations harder and causes the sharp decreasing of phonon density of states at the same phonon mean free path

P ressure decreases therm alconductivity. The e ect is connected with total hardening of all bonds and phonon m odes going away from active therm al zone.

M elting decreases therm al conductivity by another reason: the phonon's mean free path becom es sm all.

M odi ed Fourier equation of heat propagation in m icroscopic m olecular channel was obtained.

Boundary problem for tem perature distribution in channel-medium system was considered in cylindrical symmetry. The method of partial solution building for cylindrical boundary problem was proposed. Tem perature distribution isotherm swere calculated.

Concentration dependences were considered for therm alconductivity of NT-solid matrix composite. It was shown the strongly non-linear increasing of elective heat conductivity with the growth nanotubes concentration.

Phonon dynamics was considered for carbon zero-chirality nanotubes embraced by polymermolecule. V ibrational eigenmodes, density of states and amplitude distribution for a bounded tube were calculated. Phonon band parameters were obtained in linear approximation for three types of vibration.

It was shown that insertion of arm chair nanotube inside a medium or its coating by polyacetilene molecule considerably changes the structure of radial breathing phonons

This figure "fig_10_1.gif" is available in "gif" format from:

This figure "fig_11_1.gif" is available in "gif" format from:

This figure "fig_12_1.gif" is available in "gif" format from:

This figure "fig_13_1.gif" is available in "gif" format from:

This figure "fig_2_1.gif" is available in "gif" format from:

This figure "fig_3_1.gif" is available in "gif" format from:

This figure "fig_5_1.gif" is available in "gif" format from:

This figure "fig_6_1.gif" is available in "gif" format from:

This figure "fig_7_1.gif" is available in "gif" format from:

This figure "fig_8_1.gif" is available in "gif" format from:

This figure "fig_11_2.gif" is available in "gif" format from:

This figure "fig_12_2.gif" is available in "gif" format from:

This figure "fig_13_2.gif" is available in "gif" format from:

This figure "fig_3_2.gif" is available in "gif" format from:

This figure "fig_5_2.gif" is available in "gif" format from:

This figure "fig_6_2.gif" is available in "gif" format from:

This figure "fig_7_2.gif" is available in "gif" format from:

This figure "fig_8_2.gif" is available in "gif" format from:

This figure "fig_5_3.gif" is available in "gif" format from:

This figure "fig_6_3.gif" is available in "gif" format from:

This figure "fig_7_3.gif" is available in "gif" format from:

This figure "fig_8_3.gif" is available in "gif" format from:

This figure "fig_5_4.gif" is available in "gif" format from:

This figure "fig_6_4.gif" is available in "gif" format from:

This figure "fig_7_4.gif" is available in "gif" format from:

This figure "fig_8_4.gif" is available in "gif" format from:

This figure "fig_6_5.gif" is available in "gif" format from:

This figure "fig_6_6.gif" is available in "gif" format from: