

# Vortex formation in a fast rotating Bose-Einstein condensate

Tarun Kanti Ghosh

*The Abdus Salam International Centre for Theoretical Physics, Strada Costiera 11, 34100 Trieste, Italy.*  
(May 25, 2020)

We study rotational motion of an interacting atomic Bose-Einstein condensate confined in a quadratic-plus-quartic potential. We calculate the lowest energy surface mode frequency and show that a symmetric trapped (harmonic and quartic) Bose-Einstein condensate breaks the rotational symmetry of the Hamiltonian when rotational frequency is greater than one-half of the lowest energy surface mode frequency. We argue that the formation of a vortex is not possible in a non-interacting as well as in an attractive Bose-Einstein condensate confined in a harmonic trap due to the absence of the spontaneous shape deformation, but it can occur which leads to the vortex formation if we add an additional quartic potential. Moreover, the spontaneous shape deformation and consequently the formation of a vortex in an attractive system depends on the strengths of the two-body interaction and the quartic potential.

PACS numbers: 03.75.Lm, 05.30.Jp

## I. INTRODUCTION

After the creation and realization of the Bose-Einstein condensate (BEC) in an alkali atomic trapped Bose gas, a new field, namely strongly correlated atomic system, has been opened up [1]. It is always interesting to study the consequences of an external perturbation to the system. One way to perturb the system is by applying external rotation to the system. The rotation of a macroscopic quantum fluid exhibits interesting counter-intuitive phenomena. For example, an atomic interacting(repulsive) BEC confined in a rotating harmonic trap produces quantized vortices for a sufficiently large rotation [2–4]. The theoretical calculation for critical rotation frequency ( $\Omega_c$ ) based on purely thermodynamic arguments [5] is significantly smaller than the observed value of  $\Omega_c$ . For sufficiently high stirring frequencies of the trap, the system deforms spontaneously [6] to lower the energy barrier which prevents the nucleation of vortices at smaller  $\Omega$  [7] and this spontaneous deformation is detected at the ENS experiment which leads to the vortex nucleation [8]. In fact, vortices are nucleated when the stirring potential with low anisotropies resonantly excites the quadrupole mode of the condensate, and induces large amplitudes oscillations of the condensate, resulting in a dynamical instability [9].

The harmonically trapped BEC becomes singular when the rotation frequency is equal to or greater than the harmonic trap frequency, because the outward centrifugal force counteracts the inward force from the harmonic trap. We can eliminate the singularity at  $\Omega \geq 1$  if we consider an additional stiffer radial potential (say, quartic potential for simplicity). There is a growing interest about the effect of an anharmonic potential on the properties of a rotating BEC [10–14]. Recently, the quadratic-plus-quartic potential has been achieved experimentally at ENS by superimposing a blue detuned laser

beam to the magnetic harmonic trap [15].

In spite of the fact that the  $\Omega_c$  in a harmonically trapped BEC calculated from the thermodynamic arguments does not match with the experimental values, many authors have studied  $\Omega_c$  in a BEC confined in harmonic-plus-quartic trap based on the thermodynamic arguments [11–14]. In this work, we argue that a vortex formation is possible in an attractive as well as non-interacting Bose gas confined in a quadratic-plus-quartic potential and calculate the critical rotational frequencies based on the spontaneous shape deformation of rotating Bose gas.

This paper is organised as follows. In section II, we write down the model Hamiltonian and variational macroscopic wave function, from which we calculate the variational energy. By minimizing the variational energy, we derive three coupled equations of the variational parameters. In section III, we calculate the lowest energy surface mode frequency of an interacting BEC confined in a quadratic-plus-quartic potential. In section IV, we discuss how spontaneous shape deformation transfer angular momentum and vortex nucleation occurs in a harmonically trapped BEC. In section V, we discuss the possibility of the shape deformation and vortex nucleation of a BEC trapped in a quadratic-plus-quartic potential. We present summary and conclusions of our work in section VI.

## II. THE SYSTEM AND THE HAMILTONIAN

The equation of motion of the condensate wave function is described by the mean-field Gross-Pitaevskii equation,

$$i\hbar \frac{\partial \psi(\vec{r})}{\partial t} = \left[ -\frac{\hbar^2}{2m} \nabla^2 + V(\vec{r}) + g|\psi(\vec{r})|^2 - \Omega_0 L_z \right] \psi(\vec{r}), \quad (1)$$

where  $V(\vec{r}) = \frac{1}{2}m[(\omega_x^2 x^2 + \omega_y^2 y^2 + \omega_z^2 z^2) + \lambda\omega_0^2(x^2 + y^2)^2]$  is the quadratic-plus-quartic potential with the small an-harmonic term  $\lambda$ . Here, we assume  $\omega_x^2 = \omega_0^2(1 + \epsilon)$  and  $\omega_y^2 = \omega_0^2(1 - \epsilon)$ , where  $\epsilon = \frac{(\omega_x^2 - \omega_y^2)}{(\omega_x^2 + \omega_y^2)}$  is a very small deformation parameter due to the rotating deformed potential. Also,  $g = \frac{4\pi a \hbar^2}{m}$  is the strength of the mean-field interaction,  $L_z = xp_y - yp_x$  is the  $z$ -component of the angular momentum operator, and  $\Omega_0$  is the trap rotation frequency. When  $\epsilon = 0$ , it preserves the circular symmetry of the system.

For simplicity, we shall study quasi-2D system. The wave function in the  $z$ -direction is separable and is given by,

$$\psi(z) = \frac{1}{(\sqrt{\pi}a_z)^{1/2}} e^{-\frac{z^2}{2a_z^2}}, \quad (2)$$

where  $a_z^2 = \frac{\hbar}{m\omega_z}$ . After integrating out the  $z$ -component, we get the effective 2D Gross-Pitaevskii energy functional with the modified interaction strength  $g_2 = 2\sqrt{2\pi}\hbar\omega_z a_z$  [16]. One can write down the Lagrangian density corresponding to the quasi-2D system as follows:

$$\begin{aligned} \mathcal{L} = & \frac{i\hbar}{2}(\psi \frac{\partial \psi^*}{\partial t} - \psi^* \frac{\partial \psi}{\partial t}) \\ & + (\frac{\hbar^2}{2m}|\nabla\psi|^2 + V(x, y)|\psi|^2 + \frac{g_2}{2}|\psi|^4 - \Omega\psi^* L_z \psi). \end{aligned} \quad (3)$$

Here, we use the time-dependent variational method [17] to study the properties of a rotating BEC confined in a quadratic-plus-quartic potential. In order to obtain the evolution of the condensate we assume the most general Gaussian wave function,

$$\psi(X, Y, t) = C(t)e^{-\frac{1}{2}[\alpha(t)X^2 + \beta(t)Y^2 - 2i\gamma(t)XY]}, \quad (4)$$

where  $C(t) = \sqrt{\frac{N\sqrt{D}}{\pi\alpha_0^2}}$  is the normalization constant. and  $D = \alpha_1\beta_1 - \gamma_1^2$ . The new co-ordinates  $X$  and  $Y$  are the dimensionless variables,  $X = \frac{x}{a_0}$ ,  $Y = \frac{y}{a_0}$  where  $a_0 = \sqrt{\frac{\hbar}{m\omega_0}}$  is the oscillator length and  $2\omega_0^2 = \omega_x^2 + \omega_y^2$  is the mean frequency. Further,  $\alpha(t) = \alpha_1(t) + i\alpha_2(t)$ ,  $\beta(t) = \beta_1(t) + i\beta_2(t)$  and  $\gamma(t) = \gamma_1(t) + i\gamma_2(t)$  are the time-dependent dimensionless complex variational parameters. The  $\alpha_1$  and  $\beta_1$  are inverse square of the condensate widths in  $x$  and  $y$  directions, respectively.

We obtain the variational Lagrangian  $L$  by substituting Eq. (4) into Eq. (3) and integrating the Lagrangian density over the space co-ordinates,

$$\begin{aligned} \frac{L}{N\hbar\omega_0} = & \frac{1}{4D}[-(\beta_1\dot{\alpha}_2 + \alpha_1\dot{\beta}_2 - 2\gamma_1\dot{\gamma}_2) + (\alpha_1 + \beta_1)D \\ & + (\alpha_2^2 + \gamma_2^2)\beta_1 + (\beta_2^2 + \gamma_2^2)\alpha_1 \\ & - 2(\alpha_2 + \beta_2)\gamma_1\gamma_2 + (\alpha_1 + \beta_1) + \epsilon(\beta_1 - \alpha_1) \\ & + \frac{\lambda}{2} \frac{(3\alpha_1^2 + 2\alpha_1\beta_1 + 3\beta_1^2 + 4\gamma_1^2)}{D} \\ & + PD^{3/2} + 2\Omega(\gamma_1(\beta_2 - \alpha_2) + \gamma_2(\alpha_1 - \beta_1))], \end{aligned} \quad (5)$$

where  $P = 2\sqrt{\frac{2}{\pi}}\frac{(N-1)a}{a_z}$  is the dimensionless parameter that indicates the strength of the mean-field interaction and it can be positive or negative depending on the sign of the  $s$ -wave scattering length  $a$  [18].

The variational energy of the rotating condensate at equilibrium is given in terms of the inverse square width of the condensate along the  $x$  and  $y$  directions and the phase parameter  $\delta\gamma_2$ ,

$$\begin{aligned} \frac{E}{N\hbar\omega_0} = & \frac{1}{4}[(\alpha_1 + \beta_1) + \gamma_2^2(\frac{1}{\alpha_1} + \frac{1}{\beta_1}) + (\frac{1}{\alpha_1} + \frac{1}{\beta_1}) \\ & + \epsilon(\frac{1}{\alpha_1} - \frac{1}{\beta_1}) + \frac{\lambda}{2}(\frac{3}{\alpha_1^2} + \frac{2}{\alpha_1\beta_1} + \frac{3}{\beta_1^2}) \\ & + P\sqrt{\alpha_1\beta_1} + 2\Omega\gamma_2(\frac{1}{\beta_1} - \frac{1}{\alpha_1})], \end{aligned} \quad (6)$$

where  $\Omega = \Omega_0/\omega_0$ . We have used the fact that  $\alpha_2 = \beta_2 = \gamma_1 = 0$ , and also  $\gamma_2 = \delta\gamma_2$  to obtain the above equation.

One can get the equilibrium value of the variational parameters,  $\alpha_{10}$ ,  $\beta_{10}$  and  $\gamma_{20}$  by minimizing the energy with respect to the variational parameters,

$$\alpha_{10}^2(1 + \frac{P}{2}\sqrt{\frac{\beta_{10}}{\alpha_{10}}}) = \gamma_{20}^2 - 2\Omega\gamma_{20} + 1 + \epsilon + \lambda(\frac{3}{\alpha_{10}} + \frac{1}{\beta_{10}}), \quad (7)$$

$$\beta_{10}^2(1 + \frac{P}{2}\sqrt{\frac{\alpha_{10}}{\beta_{10}}}) = \gamma_{20}^2 + 2\Omega\gamma_{20} + 1 - \epsilon + \lambda(\frac{1}{\alpha_{10}} + \frac{3}{\beta_{10}}), \quad (8)$$

and

$$\gamma_{20} = \Omega \frac{(\beta_{10} - \alpha_{10})}{(\beta_{10} + \alpha_{10})}. \quad (9)$$

The Eqs. (7), (8) and (9) describes how the shape of the condensate changes due to the rotation. The relation between the phase parameter and the variational widths is also obtained in [6,19]. The above equation can be re-written as

$$\eta = \frac{\langle x^2 \rangle}{\langle y^2 \rangle} = \frac{\beta_{10}}{\alpha_{10}} = \frac{(\Omega + \gamma_{20})}{(\Omega - \gamma_{20})}. \quad (10)$$

This implies that  $\Omega > \gamma_{20}$  since  $\eta$  is always positive.

The average angular momentum per particle is given by,

$$\frac{\langle L_z \rangle}{N\hbar} = \frac{1}{2}\gamma_{20}(\frac{1}{\beta_{10}} - \frac{1}{\alpha_{10}}) = \frac{\Omega}{2} \frac{(1 - \eta)^2}{\eta(\alpha_{10} + \beta_{10})}. \quad (11)$$

This relation explicitly shows how the angular momentum transfers to a trapped BEC due to the spontaneous shape deformation and consequently how the vortices appears in the rotating BEC by calculating the average angular momentum per particle [20]. We would like to estimate the critical angular frequency from Eq.(11) by using the constraint  $\langle L_z \rangle = N\hbar$ .

### III. LOWEST ENERGY SURFACE MODE FREQUENCY

To calculate the quadrupole mode frequency of BEC confined in a quadratic-plus-quartic potential, we expand the Lagrangian in the following way:  $\alpha = \alpha_{10} + \delta\alpha_1$ ,  $\beta = \beta_{10} + \delta\beta_1$ , and  $\gamma = \delta\gamma_1$ . We keep only the second order deviations from their equilibrium values. Then the Lagrangian quadratic in the deviations takes the following form:

$$\begin{aligned}
L = & \frac{1}{4} \left[ \frac{\delta\alpha_1 \delta\dot{\alpha}_2}{\alpha_{10}^2} + \frac{\delta\beta_1 \delta\dot{\beta}_2}{\beta_{10}^2} + 2 \frac{\delta\gamma_1 \delta\dot{\gamma}_2}{\alpha_{10}\beta_{10}} \right. \\
& + \frac{\delta\alpha_2^2 + \delta\gamma_2^2}{\alpha_{10}} + \frac{\delta\beta_2^2 + \delta\gamma_2^2}{\beta_{10}} \\
& + \frac{\alpha_{10} + \beta_{10}}{\alpha_{10}\beta_{10}} \left( \frac{\delta\alpha_1^2}{\alpha_{10}^2} + \frac{\delta\beta_1^2}{\beta_{10}^2} + \frac{\delta\alpha_1\delta\beta_1}{\alpha_{10}\beta_{10}} + \frac{\delta\gamma_1^2}{\alpha_{10}\beta_{10}} \right) \\
& - \frac{(\delta\alpha_1 + \delta\beta_1)}{\alpha_{10}\beta_{10}} \left( \frac{\delta\alpha_1}{\alpha_{10}} + \frac{\delta\beta_1}{\beta_{10}} \right) \\
& + \frac{\epsilon(\beta_{10} - \alpha_{10})}{\alpha_{10}\beta_{10}} \left( \frac{\delta\alpha_1^2}{\alpha_{10}^2} + \frac{\delta\beta_1^2}{\beta_{10}^2} + \frac{\delta\alpha_1\delta\beta_1}{\alpha_{10}\beta_{10}} + \frac{\delta\gamma_1^2}{\alpha_{10}\beta_{10}} \right) \\
& - \frac{\epsilon(\delta\beta_1 - \delta\alpha_1)}{\alpha_{10}\beta_{10}} \left( \frac{\delta\alpha_1}{\alpha_{10}} + \frac{\delta\beta_1}{\beta_{10}} \right) \\
& + \frac{\lambda}{2(\alpha_{10}\beta_{10})^2} \left( (9\frac{\beta_{10}^2}{\alpha_{10}^2} + 2\frac{\beta_{10}}{\alpha_{10}}) \delta\alpha_1^2 + [(9\frac{\alpha_{10}^2}{\beta_{10}^2} + 2\frac{\alpha_{10}}{\beta_{10}}) \delta\beta_1^2 \right. \\
& + 2\delta\alpha_1\delta\beta_1 + 2(4 + 3\frac{\alpha_{10}}{\beta_{10}} + 3\frac{\beta_{10}}{\alpha_{10}}) \delta\gamma_1^2 \left. \right) \\
& - P\sqrt{\alpha_{10}\beta_{10}} \left( \frac{1}{8} \left( \frac{\delta\alpha_1}{\alpha_{10}} - \frac{\delta\beta_1}{\beta_{10}} \right)^2 + \frac{1}{2} \frac{\delta\gamma_1^2}{\alpha_{10}\beta_{10}} \right) \\
& + \frac{2\Omega}{\alpha_{10}\beta_{10}} ((\delta\beta_2 - \delta\alpha_2)\delta\gamma_1 + (\delta\alpha_1 - \delta\beta_1)\delta\gamma_2). \quad (12)
\end{aligned}$$

For simplicity, first we set  $\Omega = 0$ . Using the Euler-Lagrangian equation of motion, we can get the following two coupled equations of  $\delta\alpha_1$  and  $\delta\beta_1$ .

$$\begin{aligned}
\delta\ddot{\alpha}_1 = & \left[ \frac{P}{2}\alpha_{10}^2\sqrt{\frac{\beta_{10}}{\alpha_{10}}} - 4(1 + \epsilon) - 2\lambda\left(\frac{9}{\alpha_{10}} + \frac{2}{\beta_{10}}\right) \right] \delta\alpha_1 \\
& - \left[ \frac{P}{2}\alpha_{10}^2\sqrt{\frac{\alpha_{10}}{\beta_{10}}} + 2\lambda\frac{\alpha_{10}}{\beta_{10}^2} \right] \delta\beta_1, \quad (13)
\end{aligned}$$

$$\begin{aligned}
\delta\ddot{\beta}_1 = & - \left[ \frac{P}{2}\beta_{10}^2\sqrt{\frac{\beta_{10}}{\alpha_{10}}} + 2\lambda\frac{\beta_{10}}{\alpha_{10}^2} \right] \delta\alpha_1 \\
& + \left[ \frac{P}{2}\beta_{10}^2\sqrt{\frac{\alpha_{10}}{\beta_{10}}} - 4(1 - \epsilon) - 2\lambda\left(\frac{2}{\alpha_{10}} + \frac{9}{\beta_{10}}\right) \right] \delta\beta_1. \quad (14)
\end{aligned}$$

For an isotropic trap,  $\epsilon = 0$ , and we set  $\delta\alpha_1 = -\delta\beta_1$  to calculate the quadrupole mode ( $m_z \pm 2$ ) frequency which is given by,

$$\omega_q^2 = \left( 4 + \frac{20\lambda}{R_0} - PR_0^2 \right), \quad (15)$$

where  $R_0$  is the equilibrium radius of the system, and this can be obtained from the real solution of the cubic equation:  $(1 + \frac{P}{2})R_0^3 - R_0 - 4\lambda = 0$ . The above quadrupole frequency  $\omega_q$  is valid for all interaction (repulsive, attractive) strength with a small anharmonic term ( $\lambda$ ). For  $\lambda = 0$ , the critical value of the interaction strength (above which the system is unstable) is  $P = -2$ . For  $\lambda \neq 0$ , and  $P \leq -2$ , there is no real solution and the system collapses. Note that the quartic potential does not affect the critical strength above which the system is unstable compared to the harmonic trapped BEC. The quadrupole mode frequency  $\omega_q$  vs.  $P$  is shown in the Fig.1.




FIG. 1. Quadrupole mode frequency  $\omega_q$  vs. interaction parameter  $P$  for  $\lambda = 0.005$  (solid) and  $\lambda = 0.05$  (dashed-dot).

For  $\lambda = 0 = P$ , it reproduces the known result for the quadrupole mode frequency of a harmonically trapped non-interacting BEC. Also, for  $\lambda = 0$ , it gives the known result of the quadrupole mode frequency of a harmonically trapped interacting BEC [21]. This result shows that the  $\omega_q$  is decreased compared to the non-interacting case due to the effect of the repulsive interaction. Similarly, the  $\omega_q$  is also increased compared the non-interacting case due to the effect of the attractive interaction. But small anisotropy ( $\lambda \neq 0$ ) increases the  $\omega_q$  drastically in all cases. Note that, the  $\omega_q$  depends on two parameters,  $\lambda$  and  $P$ . For a harmonically trapped BEC,  $\omega_q$  always decreases when  $P$  increases, whereas for the additional quartic term,  $\omega_q$  first decreases for increasing  $P$ , but after some value of  $P$ , it starts increasing with  $P$ . The value of  $P$  at which  $\omega_q$  starts increasing depends on  $\lambda$ . For example,  $P_c = 40$  for  $\lambda = 0.005$  and  $P_c = 10$  for  $\lambda = 0.05$ .

The centrifugal term  $-\Omega L_z$  shifts the quadrupole mode frequency by  $\pm 2\Omega$ . The lowest energy surface mode frequency is given by,

$$\omega_{-2} = \sqrt{4 + \frac{20\lambda}{R_0} - PR_0^2} - 2\Omega. \quad (16)$$

#### IV. ROTATING BEC CONFINED IN A HARMONIC TRAP

Using the three coupled equations (7), (8) and (9) for the three variational parameters, we get the following equation for the ratio of the widths:

$$(1 + \epsilon - \Omega^2)\eta^4 + 2(1 + \epsilon - \Omega^2)\eta^3 + 2\epsilon\eta^2 + 2(\Omega^2 - 1 + \epsilon)\eta + (\Omega^2 - 1 + \epsilon) + \frac{P}{2}\sqrt{\eta}[(1 + \epsilon - \Omega^2)\eta^3 + (1 + 3\epsilon - 5\Omega^2)\eta^2 + (5\Omega^2 + 3\epsilon - 1)\eta + (\Omega^2 - 1 + \epsilon)] = 0. \quad (17)$$

The above relation for the aspect ratio of the system is valid for all interaction strength including repulsive as well as attractive system also. The same kind of relation for the phase parameter  $\gamma_2$  is obtained in [19].

First, we consider a symmetric case ( $\epsilon = 0$ ) and  $P > 0$ . One would expect naively that the ratio of the square of the widths,  $\eta$  becomes one for all rotation frequency. But this is not the case for all rotation frequencies.  $\eta$  is always one when  $0 \leq \Omega \leq \omega_q/2$ , where  $\omega_q$  is the quadrupole mode frequency which is given by

$$\omega_q = \sqrt{4 - PR_0^2}. \quad (18)$$

But  $\eta$  has three solutions when  $\Omega$  is greater than  $\frac{\omega_q}{2}$ . One solution is still one and the other two solutions are less than one and greater than one. But the system has the lowest energy when  $\eta$  is either less than or greater than one. Now we discuss how the angular momentum is transferred to the system by the quadrupolar deformation. When the rotation frequency is in the interval of  $0 \leq \Omega \leq \omega_q/2$ ,  $\eta$  is always one and it is clear from Eq. (11) that the average angular momentum per particle is zero, although we are rotating the system. In other words, the system does not respond to the external rotation till  $\Omega = \frac{\omega_q}{2}$ . This is a signature of the irrotationality of the condensate. Once the system starts deforming, it starts transferring angular momentum (see Eq. 11) and vortex starts nucleate when the average angular momentum per particle is one. The existence of the critical angular frequency at which the angular momentum per particle suddenly goes up from 0 to 1 is the direct manifestation of the superfluid of the condensate. The average angular momentum of each particle has been measured experimentally which is one when a single vortex is nucleated [3]. One can do the same analysis for  $\epsilon \neq 0$ . For large  $P$  and very small  $\epsilon$ ,  $\omega_q \sim \sqrt{2}$ , then a single vortex will form when  $\Omega \sim 1/\sqrt{2} \sim 0.7$  which is experimentally observed.

For non-interacting system, set  $P = 0$  in Eq. (17), then one can get the following quartic equation for  $\eta$ :

$$(1 + \epsilon - \Omega^2)\eta^4 + 2(1 + \epsilon - \Omega^2)\eta^3 + 2\epsilon\eta^2 + 2(\Omega^2 - 1 + \epsilon)\eta + (\Omega^2 - 1 + \epsilon) = 0. \quad (19)$$

In the non-interacting system,  $\eta$  is complex if the rotating frequency  $\Omega$  is in the interval  $[\sqrt{1 - \epsilon}, \sqrt{1 + \epsilon}]$ . Only one real solution exists for  $\Omega < \sqrt{1 - \epsilon}$  and  $\Omega > \sqrt{1 + \epsilon}$  [6,19]. For symmetric trap potential ( $\epsilon = 0$ ) it is clear that there is only one real solution ( $\eta = 1$ ) exist for any rotating frequency  $\Omega$ . The lowest energy surface mode frequency is  $2\omega_0$  and hence instability will occur when  $\Omega = \omega_0$  at which the system is destabilized. There is no spontaneous shape deformation and hence one can not produce a vortex in the non-interacting system because the angular momentum per particle is always zero. We have also checked that for finite but small  $\epsilon$ ,  $\eta$  changes very slowly with  $\Omega$  which is not sufficient to transfer one unit of angular momentum per particle to the system and hence there is no vortex formation in the non-interacting system even for finite  $\epsilon$ .

Now we consider an attractive BEC. We will always consider  $|P| < 2$ , since the condensate will be in stable state if  $|P| < 2$ . In this attractive case, the quadrupole mode frequency is given below.

$$\omega_q = \sqrt{4 + \frac{2|P|}{2 - |P|}}. \quad (20)$$

For any values of  $|P| < 2$ ,  $\omega_q > 2$ . It clearly shows and confirmed by the solution of the Eq. (17) that the shape deformation will not take place in the range of  $\Omega = 0$  to  $\Omega = 1$ . In other words,  $\eta$  is always one. Since there is no shape deformation and consequently vortex nucleation is not possible. Note that authors in Ref. [22,23] showed that the vortex formation in a rotating attractive BEC in harmonic trap is prohibited due to the fact that all the angular momentum is carried by the center of mass motion. Our analysis also shows why there is no vortex formation in an attractive BEC in harmonic trap. One can do the same analysis for  $\epsilon \neq 0$ , but the deformation is not enough to transfer unit quantum of angular momentum to each particle.

#### V. ROTATING BEC CONFINED IN A QUADRATIC-PLUS-QUARTIC POTENTIAL

In this section we will discuss the effect of the quartic potential on the possibility of a vortex formation in a harmonically trapped BEC. Using the three coupled equations (7), (8) and (9), we get the following two coupled polynomial equations:

$$0 = (\alpha_{10}^3\beta_{10}^3 + 2\alpha_{10}^4\beta_{10}^2 + \alpha_{10}^5\beta_{10})(1 + \frac{P}{2}\sqrt{\frac{\beta_{10}}{\alpha_{10}}}) - \alpha_{10}^3\beta_{10}(3\Omega^2 + 1 + \epsilon) + (\alpha_{10}\beta_{10}^3 + 2\alpha_{10}^2\beta_{10}^2)(\Omega^2 - 1 - \epsilon) - \lambda(5\alpha_{10}^2\beta_{10} + 7\alpha_{10}\beta_{10}^2 + \alpha_{10} + 3\beta_{10}^3) \quad (21)$$

and

$$\begin{aligned}
0 = & (\alpha_{10}^3 \beta_{10}^3 + 2\alpha_{10}^2 \beta_{10}^4 + \alpha_{10} \beta_{10}^5) \left(1 + \frac{P}{2} \sqrt{\frac{\alpha_{10}}{\beta_{10}}}\right) \\
& - \alpha_{10} \beta_{10}^3 (3\Omega^2 + 1 + \epsilon) + (\alpha_{10}^3 \beta_{10} + 2\alpha_{10}^2 \beta_{10}^2) (\Omega^2 - 1 - \epsilon) \\
& - \lambda (5\alpha_{10} \beta_{10}^2 + 7\alpha_{10}^2 \beta_{10} + 3\alpha_{10}^3 + \beta_{10}^3). \tag{22}
\end{aligned}$$

To show that there is a circular symmetry breaking of a BEC confined in a quadratic-plus-quartic potential, we put  $\epsilon = 0$  and consider large  $P$  (say  $P = 1000$ ) and solve those two coupled equations. We find that there is a circular symmetry breaking when the rotating frequency is greater than one-half of the quadrupole mode frequency  $\omega_q$ . We plot  $\eta$  vs.  $\Omega$  for  $\epsilon = 0$  and  $\lambda = 0.005$  which is given in Fig.2. In this case,  $\frac{\omega_q}{2} = 0.887179$  and  $\eta$  is always one for  $\Omega = 0$  to  $\Omega = 0.887179$ . When  $\Omega > 0.887179$ ,  $\eta$  has three real and positive solutions and  $\eta < 1$  or  $\eta > 1$  corresponds to the lowest energy state. As the system starts deforming itself, it transfers angular momentum to the system. When  $\Omega = 0.905$ , each particle gets a unit angular momentum, i.e.  $\langle L_z \rangle = N\hbar$ . The vortex formation can occur for  $\Omega < 1$  as well as  $\Omega > 1$ , depending on the parameter  $P$  and  $\lambda$ . For  $\lambda = 0.05$ , the critical rotation frequencies are  $\Omega_c = 1.2, 1.422, 1.554$  for  $P = 100, 500, 1000$ , respectively.




FIG. 2. The variation of  $\eta$  vs.  $\Omega$  for  $\lambda = 0.005$  and large  $P$ .

The shape deformation also occurs for non-interacting and attractive Bose gases. To discuss about the shape deformation in non-interacting system, first we set  $P = 0$  in the equations (21) and (22). We find that the  $\eta$  has three real and positive solutions when  $\Omega > \frac{\omega_q}{2}$ , like in the condensate in the harmonic trap. For quantitative discussion, we take  $\lambda = 0.005$ , then the system starts deformation when  $\Omega > \frac{\omega_q}{2} = 1.0123$ . In the previous section we have seen that the spontaneous deformation takes place due to the two-body interaction. But here spontaneous shape deformation exists in an ideal Bose system confined in a quadratic-plus-quartic potential. The average angular momentum per particle is always zero

when  $\Omega = 0$  to  $\Omega = 1.0123$ , but  $\langle L_z \rangle = N\hbar$  at  $\Omega = 1.027$ . We give another example for  $\lambda = 0.05$ . For this anharmonic strength, the system will start deforming when  $\Omega > \frac{\omega_q}{2} = 1.10895$  and the vortex will form when  $\langle L_z \rangle = N\hbar$  at  $\Omega = 1.21$ . It shows that as we increase the strength of the anharmonic term, the critical rotational frequency is also increasing. Note that the vortex formation occurs only when  $\Omega > 1$  because  $\frac{\omega_q}{2}$  is always greater than 1. The variation of  $\eta$  vs.  $\Omega$  for non-interacting BEC is shown in the Fig.3.




FIG. 3. The variation of  $\eta$  vs.  $\Omega$  for  $P = 0$ ,  $\lambda = 0.005$  and  $\epsilon = 0$ .

It is worth to point out that the shape deformation can even occur in the non-interacting Bose gas if it is confined in a quadratic-plus-quartic potential.

In the previous section, we have seen that harmonically trapped attractive BEC can not hold a vortex. This is because of the absence of the shape deformation in the rotating system. Now we will see what happens if we add the quartic potential in Hamiltonian. Once we add the quartic potential, we can access a large rotation frequency ( $\Omega \geq 1$ ) for which the rotating attractive system is not destabilized. We assume  $P = -0.015$  and  $\lambda = 0.01$  and then we get  $\frac{\omega_q}{2} = 1.02606$  for the symmetric trap. By solving the equations (21) and (22), we see that the spontaneous shape deformation starts when  $\Omega \sim 1.02606$  and transferring an angular momentum to the system. Each particle gets an unit angular momentum,  $\langle L_z \rangle = N\hbar$ , when  $\Omega = 1.052$ . We give another example. Assume,  $P = -1.5$  and  $\lambda = 0.05$  and then we get  $\frac{\omega_q}{2} = 1.66215$  for the symmetric trap. By solving the equations (21) and (22), we get three real and positive solution of  $\eta$  when  $\Omega > \frac{\omega_q}{2}$ . But the existence of the three solutions of  $\eta$  does not mean that the system will starts deforming itself. Actually, the lowest energy state can be obtained only for  $\eta = 1$ . It means there is no shape deformation for these parameters and hence a vortex can not appear under this conditions. In this attractive sys-

tem, the shape deformation can occur which leads to the vortex nucleation, depending on the strengths of the two-body interaction and the quartic potential. When there is no shape deformation for a given large  $P$  and  $\lambda$ , the center of mass motion is excited [22,23]. These results are also consistent with the recent results of Lundh *et al.* [14]. Note that  $\frac{\omega_a}{2}$  is always greater than 1, so that deformation takes place only when  $\Omega > 1$ .

## VI. SUMMARY AND CONCLUSIONS

In this paper we have calculated the quadrupole mode frequency. We have shown that when the rotating frequency of an interacting BEC confined in the quadratic-plus-quartic potential is greater than one-half of the quadrupole mode frequency, the system itself starts deforming from a circular shape to an elliptic shape and hence it breaks the rotational symmetry of the Hamiltonian. We have shown how this self-deformation transfer an angular momentum to the rotating system and consequently creates a vortex.

We have argued that the formation of the vortex is not possible due to the absence of the spontaneous shape deformation of a harmonically trapped rotating attractive BEC as well as an ideal BEC. But the shape deformation and consequently the vortex formation can occur in a rotating attractive BEC if it is confined by an additional quartic potential. We have also shown that the shape deformation occurs not only due to the two-body repulsive interaction, it can occur even in the non-interacting as well as attractive BEC if it is confined in a quadratic-plus-quartic potential and consequently it produces a vortex. However, the shape deformation and consequently forming a quantized vortex in a rotating attractive BEC depends on the strengths of the two-body interaction and the quartic potential.

Note that, in actual experiments the lowest energy surface mode is excited through the rotating potential with small anisotropies. In this work we have studied (for simplicity) the vortex formation in a symmetrically trapped BEC, but we can easily extend our work with small anisotropy  $\epsilon$ .

- 
- [1] F. Dalfovo, S. Giorgini, L. Pitaevskii and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999). A. J. Leggett, Rev. Mod. Phys. **71**, 463 (1999).
  - [2] K. W. Madison, F. Chevy, W. Wohlleben, and J. Dalibard, Phys. Rev. Lett. **84**, 806 (2000).
  - [3] F. Chevy, K. W. Madison, and J. Dalibard, Phys. Rev. Lett. **85**, 2223 (2000).
  - [4] J. R. Abo-Shaer, C. Raman, J. M. Vogels, and W. Ketterle, Science, **292**, 476 (2001); C. Raman, J. R. Abo-Shaer, J. M. Vogels, K. Xu, and W. Ketterle, Phys. Rev. Lett. **87**, 210402 (2001).
  - [5] G. Baym and C. J. Pethick, Phys. Rev. Lett. **76**, 6 (1996); F. Dalfovo and S. Stringari, Phys. Rev. A **53**, 2477 (1996); S. Sinha, Phys. Rev. A **55**, 4325 (1997).
  - [6] A. Recati, F. Zambelli, and S. Stringari, Phys. Rev. Lett. **86**, 377 (2001).
  - [7] M. Kramer, L. Pitaevskii, S. Stringari and F. Zambelli, Laser Physics **12**(1), 113 (2002).
  - [8] K. W. Madison, F. Chevy, V. Bretin, and J. Dalibard, Phys. Rev. Lett. **86**, 4443 (2001).
  - [9] S. Sinha and Y. Castin, Phys. Rev. Lett. **87**, 190402 (2001).
  - [10] A. L. Fetter, Phys. Rev. A, **64**, 063608 (2001).
  - [11] E. Lundh, Phys. Rev. A **65**, 043604 (2002).
  - [12] K. Kasamatsu, M. Tsubota, and M. Ueda, Phys. Rev. A **66**, 053606 (2002).
  - [13] G. Kavoulakis and G. Baym, New Journal of Physics **5**, 51.1 (2003).
  - [14] E. Lundh, A. Collin, K.A. Suominen, Phys. Rev. Lett. **92**, 070401 (2004).
  - [15] V. Bretin, S. Stock, Y. Seurin, and J. Dalibard, Phys. Rev. Lett. **92**, 050403 (2004).
  - [16] T. L. Ho and M. Ma, J. Low Temp. Phys. **115**, 61 (1999).
  - [17] V. Perez-Garcia *et al.* Phys. Rev. Lett. **77**, 5320 (1996).
  - [18] S. Inouye *et al.*, Nature (London), **392**, 151 (1998); J. Stenger *et al.*, Phys. Rev. Lett. **82**, 2422 (1999).
  - [19] P. Rosenbusch *et al.* Phys. Rev. Lett. **88**, 250403 (2002).
  - [20] Juan J. Garcia-Ripoll and Victor M. Perez-Garcia, Phys. Rev. A. **64**, 013602 (2001); F. Dalfovo and S. Stringari, Phys. Rev. A **63**, 011601(R) (2001).
  - [21] S. Stringari, Phys. Rev. Lett. **77**, 2360 (1996).
  - [22] N. K. Wilkin, J. M. F. Gunn, and R. A. Smith, Phys. Rev. Lett. **80**, 2265 (1998).
  - [23] B. Mottelson, Phys. Rev. Lett. **83**, 2695 (1999).