

Nanoscience with Attosecond Laser Pulses

Janina Marciak-Kozłowska^a and Mirosław Kozłowski^b

^aInstitute of Electron Technology, Al. Lotników 32/46 02-668 Warsaw
Poland

^bPhysics Department, Science Teachers College and Institute of Experimental Physics,
Warsaw University Hoża 69, 00-681 Warsaw Poland,
e-mail: mirkoz@fuw.edu.pl

Abstract

In this paper the interaction of attosecond laser pulses with matter is investigated. The scattering and potential motion of heat carriers as well as the external force are considered. Depending on the ratio of the scatterings and potential motion the heat transport is described by the thermally forced Klein-Gordon or thermally modified telegraph equation. For thermal Heisenberg type relation $V = \hbar / \tau$ (τ is the relaxation time and V is the potential) the heat transport is described by the thermally distortionless damped wave equation. In this paper Klein-Gordon, modified telegraph¹ equation and wave equation are solved.

Key words: Attosecond laser pulses; Quantum heat transport equation; Klein-Gordon thermal equation.

¹Considering history of the formulation and application of the hyperbolic partial differential equation, Oliver Heaviside for the first time wrote (in 1876) hyperbolic equation for voltage $v(x;t)$ in submarine cable. [G. Thomas and D. J. Raine, Physics to a degree, Gordon and Breach, 2000, p. 43]. $\frac{\partial^2 v(x;t)}{\partial x^2} = RC \frac{\partial v(x;t)}{\partial t} + LC \frac{\partial^2 v}{\partial t^2}$. We propose the name Heaviside equation instead of the telegraph equation for the hyperbolic partial equation (thermal wave equation with both first and second derivative).

1 Introduction

With attosecond lasers ($1 \text{ as} = 10^{-18} \text{ s}$) physicist and engineers are now close to controlling the motion of electrons on a timescale that is substantially shorter than the oscillation period of visible light. It is also possible with attosecond laser pulse to rip an electron wave-packet from the core of an atom and set it free with similar temporal precision.

Since we know that there is an absolute velocity, velocity of light it is sufficient to define the smallest time interval (e.g. 1 as) and the unit of length is a derivative quantity ($3 \cdot 10^{10} \text{ m} = 0.3 \text{ nm} = \text{Bohr radius}$).

After the standards of time and space are defined the laws of classical physics relating such parameters as distance, time, velocity, temperature are assumed to be independent of accuracy with which these parameters can be measured. It should be noted that this assumption does not enter explicitly into the formulation of the classical physics. This means that together with the assumption of the existence of the objective reality independent of any measurements (in classical physics) it was tacitly assumed that there is a possibility of unlimited increase of measurements accuracy. Bearing in mind the atomicity of time i.e. considering the smallest time period, Planck time the above statement is obviously not true. With attosecond laser pulses we are at the limit of laser time resolution.

With attosecond laser pulses we enter new Nano World where size become comparable to atomic dimension, where transport phenomena follows different laws than in macro world. This means the end of first step miniaturization (from 10^{-3} m to 10^{-6} m) and the beginning new one (from 10^{-6} to 10^{-9}). Nano World is a quantum world with all predictable and nonpredictable (yet) features.

In this paper we develop and solve the quantum relativistic heat transport equation for Nano World transport phenomena when external forces exist. This is the generalization of the results of paper [1] in which quantum relativistic hyperbolic equation was proposed and solved.

2 Relativistic hyperbolic heat transport equation

In paper [1] relativistic hyperbolic transport equation (RHT) was formulated:

$$\frac{1}{v^2} \frac{\partial^2 T}{\partial t^2} + \frac{m_0}{h} \frac{\partial T}{\partial t} = r^2 T : \quad (1)$$

In equation (1) v is the velocity of heat waves, m_0 is the mass of heat carrier and γ the Lorentz factor $\gamma = (1 - \frac{v^2}{c^2})^{-1/2}$. As was shown in paper [1] the heat energy (heaton temperature) T_h can be defined as follows:

$$T_h = m_0 v^2 : \quad (2)$$

Considering that v , thermal wave velocity equals [1]

$$v = \frac{c}{\beta} ; \quad (3)$$

where β is the coupling constant for the interactions which generate the thermal wave ($\beta = 1/137$ and $\beta = 0.15$ for electromagnetic and strong force respectively) heaton temperature equals

$$T_h = \frac{m_0 v^2}{\beta} : \quad (4)$$

From formula (4) one concludes that heaton temperature is the linear function of the mass m_0 of the heat carrier. It is quite interesting to observe that the proportionality of T_h and heat carrier mass m_0 was the first time observed in ultrahigh energy heavy ion reactions measured at CERN [2]. In paper [2] it was shown that temperature of pions, kaons and protons produced in Pb+Pb, S+S reactions are proportional to the mass of particles. Recently at Rutherford Appleton Laboratory (RAL) the VULCAN laser was used to produce the elementary particles: electrons and pions [3].

In the present paper the forced relativistic heat transport equation will be studied and solved. In paper [4] the damped thermal wave equation was developed:

$$\frac{1}{v^2} \frac{\partial^2 T}{\partial t^2} + \frac{m}{h} \frac{\partial T}{\partial t} + \frac{2V m}{h^2} T - r^2 T = 0 : \quad (5)$$

The relativistic generalization of equation (5) is quite obvious:

$$\frac{1}{v^2} \frac{\partial^2 T}{\partial t^2} + \frac{m_0}{h} \frac{\partial T}{\partial t} + \frac{2V m_0}{h^2} T - r^2 T = 0 : \quad (6)$$

It is worthwhile to note that in order to obtain nonrelativistic equation we put $\gamma = 1$.

If the nucleus contains Z protons than we have for electron temperature in atom

$$T_h^Z(\text{atom}) = m_e (Z \alpha)^2 c^2 : \quad (7)$$

Equation (7) also tell us the magnitude of the temperature of the ionization of the atom :

$$T_h^Z(\text{ionization}) = \frac{1}{2} m_e c^2 (Z \alpha)^2 \left(1 + \frac{m_e}{m_p} \right)^{-1} \approx Z^2 10^5 \text{K} :$$

The motion of electron in the atom is equivalent to the flow of an electric current in a loop of wire. With attosecond laser pulses we will be able to influence the current in the atomic "wire". This opens quite new perspective for the Nano World technology or nanotechnology. The new equation (6) is the natural candidate for the master equation which can be used in nanotechnology.

When external force is present $F(x;t)$ the forced damped heat transport is obtained instead of equation (6) (in one dimensional case):

$$\frac{1}{v^2} \frac{\partial^2 T}{\partial t^2} + \frac{m_0}{h} \frac{\partial T}{\partial t} + \frac{2V m_0}{h^2} T - \frac{\partial^2 T}{\partial x^2} = F(x;t) : \quad (8)$$

The hyperbolic relativistic quantum heat transport equation (RQHT), formula (8), describes the forced motion of heat carriers which undergo the scatterings ($\frac{m_0}{h} \frac{\partial T}{\partial t}$ term) and are influenced by potential ($\frac{2V m_0}{h^2} T$ term).

The solution of equation can be written as

$$T(x;t) = e^{-\frac{t}{\tau}} u(x;t); \quad (9)$$

where $\tau = h/(m v^2)$ is the relaxation time. After substituting formula (9) to the equation (8) we obtain new equation

$$\frac{1}{v^2} \frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} + q u(x;t) = e^{\frac{t}{\tau}} F(x;t); \quad (10)$$

and

$$q = \frac{2V m_0}{h^2} - \frac{m v^2}{2h} ; \quad (11)$$

Equation (10) can be written as:

$$\frac{\partial^2 u}{\partial t^2} - v^2 \frac{\partial^2 u}{\partial x^2} + qv^2 u(x;t) = G(x;t); \quad (12)$$

where

$$G(x;t) = v^2 e^{\frac{t}{\tau}} F(x;t):$$

When $q > 0$ equation (12) is the forced Klein-Gordon (K-G) equation. The solution of the forced Klein-Gordon equation for the initial conditions:

$$u(x;0) = f(x) \quad u_t(x;0) = g(x) \quad (13)$$

has the form : [4]

$$\begin{aligned} u(x;t) = & \frac{f(x-vt) + f(x+vt)}{2} \\ & + \frac{1}{2v} \int_{x-vt}^{x+vt} g(\xi) J_0 \left(q \sqrt{v^2 t^2 - (x-\xi)^2} \right) d\xi \\ & - \frac{p}{2qvt} \int_{x-vt}^{x+vt} f(\xi) \frac{J_1 \left(q \sqrt{v^2 t^2 - (x-\xi)^2} \right)}{\sqrt{v^2 t^2 - (x-\xi)^2}} d\xi \\ & + \frac{1}{2v} \int_0^{t-x/v} G(x-v(t-t^0), t-t^0) J_0 \left(q \sqrt{v^2 (t-t^0)^2 - (x-\xi)^2} \right) dt^0 d\xi : \end{aligned} \quad (14)$$

When $q < 0$ equation (12) is the forced modified Heaviside (telegraph) equation with the solution: [4]

$$\begin{aligned} u(x;t) = & \frac{f(x-vt) + f(x+vt)}{2} \\ & + \frac{1}{2v} \int_{x-vt}^{x+vt} g(\xi) I_0 \left(q \sqrt{v^2 t^2 - (x-\xi)^2} \right) d\xi \\ & + \frac{p}{2v} \int_{x-vt}^{x+vt} f(\xi) \frac{I_1 \left(q \sqrt{v^2 t^2 - (x-\xi)^2} \right)}{\sqrt{v^2 t^2 - (x-\xi)^2}} d\xi \\ & + \frac{1}{2v} \int_0^{t-x/v} G(x-v(t-t^0), t-t^0) I_0 \left(q \sqrt{v^2 (t-t^0)^2 - (x-\xi)^2} \right) d\xi dt^0 : \end{aligned} \quad (15)$$

When $q = 0$ equation (12) is the forced thermalequation [4].

$$\frac{\partial^2 u}{\partial t^2} - v^2 \frac{\partial^2 u}{\partial x^2} = G(x; t); \quad (16)$$

On the other hand one can say that equation (16) is the distortionless hyperbolic equation. The condition $q = 0$ can be rewrite as:

$$V = \frac{h}{8}; \quad (17)$$

The equation (17) is the analogous to the Heisenberg uncertainty relations. Considering formula (2) equation (17) can be written as:

$$V = \frac{T_h}{8}; \quad V < T_h; \quad (18)$$

One can say that the distortionless waves can be generated only if $T_h > V$. For $T_h < V$, i.e. when the "Heisenberg rule" is broken, the shape of the thermalwaves is changed.

We consider the initial and boundary value problem for the inhomogeneous thermalwave equation in semi-infinite interval [4]: that is

$$\frac{\partial^2 u}{\partial t^2} - v^2 \frac{\partial^2 u}{\partial x^2} = G(x; t); \quad 0 < x < 1; \quad t > 0; \quad (19)$$

with initial condition:

$$u(x; 0) = f(x); \quad \frac{\partial u(x; 0)}{\partial t} = g(x); \quad 0 < x < 1;$$

and boundary condition

$$au(0; t) - b \frac{\partial u(0; t)}{\partial x} = B(t); \quad t > 0; \quad (20)$$

where $a \geq 0; b \geq 0; a + b > 0$ (with a and b both equal to constants) and $F; f; g$ and B are given functions. The solution of equation (19) is of the form [4]:

$$\begin{aligned} u(x; t) = & \frac{1}{2} [f(x - vt) + f(x + vt)] + \frac{1}{2v} \int_{x-vt}^{x+vt} g(s) ds \\ & + \frac{1}{2v} \int_0^t \int_{x-v(t-t^0)}^{x+v(t-t^0)} F(s; t^0) ds dt^0; \end{aligned} \quad (21)$$

In the special case where $f = g = F = 0$ we obtain the following solution of the initial and boundary value problem (19, 20):

$$u(x;t) = \begin{cases} 0; & x > vt; \\ \frac{v}{b} \int_0^{t - \frac{x}{v}} \exp\left(\frac{va}{b} y\right) B\left(t + \frac{x}{v} - y\right) dy; & 0 < x < vt; \end{cases} \quad (22)$$

if $b \neq 0$. If $a = 0$ and $b = 1$, we have:

$$u(x;t) = \begin{cases} 0; & x > vt; \\ v \int_0^{t - \frac{x}{v}} B(y) dy; & 0 < x < vt; \end{cases}$$

whereas if $b = 0$ and $a = 1$, we obtain:

$$u(x;t) = \begin{cases} 0; & x > vt; \\ B\left(t - \frac{x}{v}\right); & 0 < x < vt; \end{cases}$$

It can be concluded that the boundary condition (20) gives rise to a wave of the form $K\left(t - \frac{x}{v}\right)$ that travels to the right with speed v . For this reason the foregoing problem is often referred to as a signaling problem for the thermal waves.

3 Looking into Nano World

With attosecond laser pulses we expect to be able to track the motion of an electron within a femtosecond. We could also follow the relaxation of the remaining bound electrons to their new equilibrium states. Once we can generate attosecond pulses with much higher energies we will be able to use them as both pumps and probes in pump-probe spectroscopy on the atomic level.

Up to now the pump-probe method allows the investigation of the heat transport on the bulk scale. The experiments triggered by Borson paper [6] disclosed the finite velocity of the heat transport i.e. the hyperbolicity of heat transport equation.

Accordingly with attosecond pump-probe method we will be able the study of heat transport in the Nano World. As was shown in [1] one can define

the electron temperature (electron temperature) in ground state hydrogen like atom :

$$T_h(\text{atom}) = m_e^2 c^2; \quad (23)$$

and in hydrogen like molecule [7, 8]

$$T_h(\text{molecule}) = \frac{2m_e}{m_p} m_e c^2; \quad (24)$$

4 Conclusion

In this paper damped forced therm al wave equation was proposed and solved. For Cauchy boundary initial condition the solution of the equation was presented. It was shown that the new equation, depending on the relation between scattering and potential motion, is of the form of the forced therm al Klein-Gordon equation or the forced modified Heaviside (telegraph) equation. For the therm al Heisenberg type relation $V = \hbar c$ the new equation is the distortionless forced wave equation. In the context of the forced wave equation the signaling solution was obtained.

R e f e r e n c e s

- [1] J.M arciak-K ozłowska, M .K ozłowski,
Lasers in Engineering, 11, (2001), p.259.
- [2] I.G .Bearden et al.,
Phys. Rev. Lett. 78, (1997), p.2080.
- [3] K.W .D .Ledingham and P.A .Norreys,
Contemporary Physics, 40, (1999), p.367.
- [4] M .K ozłowski, J.M arciak-K ozłowska,
Lasers in Engineering, 8, (1998,) p.11.
- [5] E .Zauderer,
Partial D i fferential Equation of Applied M athem atics, Second Edition,
W iley 1989.
- [6] S.D .Brorson et al.,
Phys. Rev. Lett., 59, (1987), p.1962.
- [7] M .K ozłowski, J.M arciak-K ozłowska,
Lasers in Engineering, 9, (1999), 103.
- [8] M .K ozłowski, J.M arciak-K ozłowska,
From Q uarks to Bulk M atter, H adronic P ress, U SA (2001).