

The Last Passage Problem on Graphs

Jean Desbois¹ and Olivier Benichou²

14th April 2024

¹ Laboratoire de Physique Theorique et Modeles Statistiques. Universite Paris-Sud, Bât. 100, F-91405 Orsay Cedex, France.

² Laboratoire de Physique de la Matiere Condensee, College de France, 11, place M. Berthelot, 75231 Paris Cedex 05, France.

Abstract

We consider a brownian motion on a general graph, that starts at time $t = 0$ from some vertex O and stops at time t somewhere on the graph. Denoting by g the last time when O is reached, we establish a simple expression for the Laplace Transform, L , of the probability density of g . We discuss this result for some special graphs like star, ring, tree or square lattice. Finally, we show that L can also be expressed in terms of primitive orbits when, for any vertex, all the exit probabilities are equal.

Around the years 1930, P. Levy [1] got several arc-sine laws concerning the 1D Brownian Motion on an infinite line. Let us consider such a process starting at $t = 0$ from the origin O and stopping at time t and denote by g the last time when O is reached. The second arc-sine law discovered by Levy concerns the probability law of g . It can be stated as follows:

$$P(g < u) = \frac{2}{\pi} \arcsin \sqrt{\frac{u}{t}} \quad (1)$$

$$\text{with the probability density} \quad P_t(u) \quad \frac{dP(g < u)}{du} = \frac{1}{\pi} \frac{1}{u(t-u)} \quad (2)$$

In particular, the Laplace transform of $P_t(u)$ is written:

$$\int_0^{\infty} dt e^{-\lambda t} \int_0^t du P_t(u) e^{-\lambda u} = \frac{1}{(\lambda + \dots)} \quad (3)$$

Levy also proved that the same law occurs when g represents the time spent in the region $(x > 0)$ (first arc-sine law).

The infinite line with only one point, O , specified, can be viewed as a kind of very simple graph consisting in one vertex, O , and two semi-infinite lines originating from O .

Our goal in this Letter is to get an analogue of the second arc-sine law (3) but for a quite general graph.

Let us recall that graphs properties have interested for many years physicists [2] as well as mathematicians [3]. For instance, the knowledge of spectral properties of the Laplacian operator is useful to understand superconducting networks [4] as well as organic molecules [5]. We also know that spectral determinants play a central role in the computation of weak localization corrections [6]. Recently, graphs have also been used in the context of nonequilibrium statistical physics (see [7] and references therein) and also in the field of quantum chaos [8].

The study of Brownian motion on graphs is also an active area of research. For instance, occupation times [9, 10] or local time [11] distributions have been recently computed for general graphs.

So, let us begin by setting the notations that will be used throughout this Letter. We consider a graph made of V vertices, numbered from 0 to $V-1$, linked by B bonds of finite lengths. The coordination number of vertex i is m_i ($\sum_{i=0}^{V-1} m_i = 2B$).

On each bond $[i, j]$, of length l_{ij} , we define the coordinate x that runs from 0 (vertex i) to 1 (vertex j). Sometimes we will also use the coordinate $x = 1 - x$. The set of coordinates for graph is simply noted \mathcal{X} .

An arc (i, j) is defined as the oriented bond from i to j . Each bond $[i, j]$ is therefore associated with two arcs (i, j) and (j, i) . In the sequel, we will consider the following ordering of the $2B$ arcs: $(0_1)(0_2) \dots (0_{m_0}) \dots (1_1) \dots (1_{m_1}) \dots (m-1_1) \dots (m-1_{m_{m-1}}) \dots$

Now, let us consider a brownian motion starting at some vertex O (label 0) of the graph at time $t=0$ and stopped at time t . Let g be the last time when O is visited. The probability, $P_t(g < u)$ can be written:

$$P_t(g < u) = \int_{\text{Graph}} dx^0 dx G_0(0; 0; x; u) G_D(x; u; x^0; t) \quad (4)$$

where G_0 is the free propagator on the graph and G_D is the propagator with Dirichlet boundary conditions in O . (4) simply expresses the fact that the brownian particle will not hit O between the times u and t . Let us define the probability density $P_t(u) = \frac{dP(g < u)}{du}$. In the sequel, we will be especially interested in the Laplace transform:

$$L = \int_0^{\infty} dt e^{-t} \int_0^{\infty} du P_t(g < u) e^{-u} \quad (5)$$

$$L = \int_0^{\infty} dt e^{-t} \int_0^{\infty} du P_t(u) e^{-u} = \frac{1}{1 + L} + L \quad (6)$$

With (4), we see that:

$$L = \int_{\text{Graph}} dx^0 dx \int_0^{\infty} \frac{1}{H + x} x \int_0^{\infty} \frac{1}{H_D + x^0} x^0 \int_{\text{Graph}} dx^0 dx G_0(0; x; x^0 +) G_D(x; x^0;) \quad (7)$$

On the bond $[i, j]$, H and H_D are equal to $\frac{1}{2} \frac{d^2}{dx^2}$ and we define:

$$G_0^{(i)} = \lim_{x \rightarrow 0} G_0(0; x; x^0 +) \quad ; \quad G_0^{(j)} = \lim_{x \rightarrow 0} \frac{dG_0(0; x; x^0 +)}{dx} \quad (8)$$

$$G_D^{(i)}(x) = \lim_{x^0 \rightarrow 0} G_D(x; x^0;) \quad ; \quad G_D^{(j)}(x) = \lim_{x^0 \rightarrow 0} \frac{dG_D(x; x^0;)}{dx^0} \quad (9)$$

The integration over x^0 in (7) is easily performed with the help of the equation

$$\frac{1}{2} \frac{d^2}{dx^2} + \sum_{i=1}^m G_D(x; x^0; \epsilon_i) = (x - x^0) \quad (10)$$

We get

$$\sum_{\text{Graph}} dx^0 G_D(x; x^0; \epsilon_i) = \frac{1}{2} \sum_{i=1}^m G_D^{(0, \epsilon_i)}(x) \quad (11)$$

To go further, we have to specify the behaviours of the resolvents G_0 and G_D in the neighbourhood of all the vertices.

Let us consider some vertex with its nearest neighbours $\epsilon_i, i = 1; 2; \dots; m$, on the graph. Suppose that the brownian particle reaches ϵ . It will come out towards ϵ_i with some exit probability p_{ϵ_i} . This implies the following equations to be satisfied by G_0 :

$$\frac{G_0^{(\epsilon_1)}}{p_{\epsilon_1}} = \frac{G_0^{(\epsilon_2)}}{p_{\epsilon_2}} = \dots = \frac{G_0^{(\epsilon_m)}}{p_{\epsilon_m}} \quad \text{if } \epsilon \neq 0 \quad (12)$$

Remark that the resolvent will be continuous in vertex ϵ only when the particle exits from with the same probability in all the directions.

Moreover, current conservation implies:

$$\sum_{i=1}^m G_0^{(\epsilon_i)} = 0 \quad ; \text{ if } \epsilon = 0 \quad (13)$$

$$\sum_{i=1}^m G_0^{(0, \epsilon_i)} = 2 \quad (14)$$

On the link $[\epsilon, \epsilon']$, $G_0(0; x; \epsilon; \epsilon')$ must satisfy:

$$\frac{1}{2} \frac{d^2}{dx^2} + \sum_{i=1}^m G_0(0; x; \epsilon; \epsilon') = 0 \quad (15)$$

with the solution:

$$G_0(0; x; \epsilon; \epsilon') = G_0^{(\epsilon)} \frac{\sinh \frac{p}{2(\epsilon' + \epsilon)} (1 - x)}{\sinh \frac{p}{2(\epsilon' + \epsilon)} 1} + G_0^{(\epsilon')} \frac{\sinh \frac{p}{2(\epsilon' + \epsilon)} x}{\sinh \frac{p}{2(\epsilon' + \epsilon)} 1} \quad (16)$$

So, we deduce:

$$G_0^{(\epsilon)} = \frac{q}{2(\epsilon' + \epsilon)} c(\epsilon' + \epsilon) G_0^{(\epsilon)} + s(\epsilon' + \epsilon) G_0^{(\epsilon')} \quad (17)$$

with:

$$c(\epsilon' + \epsilon) = \coth \frac{q}{2(\epsilon' + \epsilon)} 1 = c(\epsilon' + \epsilon) \quad (18)$$

$$s(\epsilon' + \epsilon) = \frac{1}{\sinh \frac{p}{2(\epsilon' + \epsilon)} 1} = s(\epsilon' + \epsilon) \quad (19)$$

Those considerations allow to write eqs.(12, 13, 14) in a matrix form :

$$M(\phi) f = \frac{1}{2(\phi + 1)} l \quad (20)$$

where f and l are two $(V-1)$ vectors. The components of f are the quantities f_i and for the components of l we have $l_j = 2 \phi_j$.

$M(\phi)$ is a $(V-1) \times (V-1)$ vertex matrix with the non vanishing elements (i runs from 0 to $V-1$):

$$M(\phi)_{ij} = \sum_{i=1}^X p_i c_i(\phi) \quad (21)$$

$$M(\phi)_{ij} = p_{ij} s_{ij}(\phi) \quad \text{if } [ij] \text{ is a bond} \quad (22)$$

$$= 0 \quad \text{otherwise} \quad (23)$$

Similar considerations hold for $G_D(x; x^0; \phi)$ and we get:

$$\frac{G_D^{(i-1)}(x)}{p_{i-1}} = \dots = \frac{G_D^{(m_0)}(x)}{p_{m_0}} \quad F(x) \quad ; \quad \text{if } \phi \neq 0 \quad (24)$$

$$G_D^{(0 \dots i)}(x) = 0 \quad ; \quad i = 1; \dots; m_0 \quad (25)$$

$$\sum_{i=1}^X G_D^{(0 \dots i)}(x) = 0 \quad ; \quad \text{if } \phi \neq 0 \quad (26)$$

$$\text{and the matrix equation} \quad M_D(\phi) F(x) = \frac{1}{\phi} l_D(x) \quad (27)$$

M_D is the matrix M with the first line and first column deleted. $F(x)$ and $l_D(x)$ are $(V-1) \times 1$ vectors. The components of $F(x)$ are the quantities $F_i(x)$, $i = 1; \dots; V-1$. The nonvanishing components of $l_D(x)$ are defined as follows.

If $x \in [ab]$ with $a \neq 0$ and $b \neq 0$ then

$$(l_D(x_{ab}))_a = s_{ab}(\phi) \sinh\left(\frac{\phi}{2} x_{ba}\right) \quad (28)$$

$$(l_D(x_{ab}))_b = s_{ab}(\phi) \sinh\left(\frac{\phi}{2} x_{ab}\right) \quad (29)$$

If $x \in [0_j]$ then

$$l_D(x_{0_j}) = s_{0_j}(\phi) \sinh\left(\frac{\phi}{2} x_{0_j}\right) \quad (30)$$

Performing the integration over x in (7), we finally get the following simple expression for L :

$$L = \frac{1}{2(\phi + 1)} \frac{M^{-1}(\phi)_{00}}{(M^{-1}(\phi))_{00}} \quad (31)$$

This is the central result of this paper.

Remark that, for the special case when G_0 and G_D are continuous Green's functions, we can recast (31) in the form ¹

$$L = \frac{1}{G_0(0;0;)} \frac{G_0(0;0; +)}{G_0(0;0;)} \quad (32)$$

To go further and deduce from (31) some general properties of the density $P_t(u)$, let us consider the following identity

$$\int_0^{\infty} dt e^{-pt} \int_0^t du e^{-u} r(u) s(t-u) = R(p) S(p) \quad (33)$$

where R and S are the Laplace Transforms of r and s .

Setting $R(p) = \frac{M^{-1}(p)_{00}}{p}$ and $S(p) = \frac{1}{p(M^{-1}(p))_{00}}$ in (33), we realize from (31) and (6) that the probability density factorizes in the following way:

$$P_t(u) = r(u) s(t-u) \quad (34)$$

This result holds for any graph.

It is now easy to analyze the limiting behaviours of $P_t(u)$ when $u \rightarrow 0^+$ and when $u \rightarrow t$.

In the first case, we can write $P_t(u) = r(u) s(t-u)$. Moreover, for large p values, $M(p) \rightarrow 1$ and $R(p) \rightarrow \frac{1}{p}$. So

$$P_t(u) \rightarrow \frac{1}{p} s(t) \quad ; \quad \text{when } u \rightarrow 0^+ \quad (35)$$

On the other hand, when $u \rightarrow t$, $P_t(u) = r(t) s(t-u)$. For large p values, $S(p) \rightarrow \frac{1}{p}$ and finally

$$P_t(u) \rightarrow \frac{1}{p(t-u)} r(t) \quad ; \quad \text{when } u \rightarrow t \quad (36)$$

Let us now develop some examples.

For the graph of Figure 1, we get:

$$L = \frac{1}{p(1+)} \frac{\prod_{i=1}^n p_{0i} \text{th}(\frac{p}{2} l_{0i})}{\prod_{i=1}^n p_{0i} \text{th}(\frac{p}{2(1+)} l_{0i})} \quad (37)$$

$$\rightarrow \frac{1}{p(1+)} \quad ; \quad \text{when } l_{0i} \rightarrow 1 \quad ; \quad i = 1, \dots, n \quad (38)$$

¹ This is readily done with the help of the Dirichlet Green's function:

$$G_D(x; x^0;) = G_0(x; x^0;) - \frac{G_0(x; 0;) G_0(0; x^0;)}{G_0(0; 0;)}$$

Using completeness and also the relationship

$$\int_{\text{Graph}} dx^0 G_0(x; x^0;) = \frac{1}{p}$$

all the integrations in (7) can be performed, leading to (32).




Figure 1: a star-graph with n legs of lengths l_{0i} , $i = 1;:::n$, originating from O .

When all the legs become infinite, L does not depend on the exit probabilities from O , p_{0i} , because in that case all the legs are equivalent. The same behaviour occurs for a finite symmetric star ($l_{01} = \dots = l_{0n} = 1$) where we get

$$L = \frac{1}{p} \frac{\text{th}(\frac{p}{2} - 1)}{\text{th}(\frac{p}{2} + 1)} \quad (39)$$

that leads to the density

$$P_t(u) = \frac{1}{p} \frac{1}{u(t-u)} \sum_{p=1}^{\infty} \left(\frac{1}{p} \right)^p e^{\frac{2p^2 t^2}{u}} A \sum_{q=1}^{\infty} \left(\frac{1}{q} \right)^q e^{\frac{2q^2 t^2}{u}} A \quad (40)$$

$$= \frac{1}{2t^2} \sum_{m=1}^{\infty} \left(\frac{1}{m} \right)^m e^{\frac{(2m-1)^2 t^2}{8t^2}} \sum_{n=1}^{\infty} \left(\frac{1}{n} \right)^n e^{\frac{n^2 t^2}{2t^2}} \quad (41)$$

This last formula allows for studying the large u and t behaviours of $P_t(u)$. For instance:

$$P_t(u) \sim \frac{1}{t^2} e^{-\frac{t^2(t-u)}{8t^2}} \quad \text{when } u \sim t^2 \text{ and } (t-u) \sim 1 \quad (42)$$

In Figure 2, we consider a ring where the vertices O and 1 are linked by two bonds of lengths $l_{01}^{(1)}$ and $l_{01}^{(2)}$ ($1 = l_{01}^{(1)} + l_{01}^{(2)}$). The exit probabilities from O (resp. 1) are $p_{01}^{(1)}$ and $p_{01}^{(2)}$ (resp. $p_{10}^{(1)}$ and $p_{10}^{(2)}$) and we define $c_{01}^{(i)}(\cdot) = \coth \frac{p}{2} l_{01}^{(i)}$, $s_{01}^{(i)}(\cdot) = \frac{1}{\sinh \frac{p}{2} l_{01}^{(i)}}$, $i = 1;2$.




Figure 2: a ring with two vertices, O and 1 , linked by two bonds (1) and (2).

Adding intermediate vertices on each bond, we easily show that the matrix M writes:

$$M = \begin{pmatrix} 0 & \prod_{i=1}^2 p_{01}^{(i)} c_{01}^{(i)} & \prod_{i=1}^2 p_{10}^{(i)} s_{01}^{(i)} \\ \prod_{i=1}^2 p_{01}^{(i)} s_{01}^{(i)} & \prod_{i=1}^2 p_{10}^{(i)} c_{01}^{(i)} & 1 \end{pmatrix}$$

When $p_{10}^{(1)} = p_{10}^{(2)} = 1/2$, L does not depend on $p_{01}^{(i)}$ and we get:

$$L = \frac{1}{\left(\frac{1}{2} + \frac{1}{2} \right) \frac{\text{th} \left(\frac{1}{2} \right)}{\text{th} \left(\frac{1}{2} \right) + \frac{1}{2}}} \quad (43)$$

In that case, the ring can be viewed as a star with two legs of equal lengths, $l=2$, glued at their endpoints. Thus, (43) is simply (39) where l has been replaced by $l=2$.

Remark, however, that we should obtain a quite different result when taking $p_{01}^{(1)} = p_{01}^{(2)} = 1/2$:

$$L = \frac{1}{\left(\frac{1}{2} + \frac{1}{2} \right) \frac{\coth \left(\frac{1}{2} \right) + F \left(\frac{1}{2} \right)}{\coth \left(\frac{1}{2} \right) + F \left(\frac{1}{2} \right)}} \quad (44)$$

with

$$F \left(\frac{1}{2} \right) = \frac{p_{10}^{(1)} p_{10}^{(2)} \sinh \frac{1}{2} l_{01}^{(2)} \frac{1}{2}}{\cosh \frac{1}{2} l_{01}^{(2)} + 1} \quad (45)$$

Let us now turn to a brownian motion starting at the root O of a regular tree of coordination number z and depth n (see Figure 3 part a) for $z = 3$ and $n = 2$). We suppose that all the links have the same length l and also that

$$p_i = \frac{1}{m} = \frac{1}{z} \quad \text{for any vertex inside the graph} \quad (46)$$

$$= 1 \quad \text{otherwise} \quad (47)$$




Figure 3: part a) a tree with coordination number $z = 3$ and depth $n = 2$; part b) the equivalent linear graph with exit probabilities given in (48) and (49).

For the last passage problem in O this tree is equivalent to a $(n+1)$ linear graph with the exit probabilities

$$p_{01} = 1 = p_{n,n-1} \quad (48)$$

$$p_{i,i-1} = \frac{1}{z} ; \quad p_{i,i+1} = 1 - \frac{1}{z} ; \quad i = 1; \dots; n-1 \quad (49)$$

With $c(\cdot) = \coth \frac{p}{2} 1$ and $s(\cdot) = \frac{1}{\sinh \frac{p}{2} 1}$, we can write the relations

$$\det M(\cdot) = c(\cdot) \det M_{n-1}(\cdot) - \frac{1}{z} (s(\cdot))^2 \det M_{n-1}(\cdot) \quad (50)$$

$$\det M_{n-k}(\cdot) = c(\cdot) \det M_{n-k-1}(\cdot) + \frac{1}{z} \frac{1}{z} 1 (s(\cdot))^2 \det M_{n-k-2}(\cdot) ; \quad 0 \leq k \leq n \quad (51)$$

$$\det M_2(\cdot) = 1 + \frac{1}{z} (s(\cdot))^2 \quad (52)$$

$$\det M_1(\cdot) = c(\cdot) \quad (53)$$

M_{n-k} is the matrix M where the $(k+1)$ first lines and first columns have been deleted. Recursion relations can also be written for $M^{-1}_{00} = \frac{\det M_n}{\det M}$. In the limit $n \rightarrow \infty$, we get:

$$L = \frac{1}{p \frac{1}{(1+z)}} \frac{c(\cdot) 1 - \frac{2}{z} + \frac{1}{1 + 1 - \frac{2}{z}} (s(\cdot))^2}{c(\cdot) 1 - \frac{2}{z} + \frac{1}{1 + 1 - \frac{2}{z}} (s(\cdot))^2} \quad (54)$$

As expected, $z = 2$ leads to the second arc-sine Levy's law.

As an example, let us consider an infinite 2D square lattice of stepsize 1 with exit probabilities equal to $1/4$ in any vertex. A standard computation based on the tight binding model [12] leads to:

$$L = \frac{1}{p \frac{1}{(1+z)}} \frac{\frac{p}{2(1+z)} 1}{\frac{p}{2} 1} \frac{K \frac{1}{\cosh \frac{p}{2(1+z)} 1}}{K \frac{1}{\cosh \frac{p}{2} 1}} \quad (55)$$

$K(\cdot)$ is a complete elliptic integral of first kind [13].

To complete this work, let us come back to a general graph and choose equal exit probabilities in each vertex:

$$p_{ij} = \frac{1}{m} ; \quad i = 1; \dots; m \quad (56)$$

In that case, L can be expressed in terms of primitive orbits on the graph. Recall that an orbit \mathcal{C} is said to be primitive if it cannot be decomposed as a repetition of any smaller orbit. From [14], we get:

$$\sum_{\mathcal{C}} \frac{1}{l(\mathcal{C})} e^{p \frac{1}{2} l(\mathcal{C})} = \frac{1}{4} \sum_{\mathcal{C}} \frac{1}{l(\mathcal{C})} e^{p \frac{1}{2} l(\mathcal{C})} \frac{1}{\det M(\cdot)} \quad (57)$$

The product on the left hand-side is taken over all the primitive orbits \mathcal{C} of the graph. $l(\mathcal{C})$ is the length of \mathcal{C} and the weight $w(\mathcal{C})$ is the product of all the reflection { or transmission } factors along \mathcal{C} . (An orbit is a closed succession of arcs $[(i, j)]$ and a reflection (transmission)

occurs in \mathcal{C} if $\mathcal{C} = (\mathcal{C}_0)$. The corresponding factors are $\frac{2}{m} - 1$ for a reflection and $\frac{2}{m}$ for a transmission).

Now, imposing Dirichlet boundary conditions in O , we get, still with [14]:

$$Y_{\mathcal{C}}^{(0)} = \frac{1}{\det M_D(\mathcal{C})} \sum_{\mathcal{C}_0} \frac{e^{p \frac{2}{m} l(\mathcal{C}_0)}}{e^{p \frac{2}{m} l(\mathcal{C})}} = \frac{1}{\det M_D(\mathcal{C})} \sum_{\mathcal{C}_0} \frac{e^{p \frac{2}{m} l(\mathcal{C}_0)}}{e^{p \frac{2}{m} l(\mathcal{C})}} \quad (58)$$

$Y_{\mathcal{C}}^{(0)}$ is the same as $Y_{\mathcal{C}}$ except for orbits visiting O where the reflection and transmission factors are respectively -1 and 0 .

It is now easy to realize that the ratio $\frac{\det M_D(\mathcal{C})}{\det M(\mathcal{C})}$ only involves the primitive orbits \mathcal{C}_0 that visit O at least once. Thus:

$$M^{-1}(\mathcal{C})_{00} = \frac{\sum_{\mathcal{C}_0} \frac{1}{\det M(\mathcal{C}_0)} e^{p \frac{2}{m} l(\mathcal{C}_0)}}{\sum_{\mathcal{C}_0} \frac{1}{\det M(\mathcal{C}_0)} e^{p \frac{2}{m} l(\mathcal{C}_0)}} \quad (59)$$

Finally:

$$L = \frac{1}{\det M(\mathcal{C})} \sum_{\mathcal{C}_0} \frac{Y_{\mathcal{C}_0}^{(0)}}{\det M(\mathcal{C}_0)} e^{p \frac{2}{m} l(\mathcal{C}_0)} = \frac{1}{\det M(\mathcal{C})} \sum_{\mathcal{C}_0} \frac{Y_{\mathcal{C}_0}^{(0)}}{\det M(\mathcal{C}_0)} e^{p \frac{2}{m} l(\mathcal{C}_0)} \quad (60)$$

$Y_{\mathcal{C}_0}^{(0)}$

means that we only consider primitive orbits with reflections in O .

\mathcal{C}_0

As an example, let us see how (60) works with the star-graph of Figure 1 with $n = 2$ legs. Only one primitive orbit, with length $2(l_{01} + l_{02})$ and two reflection factors equal to -1 , contributes to the product.

\mathcal{C}_0

Concerning $Y_{\mathcal{C}_0}^{(0)}$, two orbits, with lengths $2l_{01}$ and $2l_{02}$, will contribute. For each one, the two

\mathcal{C}_0

reflection factors are (-1) and 1 . For L , we get the result:

$$L = \frac{1}{\det M(\mathcal{C})} \frac{(1 + e^{p \frac{2}{m} l_{01}})(1 + e^{p \frac{2}{m} l_{02}})}{(1 + e^{p \frac{2}{m} l_{01}})(1 + e^{p \frac{2}{m} l_{02}})} \frac{(1 - e^{p \frac{2}{m} (l_{01} + l_{02})})}{(1 - e^{p \frac{2}{m} (l_{01} + l_{02})})} \quad (61)$$

$$= \frac{1}{\det M(\mathcal{C})} \frac{\cosh \frac{p}{2} l_{01} \cosh \frac{p}{2} l_{02}}{\cosh \frac{p}{2} l_{01} \cosh \frac{p}{2} l_{02}} \frac{\sinh \frac{p}{2} (l_{01} + l_{02})}{\sinh \frac{p}{2} (l_{01} + l_{02})} \quad (62)$$

in agreement with (37).

For the ring of Figure 2, two (time-reversed) orbits of length l should contribute to $Y_{\mathcal{C}_0}$ and only

one of length $2l$ should contribute to $Y_{\mathcal{C}_0}^{(0)}$. So:

$$L = \frac{1}{p(+) } \frac{(1 - e^{\frac{p}{2(+)1}})}{(1 - e^{\frac{p}{2}1})} \frac{(1 - e^{\frac{p}{2}1})^2}{(1 - e^{\frac{p}{2(+)1}2}} \quad (63)$$

$$= \frac{1}{p(+) } \frac{\text{th}^{\frac{p}{2}1}}{\text{th}^{\frac{p}{2(+)1}2}} \quad (64)$$

as given by (43).

References

- [1] Levy P 1948 *Processus Stochastiques et Mouvement Brownien* (Paris: Editions Jacques Gabay)
- [2] RammalR 1984 *J.Phys. I (France)* 45 191
 Doucot B and RammalR 1985 *Phys.Rev.Lett.* 55 (10) 1148 ; 1986 *J.Physique* 47 973
 Avron J E and Sadun L 1991 *Ann.Phys. (N.Y.)* 206 440
 Texier C and Montambaux G 2001 *J.Phys.A :M ath.Gen.* 34 10307
- [3] Ihara Y 1966 *J.M ath.Soc. Japan* 18 219
 Roth J P 1983 *C .R .Acad.Sc.Paris* 296 793
 Stark H M and Terras A A 1996 *Adv. in M ath.* 121 124
 W u F Y and Kunz H 1999 *Ann.Combin.* 3 475
- [4] Alexander S 1983 *Phys.Rev.B* 27 (3) 1541
- [5] Rudenberg K and Scherr C 1953 *J.Chem .Phys.* 21 1565
- [6] Pascaud M and Montambaux G 1999 *Phys.Rev.Lett.* 82 4512
- [7] Dhar D 1999 *cond-mat/9909009*
- [8] Kottos T and Smilansky U 1997 *Phys.Rev.Lett.* 79 (24) 4794 ; 1999 *Ann.Phys. (N.Y.)* 274 (1) 76
- [9] Barlow M , Pitman J and Yor M 1989 *Sem .Probabilites XX III, Lect.Notes in M aths.* 1372 (Springer) page 294
- [10] Desbois J 2002 *J.Phys.A :M ath.Gen.* 35 L673
- [11] Comtet A , Desbois J and Majumdar S N 2002 *J.Phys.A :M ath.Gen.* 35 L687
- [12] Economou E N 1983 *Green's Functions in Quantum Physics* (Berlin: Springer)
- [13] Gradshteyn I S and Ryzhik I M 1980 *Table of Integrals, Series, and Products* (New York: Academic)
- [14] Desbois J 2001 *Eur.Phys.J.B* 24 261