

The Last Passage Problem on Graphs

Jean Desbois¹ and Olivier Bénichou²

25th May 2020

¹ Laboratoire de Physique Théorique et Modèles Statistiques. Université Paris-Sud, Bât. 100, F-91405 Orsay Cedex, France.

² Laboratoire de Physique de la Matière Condensée, Collège de France, 11, place M. Berthelot, 75231 Paris Cedex 05, France.

Abstract

We consider a brownian motion on a general graph, that starts at time $t = 0$ from some vertex O and stops at time t somewhere on the graph. Denoting by g the last time when O is reached, we establish a simple expression for the Laplace Transform, L , of the probability density of g . We discuss this result for some special graphs like star, ring, tree or square lattice. Finally, we show that L can also be expressed in terms of primitive orbits when, for any vertex, all the exit probabilities are equal.

Around the years 1930, P Levy [1] got several arc-sine laws concerning the 1D Brownian Motion on an infinite line. Let us consider such a process starting at $t = 0$ from the origin O and stopping at time t and denote by g the last time when O is reached. The second arc-sine law discovered by Levy concerns the probability law of g . It can be stated as follows:

$$P(g < u) = \frac{2}{\pi} \arcsin \sqrt{\frac{u}{t}} \quad (1)$$

$$\text{with the probability density} \quad \mathcal{P}_t(u) \equiv \frac{dP(g < u)}{du} = \frac{1}{\pi} \frac{1}{\sqrt{u(t-u)}} \quad (2)$$

In particular, the Laplace transform of $\mathcal{P}_t(u)$ is written:

$$\int_0^\infty dt e^{-\gamma t} \int_0^t du \mathcal{P}_t(u) e^{-\xi u} = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \quad (3)$$

Levy also proved that the same law occurs when g represents the time spent in the region ($x > 0$) (first arc-sine law).

The infinite line with only one point, O , specified, can be viewed as a kind of very simple graph consisting in one vertex, O , and two semi-infinite lines originating from O .

Our goal in this Letter is to get an analogue of the second arc-sine law (3) but for a quite general graph.

Let us recall that graphs properties have interested for many years physicists [2] as well as mathematicians [3]. For instance, the knowledge of spectral properties of the Laplacian operator is useful to understand superconducting networks [4] as well as organic molecules [5]. We also know that spectral determinants play a central role in the computation of weak localization corrections [6]. Recently, graphs have also been used in the context of nonequilibrium statistical physics (see [7] and references therein) and also in the field of quantum chaos [8].

The study of Brownian motion on graphs is also an active area of research. For instance, occupation times [9, 10] or local time [11] distributions have been recently computed for general graphs.

So, let us begin by setting the notations that will be used throughout this Letter. We consider a graph made of V vertices, numbered from 0 to $V - 1$, linked by B bonds of finite lengths. The coordination number of vertex α is m_α ($\sum_{\alpha=0}^{V-1} m_\alpha = 2B$).

On each bond $[\alpha\beta]$, of length $l_{\alpha\beta}$, we define the coordinate $x_{\alpha\beta}$ that runs from 0 (vertex α) to $l_{\alpha\beta}$ (vertex β). Sometimes we will also use the coordinate $x_{\beta\alpha} \equiv l_{\alpha\beta} - x_{\alpha\beta}$. The set of coordinates $\{x_{\alpha\beta}\}$ is simply noted x .

An arc $(\alpha\beta)$ is defined as the oriented bond from α to β . Each bond $[\alpha\beta]$ is therefore associated with two arcs $(\alpha\beta)$ and $(\beta\alpha)$. In the sequel, we will consider the following ordering of the $2B$ arcs: $(0\mu_1)(0\mu_2) \dots (0\mu_{m_0}) \dots (\alpha\beta_1) \dots (\alpha\beta_i) \dots (\alpha\beta_{m_\alpha}) \dots$

Now, let us consider a brownian motion starting at some vertex O (label 0) of the graph at time $t = 0$ and stopped at time t . Let g be the last time when O is visited. The probability, $P_t(g < u)$ can be written:

$$P_t(g < u) = \int_{\text{Graph}} dx' dx G_0(0, 0; x, u) G_D(x, u; x', t) \quad (4)$$

where G_0 is the free propagator on the graph and G_D is the propagator with Dirichlet boundary conditions in O . (4) simply expresses the fact that the brownian particle will not hit O between the times u and t . Let us define the probability density $\mathcal{P}_t(u) \equiv \frac{dP(g < u)}{du}$. In the sequel, we will be especially interested in the Laplace transforms:

$$\mathcal{L} \equiv \int_0^\infty dt e^{-\gamma t} \int_0^t du P_t(g < u) e^{-\xi u} \quad (5)$$

$$L \equiv \int_0^\infty dt e^{-\gamma t} \int_0^t du \mathcal{P}_t(u) e^{-\xi u} = \frac{1}{\gamma + \xi} + \xi \mathcal{L} \quad (6)$$

With (4), we see that:

$$\mathcal{L} = \int_{\text{Graph}} dx' dx \left\langle 0 \left| \frac{1}{H + \gamma + \xi} \right| x \right\rangle \left\langle x \left| \frac{1}{H_D + \gamma} \right| x' \right\rangle \equiv \int_{\text{Graph}} dx' dx \mathcal{G}_0(0, x; \gamma + \xi) \mathcal{G}_D(x, x'; \gamma) \quad (7)$$

On the bond $[\alpha\beta]$, H and H_D are equal to $-\frac{1}{2} \frac{d^2}{dx_{\alpha\beta}^2}$ and we define:

$$\mathcal{G}_0^{(\alpha\beta)} = \lim_{x_{\alpha\beta} \rightarrow 0} \mathcal{G}_0(0, x_{\alpha\beta}; \gamma + \xi) \quad ; \quad \mathcal{G}'_0^{(\alpha\beta)} = \lim_{x_{\alpha\beta} \rightarrow 0} \frac{d\mathcal{G}_0(0, x_{\alpha\beta}; \gamma + \xi)}{dx_{\alpha\beta}} \quad (8)$$

$$\mathcal{G}_D^{(\alpha\beta)}(x) = \lim_{x'_{\alpha\beta} \rightarrow 0} \mathcal{G}_D(x, x'_{\alpha\beta}; \gamma) \quad ; \quad \mathcal{G}'_D^{(\alpha\beta)}(x) = \lim_{x'_{\alpha\beta} \rightarrow 0} \frac{d\mathcal{G}_D(x, x'_{\alpha\beta}; \gamma)}{dx'_{\alpha\beta}} \quad (9)$$

The integration over x' in (7) is easily performed with the help of the equation

$$\left(-\frac{1}{2} \frac{d^2}{dx_{\alpha\beta}^2} + \gamma\right) \mathcal{G}_D(x, x'_{\alpha\beta}; \gamma) = \delta(x - x'_{\alpha\beta}) \quad (10)$$

We get

$$\int_{\text{Graph}} dx' \mathcal{G}_D(x, x'; \gamma) = \frac{1}{\gamma} - \frac{1}{2\gamma} \sum_{i=1}^{m_0} \mathcal{G}'_D^{(0\mu_i)}(x) \quad (11)$$

To go further, we have to specify the behaviours of the resolvents \mathcal{G}_0 and \mathcal{G}_D in the neighbourhood of all the vertices.

Let us consider some vertex α with its nearest neighbours β_i , $i = 1, 2, \dots, m_\alpha$, on the graph. Suppose that the brownian particle reaches α . It will come out towards β_i with some exit probability $p_{\alpha\beta_i}$. This implies the following equations to be satisfied by \mathcal{G}_0 :

$$\frac{\mathcal{G}_0^{(\alpha\beta_1)}}{p_{\alpha\beta_1}} = \frac{\mathcal{G}_0^{(\alpha\beta_2)}}{p_{\alpha\beta_2}} = \dots = \frac{\mathcal{G}_0^{(\alpha\beta_{m_\alpha})}}{p_{\alpha\beta_{m_\alpha}}} \equiv f_\alpha \quad , \quad \forall \alpha \quad (12)$$

Remark that the resolvent will be continuous in vertex α only when the particle exits from α with the same probability in all the directions.

Moreover, current conservation implies:

$$\sum_{i=1}^{m_\alpha} \mathcal{G}'_0^{(\alpha\beta_i)} = 0 \quad , \quad \text{if } \alpha \neq 0 \quad (13)$$

$$\sum_{i=1}^{m_0} \mathcal{G}'_0^{(0\mu_i)} = -2 \quad (14)$$

On the link $[\alpha\beta]$, $\mathcal{G}_0(0, x_{\alpha\beta}; \gamma + \xi)$ must satisfy:

$$\left(-\frac{1}{2} \frac{d^2}{dx_{\alpha\beta}^2} + \gamma + \xi\right) \mathcal{G}_0(0, x_{\alpha\beta}; \gamma + \xi) = 0 \quad (15)$$

with the solution:

$$\mathcal{G}_0(0, x_{\alpha\beta}; \gamma + \xi) = \mathcal{G}_0^{(\alpha\beta)} \frac{\sinh \sqrt{2(\gamma + \xi)}(l_{\alpha\beta} - x_{\alpha\beta})}{\sinh \sqrt{2(\gamma + \xi)}l_{\alpha\beta}} + \mathcal{G}_0^{(\beta\alpha)} \frac{\sinh \sqrt{2(\gamma + \xi)}x_{\alpha\beta}}{\sinh \sqrt{2(\gamma + \xi)}l_{\alpha\beta}} \quad (16)$$

So, we deduce:

$$\mathcal{G}'_0^{(\alpha\beta)} = \sqrt{2(\gamma + \xi)} \left(-c_{\beta\alpha}(\gamma + \xi) \mathcal{G}_0^{(\alpha\beta)} + s_{\alpha\beta}(\gamma + \xi) \mathcal{G}_0^{(\beta\alpha)}\right) \quad (17)$$

with:

$$c_{\alpha\beta}(\gamma + \xi) = \coth \sqrt{2(\gamma + \xi)}l_{\alpha\beta} = c_{\beta\alpha}(\gamma + \xi) \quad (18)$$

$$s_{\alpha\beta}(\gamma + \xi) = \frac{1}{\sinh \sqrt{2(\gamma + \xi)}l_{\alpha\beta}} = s_{\beta\alpha}(\gamma + \xi) \quad (19)$$

Those considerations allow to write eqs.(12, 13, 14) in a matrix form:

$$M(\gamma + \xi) f = \frac{1}{\sqrt{2(\gamma + \xi)}} l \quad (20)$$

where f and l are two $(V \times 1)$ vectors. The components of f are the quantities f_α and for the components of l we have $l_j = 2\delta_{j0}$.

$M(\gamma + \xi)$ is a $(V \times V)$ vertex matrix with the non vanishing elements (α runs from 0 to $V - 1$):

$$M(\gamma + \xi)_{\alpha\alpha} = \sum_{i=1}^{m_\alpha} p_{\alpha\beta_i} c_{\alpha\beta_i}(\gamma + \xi) \quad (21)$$

$$M(\gamma + \xi)_{\alpha\beta} = -p_{\beta\alpha} s_{\alpha\beta}(\gamma + \xi) \quad \text{if } [\alpha\beta] \text{ is a bond} \quad (22)$$

$$= 0 \quad \text{otherwise} \quad (23)$$

Similar considerations hold for $\mathcal{G}_D(x, x'; \gamma)$ and we get:

$$\frac{\mathcal{G}_D^{(\alpha\beta_1)}(x)}{p_{\alpha\beta_1}} = \dots = \frac{\mathcal{G}_D^{(\alpha\beta_{m_\alpha})}(x)}{p_{\alpha\beta_{m_\alpha}}} \equiv F_\alpha(x) \quad , \quad \text{if } \alpha \neq 0 \quad (24)$$

$$\mathcal{G}_D^{(0\mu_i)}(x) = 0 \quad ; \quad i = 1, \dots, m_0 \quad (25)$$

$$\sum_{i=1}^{m_\alpha} \mathcal{G}'^{(\alpha\beta_i)}(x) = 0 \quad , \quad \text{if } \alpha \neq 0 \quad (26)$$

$$\text{and the matrix equation} \quad M_D(\gamma)F(x) = \frac{1}{\gamma} l_D(x) \quad (27)$$

M_D is the matrix M with the first line and first column deleted. $F(x)$ and $l_D(x)$ are $(V - 1) \times 1$ vectors. The components of $F(x)$ are the quantities $F_\alpha(x)$, $\alpha = 1, \dots, V - 1$. The nonvanishing components of $l_D(x)$ are defined as follows.

If $x \in [ab]$ with $a \neq 0$ and $b \neq 0$ then

$$(l_D(x_{ab}))_a = s_{ab}(\gamma) \sinh(\sqrt{2\gamma} x_{ba}) \quad (28)$$

$$(l_D(x_{ab}))_b = s_{ab}(\gamma) \sinh(\sqrt{2\gamma} x_{ab}) \quad (29)$$

If $x \in [0\mu_j]$ then

$$(l_D(x_{0\mu_j}))_{\mu_j} = s_{0\mu_j}(\gamma) \sinh(\sqrt{2\gamma} x_{0\mu_j}) \quad (30)$$

Performing the integration over x in (7), we finally get the following simple expression for L :

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{(M^{-1}(\gamma + \xi))_{00}}{(M^{-1}(\gamma))_{00}} \quad (31)$$

This is the central result of this paper.

Remark that, for the special case when \mathcal{G}_0 and \mathcal{G}_D are continuous Green's functions, we can recast (31) in the form ¹

$$L = \frac{1}{\gamma} \frac{\mathcal{G}_0(0, 0; \gamma + \xi)}{\mathcal{G}_0(0, 0; \gamma)} \quad (32)$$

To go further and deduce from (31) some general properties of the density $\mathcal{P}_t(u)$, let us consider the following identity

$$\int_0^\infty dt e^{-\gamma t} \int_0^t du e^{-\xi u} r(u) s(t-u) = \mathcal{R}(\gamma + \xi) \mathcal{S}(\gamma) \quad (33)$$

where \mathcal{R} and \mathcal{S} are the Laplace Transforms of r and s .

Setting $\mathcal{R}(p) = \frac{(M^{-1}(p))_{00}}{\sqrt{p}}$ and $\mathcal{S}(p) = \frac{1}{\sqrt{p} (M^{-1}(p))_{00}}$ in (33), we realize from (31) and (6) that the probability density factorizes in the following way:

$$\mathcal{P}_t(u) = r(u).s(t-u) \quad (34)$$

This result holds for any graph.

It is now easy to analyze the limiting behaviours of $\mathcal{P}_t(u)$ when $u \rightarrow 0^+$ and when $u \rightarrow t^-$.

In the first case, we can write $\mathcal{P}_t(u) \approx r(u)s(t)$. Moreover, for large p values, $M(p) \rightarrow 1$ and $\mathcal{R}(p) \sim \frac{1}{\sqrt{p}}$. So

$$\mathcal{P}_t(u) \sim \frac{1}{\sqrt{\pi u}} s(t) \quad , \quad \text{when } u \rightarrow 0^+ \quad (35)$$

On the other hand, when $u \rightarrow t^-$, $\mathcal{P}_t(u) \approx r(t)s(t-u)$. For large p values, $\mathcal{S}(p) \sim \frac{1}{\sqrt{p}}$ and finally

$$\mathcal{P}_t(u) \sim \frac{1}{\sqrt{\pi (t-u)}} r(t) \quad , \quad \text{when } u \rightarrow t^- \quad (36)$$

Let us now develop some examples.

For the graph of Figure 1, we get:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{\sum_{i=1}^n p_{0i} \text{th}(\sqrt{2\gamma} l_{0i})}{\sum_{i=1}^n p_{0i} \text{th}(\sqrt{2(\gamma + \xi)} l_{0i})} \quad (37)$$

$$\rightarrow \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \quad , \quad \text{when } l_{0i} \rightarrow \infty \quad , \quad i = 1, \dots, n \quad (38)$$

¹This is readily done with the help of the Dirichlet Green's function:

$$\mathcal{G}_D(x, x'; \gamma) = \mathcal{G}_0(x, x'; \gamma) - \frac{\mathcal{G}_0(x, 0; \gamma) \mathcal{G}_0(0, x'; \gamma)}{\mathcal{G}_0(0, 0; \gamma)}$$

Using completeness and also the relationship

$$\int_{\text{Graph}} dx' \mathcal{G}_0(x, x'; \lambda) = \frac{1}{\lambda}$$

all the integrations in (7) can be performed, leading to (32).




Figure 1: a star-graph with n legs of lengths l_{0i} , $i = 1, \dots, n$, originating from O .

When all the legs become infinite, L does not depend on the exit probabilities from O , p_{0i} , because in that case all the legs are equivalent. The same behaviour occurs for a finite symmetric star ($l_{01} = \dots = l_{0n} \equiv l$) where we get

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{\text{th}(\sqrt{2\gamma}l)}{\text{th}(\sqrt{2(\gamma + \xi)}l)} \quad (39)$$

that leads to the density

$$\mathcal{P}_t(u) = \frac{1}{\pi\sqrt{u(t-u)}} \left(\sum_{p=-\infty}^{+\infty} (-1)^p e^{-\frac{2p^2 l^2}{t-u}} \right) \left(\sum_{q=-\infty}^{+\infty} e^{-\frac{2q^2 l^2}{u}} \right) \quad (40)$$

$$= \frac{1}{2l^2} \left(\sum_{m=-\infty}^{+\infty} e^{-\frac{(2m-1)^2 \pi^2 (t-u)}{8l^2}} \right) \left(\sum_{n=-\infty}^{+\infty} e^{-\frac{n^2 \pi^2 u}{2l^2}} \right) \quad (41)$$

This last formula allows for studying the large u and t behaviours of $\mathcal{P}_t(u)$. For instance:

$$\mathcal{P}_t(u) \sim \frac{1}{l^2} e^{-\frac{\pi^2(t-u)}{8l^2}} \quad \text{when } u \gg l^2 \text{ and } (t-u) \gg l^2 \quad (42)$$

In Figure 2, we consider a ring where the vertices O and 1 are linked by two bonds of lengths $l_{01}^{(1)}$ and $l_{01}^{(2)}$ ($l \equiv l_{01}^{(1)} + l_{01}^{(2)}$). The exit probabilities from O (resp. 1) are $p_{01}^{(1)}$ and $p_{01}^{(2)}$ (resp. $p_{10}^{(1)}$ and $p_{10}^{(2)}$) and we define $c_{01}^{(i)}(\gamma) = \coth \sqrt{2\gamma}l_{01}^{(i)}$, $s_{01}^{(i)}(\gamma) = \frac{1}{\sinh \sqrt{2\gamma}l_{01}^{(i)}}$, $i = 1, 2$.




Figure 2: a ring with two vertices, O and 1 , linked by two bonds (1) and (2).

Adding intermediate vertices on each bond, we easily show that the matrix M writes:

$$M = \begin{pmatrix} \sum_{i=1}^2 p_{01}^{(i)} c_{01}^{(i)} & -\sum_{i=1}^2 p_{10}^{(i)} s_{01}^{(i)} \\ -\sum_{i=1}^2 p_{01}^{(i)} s_{01}^{(i)} & \sum_{i=1}^2 p_{10}^{(i)} c_{01}^{(i)} \end{pmatrix}$$

When $p_{10}^{(1)} = p_{10}^{(2)} = 1/2$, L does not depend on $p_{01}^{(i)}$ and we get:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{\text{th}(\sqrt{2\gamma} \frac{l}{2})}{\text{th}(\sqrt{2(\gamma + \xi)} \frac{l}{2})} \quad (43)$$

In that case, the ring can be viewed as a star with two legs of equal lengths, $l/2$, glued at their endpoints. Thus, (43) is simply (39) where l has been replaced by $l/2$.

Remark, however, that we should obtain a quite different result when taking $p_{01}^{(1)} = p_{01}^{(2)} = 1/2$:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{\coth(\sqrt{2(\gamma + \xi)} \frac{l}{2}) + F(\gamma + \xi)}{\coth(\sqrt{2\gamma} \frac{l}{2}) + F(\gamma)} \quad (44)$$

with

$$F(\gamma) = \frac{(p_{10}^{(1)} - p_{10}^{(2)}) \sinh \sqrt{2\gamma} (l_{01}^{(2)} - l_{01}^{(1)})}{\cosh \sqrt{2\gamma} l - 1} \quad (45)$$

Let us now turn to a brownian motion starting at the root O of a regular tree of coordination number z and depth n (see Figure 3 part a) for $z = 3$ and $n = 2$). We suppose that all the links have the same length l and also that

$$p_{\alpha\beta_i} = \frac{1}{m_\alpha} = \frac{1}{z} \quad \text{for any vertex inside the graph} \quad (46)$$

$$= 1 \quad \text{otherwise} \quad (47)$$




Figure 3: part a) a tree with coordination number $z = 3$ and depth $n = 2$; part b) the equivalent linear graph with exit probabilities given in (48) and (49).

For the last passage problem in O this tree is equivalent to a $(n + 1)$ linear graph with the exit probabilities

$$p_{01} = 1 = p_{n,n-1} \quad (48)$$

$$p_{i,i-1} = \frac{1}{z} \quad , \quad p_{i,i+1} = 1 - \frac{1}{z} \quad , \quad i = 1, \dots, n-1 \quad (49)$$

With $c(\gamma) = \coth \sqrt{2\gamma}l$ and $s(\gamma) = \frac{1}{\sinh \sqrt{2\gamma}l}$, we can write the relations

$$\det M(\gamma) = c(\gamma) \det M_n(\gamma) - \frac{1}{z}(s(\gamma))^2 \det M_{n-1}(\gamma) \quad (50)$$

$$\det M_{n-k}(\gamma) = c(\gamma) \det M_{n-k-1}(\gamma) + \frac{1}{z} \left(\frac{1}{z} - 1 \right) (s(\gamma))^2 \det M_{n-k-2}(\gamma) \quad , \quad 0 \leq k \leq n-1 \quad (51)$$

$$\det M_2(\gamma) = 1 + \frac{1}{z}(s(\gamma))^2 \quad (52)$$

$$\det M_1(\gamma) = c(\gamma) \quad (53)$$

M_{n-k} is the matrix M where the $(k+1)$ first lines and first columns have been deleted. Recursion relations can also be written for $(M^{-1})_{00} \equiv \frac{\det M_n}{\det M}$. In the limit $n \rightarrow \infty$, we get:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{c(\gamma) \left(1 - \frac{2}{z}\right) + \sqrt{1 + \left(1 - \frac{2}{z}\right)^2 (s(\gamma))^2}}{c(\gamma + \xi) \left(1 - \frac{2}{z}\right) + \sqrt{1 + \left(1 - \frac{2}{z}\right)^2 (s(\gamma + \xi))^2}} \quad (54)$$

As expected, $z = 2$ leads to the second arc-sine Levy's law.

As a final example, let us consider an infinite 2D square lattice of stepsize l with exit probabilities equal to $1/4$ in any vertex. A standard computation based on the tight binding model [12] leads to:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{\text{th} \sqrt{2(\gamma + \xi)l}}{\text{th} \sqrt{2\gamma}l} \frac{K \left(\frac{1}{\cosh \sqrt{2(\gamma + \xi)l}} \right)}{K \left(\frac{1}{\cosh \sqrt{2\gamma}l} \right)} \quad (55)$$

$K(\lambda)$ is a complete elliptic integral of first kind [13].

To complete this work, let us come back to a general graph and choose equal exit probabilities in each vertex:

$$\forall \alpha \quad , \quad p_{\alpha\beta_i} = \frac{1}{m_\alpha} \quad ; \quad i = 1, \dots, m_\alpha \quad (56)$$

In that case, L can be expressed in terms of primitive orbits on the graph. Recall that an orbit \tilde{C} is said to be primitive if it cannot be decomposed as a repetition of any smaller orbit. From [14], we get:

$$\prod_{\tilde{C}} \left(1 - \mu(\tilde{C}) e^{-\sqrt{2\gamma}l(\tilde{C})} \right) = \left[\prod_{[\alpha\beta]} \left(1 - e^{-2\sqrt{2\gamma}l_{\alpha\beta}} \right) \right] \det M(\gamma) \quad (57)$$

The product on the left hand-side is taken over all the primitive orbits \tilde{C} of the graph. $l(\tilde{C})$ is the length of \tilde{C} and the weight $\mu(\tilde{C})$ is the product of all the reflection – or transmission – factors along \tilde{C} . (An orbit is a closed succession of arcs $\dots(\tau\alpha)(\alpha\beta)\dots$ and a reflection (transmission)

occurs in α if $\tau = \beta$ ($\tau \neq \beta$). The corresponding factors are $\left(\frac{2}{m_\alpha} - 1\right)$ for a reflection and $\left(\frac{2}{m_\alpha}\right)$ for a transmission).

Now, imposing Dirichlet boundary conditions in O , we get, still with [14]:

$$\prod_{\tilde{C}} \left(1 - \mu^{(0)}(\tilde{C})e^{-\sqrt{2\gamma}l(\tilde{C})}\right) = \left[\prod_{[\alpha\beta]} \left(1 - e^{-2\sqrt{2\gamma}l_{\alpha\beta}}\right) \right] \det M_D(\gamma) \quad (58)$$

$\mu^{(0)}(\tilde{C})$ is the same as $\mu(\tilde{C})$ except for orbits visiting O where the reflection and transmission factors are respectively -1 and 0 .

It is now easy to realize that the ratio $\frac{\det M_D(\gamma)}{\det M(\gamma)}$ only involves the primitive orbits \tilde{C}_0 that visit O at least once. Thus:

$$\left(M^{-1}(\gamma)\right)_{00} = \frac{\prod_{\tilde{C}_0} \left(1 - \mu^{(0)}(\tilde{C}_0)e^{-\sqrt{2\gamma}l(\tilde{C}_0)}\right)}{\prod_{\tilde{C}_0} \left(1 - \mu(\tilde{C}_0)e^{-\sqrt{2\gamma}l(\tilde{C}_0)}\right)} \quad (59)$$

Finally:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \prod'_{\tilde{C}_0} \left(\frac{1 - \mu^{(0)}(\tilde{C}_0)e^{-\sqrt{2(\gamma+\xi)l(\tilde{C}_0)}}}{1 - \mu(\tilde{C}_0)e^{-\sqrt{2\gamma}l(\tilde{C}_0)}} \right) \prod_{\tilde{C}_0} \left(\frac{1 - \mu(\tilde{C}_0)e^{-\sqrt{2\gamma}l(\tilde{C}_0)}}{1 - \mu(\tilde{C}_0)e^{-\sqrt{2(\gamma+\xi)l(\tilde{C}_0)}}} \right) \quad (60)$$

$\prod'_{\tilde{C}_0}$ means that we only consider primitive orbits with reflections in O .

As an example, let us see how (60) works with the star-graph of Figure 1 with $n = 2$ legs. Only one primitive orbit, with length $2(l_{01} + l_{02})$ and two reflection factors equal to 1, contributes to the product $\prod_{\tilde{C}_0}$.

Concerning $\prod'_{\tilde{C}_0}$, two orbits, with lengths $2l_{01}$ and $2l_{02}$, will contribute. For each one, the two reflection factors are (-1) and 1 . For L , we get the result:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{(1 + e^{-2\sqrt{2(\gamma+\xi)l_{01}}})(1 + e^{-2\sqrt{2(\gamma+\xi)l_{02}}})}{(1 + e^{-2\sqrt{2\gamma}l_{01}})(1 + e^{-2\sqrt{2\gamma}l_{02}})} \frac{(1 - e^{-2\sqrt{2\gamma}(l_{01}+l_{02})})}{(1 - e^{-2\sqrt{2(\gamma+\xi)(l_{01}+l_{02})})} \quad (61)$$

$$= \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{\cosh \sqrt{2(\gamma + \xi)l_{01}} \cosh \sqrt{2(\gamma + \xi)l_{02}}}{\cosh \sqrt{2\gamma}l_{01} \cosh \sqrt{2\gamma}l_{02}} \frac{\sinh \sqrt{2\gamma}(l_{01} + l_{02})}{\sinh \sqrt{2(\gamma + \xi)(l_{01} + l_{02})} \quad (62)$$

in agreement with (37).

For the ring of Figure 2, two (time-reversed) orbits of length l should contribute to $\prod_{\tilde{C}_0}$ and only

one of length $2l$ should contribute to $\prod'_{\tilde{C}_0}$. So:

$$L = \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{(1 - e^{-2\sqrt{2(\gamma+\xi)l}})}{(1 - e^{-2\sqrt{2\gamma}l})} \frac{(1 - e^{-\sqrt{2\gamma}l})^2}{(1 - e^{-\sqrt{2(\gamma+\xi)l})^2}} \quad (63)$$

$$= \frac{1}{\sqrt{\gamma(\gamma + \xi)}} \frac{\text{th}\sqrt{2\gamma}\frac{l}{2}}{\text{th}\sqrt{2(\gamma + \xi)}\frac{l}{2}} \quad (64)$$

as given by (43).

References

- [1] Levy P 1948 *Processus Stochastiques et Mouvement Brownien* (Paris: Editions Jacques Gabay)
- [2] Rammal R 1984 *J. Phys. I (France)* **45** 191
 Douçot B and Rammal R 1985 *Phys. Rev. Lett.* **55**(10) 1148 ; 1986 *J. Physique* **47** 973
 Avron J E and Sadun L 1991 *Ann. Phys. (N.Y.)* **206** 440
 Texier C and Montambaux G 2001 *J. Phys. A: Math.Gen.* **34** 10307
- [3] Ihara Y 1966 *J. Math. Soc. Japan* **18** 219
 Roth J P 1983 *C. R. Acad. Sc. Paris* **296** 793
 Stark H M and Terras A A 1996 *Adv. in Math.* **121** 124
 Wu F Y and Kunz H 1999 *Ann. Combin.* **3** 475
- [4] Alexander S 1983 *Phys. Rev. B* **27**(3) 1541
- [5] Rudenberg K and Scherr C 1953 *J. Chem. Phys.* **21** 1565
- [6] Pascaud M and Montambaux G 1999 *Phys. Rev. Lett.* **82** 4512
- [7] Dhar D 1999 *cond-mat/9909009*
- [8] Kottos T and Smilansky U 1997 *Phys. Rev. Lett.* **79**(24) 4794 ; 1999 *Ann. Phys. (N.Y.)* **274**(1) 76
- [9] Barlow M, Pitman J and Yor M 1989 *Sém. Probabilités XXIII, Lect. Notes in Maths.* 1372 (Springer) page 294
- [10] Desbois J 2002 *J. Phys. A: Math. Gen.* **35** L673
- [11] Comtet A, Desbois J and Majumdar S N 2002 *J. Phys. A: Math. Gen.* **35** L687
- [12] Economou E N 1983 *Green's Functions in Quantum Physics* (Berlin: Springer)
- [13] Gradshteyn I S and Ryzhik I M 1980 *Table of Integrals, Series, and Products* (New York: Academic)
- [14] Desbois J 2001 *Eur. Phys. J. B* **24** 261