

Emptiness Formation Probability for the Anisotropic XY Spin Chain in a Magnetic Field

Alexander G. Abanov, Fabio Franchini

*Physics & Astronomy Department,
State University of New York,
Stony Brook, New York 11794-3800*

Abstract

We study an asymptotic behavior of the probability of formation of a ferromagnetic string (referred to as EFP) of length n in a ground state of the one-dimensional anisotropic XY model in a transversal magnetic field as $n \rightarrow \infty$. We find that it is exponential everywhere in the phase diagram of the XY model except at the critical lines where the spectrum is gapless. One of those lines corresponds to the isotropic XY model where EFP decays in a Gaussian way, as was shown in Ref.[1]. The other lines are at the critical value of the magnetic field. There, we show that EFP is still exponential but acquires a non-trivial power-law prefactor with a universal exponent.

Key words: Emptiness Formation Probability, Integrable Models, XY Spin Chain, Toeplitz Determinants, Fisher-Hartwig Conjecture
PACS: 75.10.Pq, 02.30.Ik, 02.30.Tb

1 Introduction

Emptiness Formation Probability (EFP) is a special correlation function which plays an important role in the theory of integrable models [2]. In the case of spin- $\frac{1}{2}$ chains at zero temperature it is defined as

$$P(n) \equiv \left\langle \prod_{j=1}^n \frac{1 - \sigma_j^z}{2} \right\rangle \quad (1)$$

where $\sigma_i^{x,y,z}$ are Pauli matrices, the integer j labels the sites of the 1d lattice, and the average in (1) is performed over the ground state of the Hamiltonian

of the spin chain. EFP measures the probability that n consecutive spin sites are all found aligned downward in the ground state. The asymptotic behavior of $P(n)$ at large n is, probably, the most interesting one from the physical point of view, as it is believed that it measures the tendency of the system to a phase separation.

There has been considerable progress in finding this asymptotic behavior in the case of the integrable XXZ spin chain in critical regime [2,3,1,4,5,6,7]. At finite temperature the asymptotic behavior of EFP is exponential [2] $P(n) \sim e^{-\gamma n}$, while at zero temperature in critical regime it is Gaussian $P(n) \sim e^{-\alpha n^2}$. The qualitative argument in favor of a Gaussian decay of EFP at zero temperature was given in Ref. [8] in the framework of a field theoretical approach. There, it was argued that an optimal fluctuation of the field corresponding to EFP in the critical model will have (very roughly) the form of a round droplet in space-time with area $\sim n^2$ and will have an action proportional to n^2 . The asymptotic form of $P(n)$ is defined by the action of this optimal configuration and is proportional to $P(n) \sim e^{-\alpha n^2}$, which explains the Gaussian decay of EFP at zero temperature. Similarly, at finite temperature T in the limit $n \gg 1/T$ the droplet becomes rectangular (one dimension n is replaced by the inverse temperature $1/T$) and the action cost is proportional to n , giving $P(n) \sim e^{-\gamma n}$. This argument is tied to the criticality of the theory. Indeed, for noncritical theory one should not expect the same scaling of time and space dimensions of an optimal configuration. It is, therefore, very interesting to see how the asymptotic behavior of EFP is related to the criticality of a theory.

In this paper we examine the relation between an asymptotic behavior of EFP and criticality using the example of the Anisotropic XY spin-1/2 chain in a transverse magnetic field h

$$H = \sum_j \left[\left(\frac{1+\gamma}{2} \right) \sigma_j^x \sigma_{j+1}^x + \left(\frac{1-\gamma}{2} \right) \sigma_j^y \sigma_{j+1}^y \right] - h \sum_j \sigma_j^z \quad (2)$$

This model has been solved in [9] in the case of zero magnetic field and in [10] in the presence of a magnetic field. The phase diagram is depicted in Fig. 1. The phase diagram of the XY model has obvious symmetries $\gamma \rightarrow -\gamma$ and $h \rightarrow -h$. However, the latter one is broken by EFP itself. Therefore, we show only the part of the diagram corresponding to $\gamma \geq 0$. The phase diagram has both critical and non-critical regimes. The first critical regime – the Isotropic ($\gamma = 0$) XY model – is labeled Ω_0 . There is another critical line $h = 1$ which is labeled Ω_+ where excitations are gapless (and similarly line Ω_- at $h = -1$). The rest of the phase diagram is non-critical with finite gap for excitations and is divided into domains Σ_+ , Σ_0 , and Σ_- for $h > 1$, $-1 < h < 1$, and $h < -1$ respectively. Fig. 1 includes the line $\gamma = 1$ (Γ_I) corresponding to an Ising model in transverse magnetic field and the line $\gamma^2 + h^2 = 1$ (Γ_E) on which the wave function of the ground state is factorized into a product of




Fig. 1. Phase diagram of the XY Model.

single spin states [11].

2 The Emptiness Formation Probability in the XY Model

It was shown in Refs. [9,10] that spin-spin correlation functions in the XY model can be represented as determinants of some Toeplitz matrices so that one can find their asymptotic behavior using known theorems from the theory of Toeplitz matrices. This approach has been applied to EFP for the isotropic ($\gamma = 0$) XY model in magnetic field in Ref. [1]. In this section we (slightly) generalize the derivation of Ref. [1] to the case of $\gamma \neq 0$ and represent EFP as a determinant of a Toeplitz matrix.

We follow the standard prescription [9] and reformulate the Hamiltonian (2) in terms of spinless fermions ψ_i by means of a Jordan-Wigner transformation:

$$\sigma_j^+ = \psi_j^\dagger e^{i\pi \sum_{k<i} \psi_k^\dagger \psi_k} = \psi_j^\dagger \prod_{k<j} (2\psi_k^\dagger \psi_k - 1), \quad (3)$$

$$\sigma_j^z = 2\psi_j^\dagger \psi_j - 1, \quad (4)$$

where, as usual, $\sigma^\pm = (\sigma^x \pm i\sigma^y)/2$. In Fourier components $\psi_j = \sum_q \psi_q e^{iqj}$ we have:

$$H = \sum_q \left[2(\cos q - h) \psi_q^\dagger \psi_q + i\gamma \sin q \psi_q^\dagger \psi_{-q}^\dagger - i\gamma \sin q \psi_{-q} \psi_q \right]. \quad (5)$$

The Bogoliubov transformation

$$\chi_q = \cos \frac{\vartheta_q}{2} \psi_q + i \sin \frac{\vartheta_q}{2} \psi_{-q}^\dagger \quad (6)$$

with “rotation angle” ϑ_q

$$e^{i\vartheta_q} = \frac{1}{\varepsilon_q} (\cos q - h + i\gamma \sin q), \quad (7)$$

brings the Hamiltonian (5) to a diagonal form $\sum_q \varepsilon_q \chi_q^\dagger \chi_q$ with the quasiparticle spectrum

$$\varepsilon_q = \sqrt{(\cos q - h)^2 + \gamma^2 \sin^2 q}. \quad (8)$$

The fermionic correlation functions are also easy to obtain from (5). In the thermodynamic limit they read [9,10]

$$F_{jk} \equiv i\langle\psi_j\psi_k\rangle = -i\langle\psi_j^\dagger\psi_k^\dagger\rangle = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \frac{1}{2} \sin \vartheta_q e^{iq(j-k)} \quad (9)$$

$$G_{jk} \equiv \langle\psi_j\psi_k^\dagger\rangle = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \frac{1 + \cos \vartheta_q}{2} e^{iq(j-k)}. \quad (10)$$

We express EFP as the fermionic correlator [1]

$$P(n) = \langle \prod_{j=1}^n \psi_j \psi_j^\dagger \rangle. \quad (11)$$

and using Wick's theorem on the r.h.s of (11), we obtain

$$P(n) = \text{Pf}(\mathbf{M}) \equiv \sqrt{\det(\mathbf{M})}, \quad (12)$$

where $\text{Pf}(\mathbf{M})$ is the *Pfaffian* of the $2n \times 2n$ skew-symmetric matrix \mathbf{M} of correlation functions

$$\mathbf{M} = \begin{pmatrix} -i\mathbf{F} & \mathbf{G} \\ -\mathbf{G} & i\mathbf{F} \end{pmatrix}. \quad (13)$$

Here \mathbf{F} and \mathbf{G} are $n \times n$ matrices with matrix elements given by F_{jk} and G_{jk} from (9,10) respectively.

We perform a unitary transformation

$$\mathbf{M}' = \mathbf{U}\mathbf{M}\mathbf{U}^\dagger = \begin{pmatrix} 0 & \mathbf{S}_n \\ -\mathbf{S}_n^\dagger & 0 \end{pmatrix}, \quad \mathbf{U} = \frac{1}{\sqrt{2}} \begin{pmatrix} \mathbf{I} & -\mathbf{I} \\ \mathbf{I} & \mathbf{I} \end{pmatrix}, \quad (14)$$

where \mathbf{I} is the $n \times n$ unit matrix and $\mathbf{S}_n = \mathbf{G} + i\mathbf{F}$ and $\mathbf{S}_n^\dagger = \mathbf{G} - i\mathbf{F}$. This allows us to calculate the determinant of \mathbf{M} as

$$\det(\mathbf{M}) = \det(\mathbf{M}') = \det(\mathbf{S}_n) \cdot \det(\mathbf{S}_n^\dagger) = |\det(\mathbf{S}_n)|^2. \quad (15)$$

The matrix \mathbf{S}_n is a $n \times n$ Toeplitz matrix (its matrix elements depend only on the difference of row and column indices [15]). The generating function $\sigma(q)$ of a Toeplitz matrix is defined by

$$(\mathbf{S}_n)_{jk} = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \sigma(q) e^{iq(j-k)} \quad (16)$$

and can be found from (9,10) as

$$\sigma(q) = \frac{1}{2} (1 + e^{i\vartheta_q}) = \frac{1}{2} + \frac{\cos q - h + i\gamma \sin q}{2\sqrt{(\cos q - h)^2 + \gamma^2 \sin^2 q}}. \quad (17)$$

Thus, the problem of calculation of EFP

$$P(n) = |\det(\mathbf{S}_n)|, \quad (18)$$

is reduced (exactly) to the calculation of the determinant of the $n \times n$ Toeplitz matrix \mathbf{S}_n defined by the generating function (16,17). The representation (18) is exact and valid for any n . In our study we are interested in finding an asymptotic behavior at large $n \rightarrow \infty$.

In the following section we apply known theorems on the asymptotic behavior of Toeplitz determinants to extract the corresponding asymptotes of $P(n)$ at $n \rightarrow \infty$. The reader who is interested in results can skip the following section and go directly to Sec.4 where we present and discuss our results.

3 The results for the asymptotic behavior of EFP

The asymptotic behavior of EFP at $n \rightarrow \infty$ is exactly related to the asymptotic behavior of the determinant of the corresponding Toeplitz matrix (16,17,18) and can be extracted from known theorems and conjectures in the theory of Toeplitz matrices. These types of calculations have been done first in [9,10] for spin-spin correlation functions. It is well known that the asymptotic behavior of the determinant of a Toeplitz matrix as the size of the matrix goes to infinity strongly depends upon the zeros of the generating function of the matrix and upon its phase.

In the case when the generating function $\sigma(q)$ has only isolated singularities (zeros and phase discontinuities) the most general known result for the asymptotic behavior of a Toeplitz determinant is called *generalized Fisher-Hartwig conjecture* (gFH) and is due to Basor and Tracy [15]. The result can be written as

$$P(n) \sim \sum_i E_i n^{-\lambda_i} e^{-\beta_i n}, \quad (19)$$

where the decay rates β_i , the exponents λ_i , and the coefficients E_i are known functionals of $\sigma(q)$ [15]. The original *Fisher-Hartwig conjecture* (FH) [13] consists of just one term of the sum in (19). We apply gFH to the generating

function (17) and extract the asymptotic behavior of $P(n)$. All necessary theorems on the asymptotic behavior of Toeplitz determinants can be found in Ref.[12,13,14,15,17,16]. For the most recent review of gFH see also Ref.[18].

In some parts of the phase diagram we have to rely on “conjectures”, i.e., results which are not proven yet. Therefore, it is prudent to look for some independent ways of checking those results. We did some numeric calculations on these Toeplitz determinants and the results of these calculations support the analytic results presented below [19]. Also, we used some limiting cases where EFP or an asymptotic behavior of Toeplitz determinant can be obtained in an independent way to confirm our conclusions. E.g., on the line Γ_E EFP can be calculated exactly and the asymptotic behavior of the result is the same as the one obtained using FH for the whole region Σ_0 of the phase diagram.

In the rest of this section we present the results for the asymptotic behavior of $P(n)$ in different regions of the phase diagram of the XY model (Fig. 1). In this letter we concentrate on the calculation of λ and $\beta(h, \gamma)$ leaving our findings for the coefficient $E(h, \gamma)$ as well as the results of our numeric calculations for a future, more extended publication [19].

3.1 The non-critical theory (Σ_{\pm} and Σ_0)

3.1.1 The region Σ_-

In this region ($\gamma \neq 0, h < -1$) the generating function (17) is nonzero for all q and the FH conjecture is reduced to a (rigorously proven) *Strong Szegő Limit Theorem*. It gives

$$P(n) \sim E(h, \gamma)e^{-\beta(h, \gamma)n} \quad (20)$$

with

$$\begin{aligned} \beta(h, \gamma) &= - \int_{-\pi}^{\pi} \frac{dq}{2\pi} \log |\sigma(q)| \\ &= - \int_0^{\pi} \frac{dq}{2\pi} \log \left[\frac{1}{2} \left(1 + \frac{\cos q - h}{\sqrt{(\cos q - h)^2 + \gamma^2 \sin^2 q}} \right) \right]. \end{aligned} \quad (21)$$

The integral in (21) is convergent for all h and all $\gamma \neq 0$ and $\beta(h, \gamma)$ is a continuous function of its parameters. In fact, it turns out (see below) that this expression for $\beta(h, \gamma)$ describes correctly the decay rate of EFP everywhere in the phase diagram except for $\gamma = 0$. However, both pre-exponential factors




Fig. 2. Plot of the decay factor β as a function of the parameters γ and h . The function diverges for $\gamma = 0$ and is continuous for $h = \pm 1$ (although has a weak singularity at $h = \pm 1$).

and the mathematical status of the results are quite different in different parts of the phase diagram (see below).

In Figure 2, $\beta(h, \gamma)$ is plotted as a function of h at several values of γ . One can see that $\beta(h, \gamma)$ is continuous but has a weak (logarithmic) singularity at $h = \pm 1$. This is obviously one of the effects of the criticality of the model on the asymptotic behavior of EFP.

3.1.2 The region Σ_0

In Σ_0 ($\gamma \neq 0, -1 < h < 1$) the generating function $\sigma(q)$ vanishes at $q = \pm\pi$ and acquires an overall phase of π as q goes from $-\pi$ to π . The Szegő theorem is not applicable anymore. One has to rely on the FH conjecture (which, in fact is proven for this case – the case of one singularity [17]) to derive the result, which is identical to (20,21). The expression for the coefficient $E(h, \gamma)$ is different from the one in Σ_- but its form is beyond the scope of this paper.

3.1.3 The region Σ_+

In Σ_+ ($\gamma \neq 0, h > 1$) the generating function $\sigma(q)$ vanishes at the points $q = 0$ and $q = \pm\pi$, and its phase has jumps at these points. In this case an application of FH leads to some ambiguity, e.g., one obtains two values for

$\beta(h, \gamma)$: $\beta_1 = \beta$ and $\beta_2 = \beta + \pi i$ with β from (21). This ambiguity is resolved in the use of the gFH which gives EFP as a sum of two terms so that both values of β are used.

We obtain

$$P(n) \sim E(h, \gamma) [1 + (-1)^n A(h, \gamma)] e^{-n\beta(h, \gamma)}. \quad (22)$$

Thus, in this region we again have an exponential asymptotic decay of EFP but, the constant prefactor is different for even and odd n .

3.1.4 The line Γ_2 : an exact calculation

Before proceeding to the analysis of EFP on the critical lines let us check our results (20,21) on the special line¹ in the phase diagram defined by

$$h^2 + \gamma^2 = 1. \quad (23)$$

It was shown in Ref.[11] that on this line the ground state is a product of single spin states and is given by

$$|G\rangle = \prod_j |\theta, j\rangle = \prod_j \left[\cos\left(\frac{\theta}{2}\right) |\uparrow, j\rangle + (-1)^j \sin\left(\frac{\theta}{2}\right) |\downarrow, j\rangle \right], \quad (24)$$

where $|\uparrow, j\rangle$ is an up-spin state at the lattice site j etc. One can directly check that the state (24) is an eigenstate of (2) if the value of the parameter θ is

$$\cos^2 \theta = \frac{1 - \gamma}{1 + \gamma} \quad (25)$$

and (23) is satisfied. It is also easy to show [11] that this state is, in fact, the ground state of (2).

The probability of formation of a ferromagnetic string in the state (24) is obviously

$$P(n) = \sin^{2n}\left(\frac{\theta}{2}\right) = \left(\frac{1}{2} - \frac{1}{2}\sqrt{\frac{1 - \gamma}{1 + \gamma}}\right)^n \quad (26)$$

¹ We are grateful to Fabian Essler who suggested us to check our results on this special line and pointed out the reference [11] to us.

which is an exact result on the line (23). The value of $\beta(h, \gamma)$ which immediately follows from this exact result is

$$\beta(h = \sqrt{1 - \gamma^2}, \gamma) = -\log \left(\frac{1}{2} - \frac{1}{2} \sqrt{\frac{1 - \gamma}{1 + \gamma}} \right) \quad (27)$$

Indeed, one can easily check that the integral in the general formula for the decay rate (21) under the condition (23) gives precisely (27). In fact, the Toeplitz matrix generated by (17) becomes triangular on the line (23) with diagonal matrix element $(S_n)_{jj} = \sin^2(\theta/2)$ and the determinant of \mathbf{S}_n is exactly (26).

3.2 The critical line Ω_0 ($\gamma = 0$)

The case $\gamma = 0$, corresponding to the Isotropic XY Model, has been extensively studied in Ref.[1]. For $\gamma = 0$ the generating functions (17) is reduced to the one found in [1].

For $h < 1$, the generating function $\sigma(q)$ has a limited support: to find the asymptotic behavior of the determinant of the Toeplitz matrix one can apply Widom's Theorem [16] and obtain [1]:

$$P(n) \sim 2^{\frac{5}{24}} e^{3\zeta'(-1)} (1 - h)^{-\frac{1}{8}} n^{-\frac{1}{4}} \left(\frac{1 + h}{2} \right)^{\frac{n^2}{2}} \quad (28)$$

and we see that in this case, EFP decay as a Gaussian with an additional power-law pre-factor.²

For $|h| > 1$, the theory is not critical anymore and the ground state is completely polarized in z direction, giving a trivial EFP $P(n) = 0$ for $h > 1$ and $P(n) = 1$ for $h < -1$.

3.3 The critical lines $\Omega_{\pm}(h = \pm 1)$

3.3.1 The critical line $\Omega_+(h = 1)$

For $h = 1$ the generating function $\sigma(q)$ vanishes at $q = \pm\pi$ and its phase has jumps at $q = 0, \pi$. The gFH in this case produces many terms of the form (19).

² Notice, that the formula (28) was in fact obtained in [20] in the context of unitary random matrices (which are known to be equivalent to free fermions).

However, in contrast to the Σ_+ region, all terms are suppressed by power law factors of n with respect to the leading one. The leading term has β from (21) and critical exponent $\lambda = \frac{1}{16}$:

$$P(n) \sim E(\gamma) n^{-\frac{1}{16}} e^{-n\beta(1,\gamma)}. \quad (29)$$

The next subleading term (faster decay at $n \rightarrow \infty$) is given by

$$E'(\gamma)(-1)^n n^{-\frac{9}{16}} e^{-n\beta(1,\gamma)}. \quad (30)$$

Although inclusion of the latter (subleading) term is somewhat beyond even the generalized FH, we write the sum of these two terms as a conjecture for EFP at $h = 1$ and obtain

$$P(n) \sim E(\gamma) n^{-\frac{1}{16}} \left[1 + (-1)^n A(\gamma)/n^{\frac{1}{2}} \right] e^{-n\beta(1,\gamma)}. \quad (31)$$

As this result relies on this unproven conjecture, we present our numeric results for this case in Figure 3. Indeed, we see that the form (31) is in good agreement with numerics. We conclude that at $h = 1$ EFP decays exponentially at $n \rightarrow \infty$ but with an additional power law pre-factor and a damped oscillatory component.

Remark. It is curious to notice that $1/16$ and $9/16$ remind of the scaling dimensions of spins σ^x and σ^y .³ It looks as if EFP operator (1) among other things has inserted square roots of spins at the ends of the string.

3.3.2 The critical line $\Omega_-(h = -1)$

For $h = -1$ the generating function $\sigma(q)$ does not vanish but has a phase discontinuity at $q = \pm\pi$. We apply FH to obtain

$$P(n) \sim E(\gamma) n^{-\frac{1}{16}} e^{-n\beta(-1,\gamma)}. \quad (32)$$

and stretching gFH the same way as we did for $h = +1$ we include a subleading term

$$P(n) \sim E(\gamma) n^{-\frac{1}{16}} \left[1 + A(\gamma)/n^{\frac{1}{2}} \right] e^{-n\beta(-1,\gamma)}. \quad (33)$$

This form is also in good agreement with our numeric results.

³ See Ref.[10] where it was shown that power laws are $1/4$ and $9/4$ for σ^x and σ^y correlators respectively.




Fig. 3. Results of the numeric calculation of the Toeplitz determinant is shown as $P(n)e^{\beta n}n^{1/16}$ vs n at $\gamma = 1, h = 1$. The value of $\beta = \log 2 + 2G/\pi$ with Catalan's constant G is obtained from (21). The solid line is the analytic result $E(1 + (-1)^n A/n^{1/2})$ with $E = 0.8065\dots, A = 0.2399\dots$ obtained by fitting at large n . To make the plot more readable we show only every 11th point (for $n = 1, 12, 23, \dots$).

4 Discussion and Conclusions

An asymptotic behavior of Emptiness Formation Probability $P(n)$ as $n \rightarrow \infty$ for the Anisotropic XY model in a transverse magnetic field as a function of the anisotropy γ and the magnetic field h has been studied. We summarize our results in Table 1. All these results are new except for the isotropic case Ω_0 ($\gamma = 0$) which was studied in Ref.[1].

Our main motivation was to study the relation between the criticality of the theory and the asymptotics of EFP. Let us look at the results on the critical lines Ω_0 and Ω_{\pm} . The Gaussian behavior on Ω_0 is in accord with the qualitative argument of Ref.[8]. However, on Ω_{\pm} the decay of EFP is exponential instead of Gaussian, and apparently contradicts the qualitative picture of Ref.[8] (which is reproduced in the introduction to this paper for the sake of completeness). The reason for this disagreement is that at $h = \pm 1$ the model is critical for the *quasiparticles* χ defined in (6). However, we are interested in EFP for the “original” Jordan-Wigner fermions ψ , and this correlator has clearly a different expression in terms of χ than the simple one (11) which holds at $\gamma = 0$. From the technical point of view, the difference is that in the qualitative argument in favor of a Gaussian decay of EFP for critical systems there is an implicit

Table 1

Asymptotic behavior of EFP in the different regimes. The exponential decay rate β is given by Eq. (21) for all regimes. The coefficients E, A are functions of h and γ as appropriate.

EFP for an Anisotropic XY model					
Region	γ, h	criticality	$P(n)$	Eq.#	Theorem
Σ_-	$h < -1$	No	$Ee^{-n\beta}$	20	Szegö
Σ_0	$-1 < h < 1$	No	$Ee^{-n\beta}$	20	FH
Σ_+	$h > 1$	No	$E [1 + (-1)^n A] e^{-n\beta}$	22	gFH
Γ_2	$\gamma^2 + h^2 = 1$	No	$Ee^{-n\beta}$	26	Exact
Ω_+	$h = 1$	Yes	$E n^{-1/16} [1 + (-1)^n A/\sqrt{n}] e^{-n\beta}$	31	gFH
Ω_-	$h = -1$	Yes	$E n^{-1/16} [1 + A/\sqrt{n}] e^{-n\beta}$	33	gFH
Ω_0	$\gamma = 0, -1 < h < 1$	Yes	$En^{-1/4}e^{-n^2\alpha}$	28	Widom

assumption that the density of fermions (or magnetization) is related in a local way to the field responsible for critical degrees of freedom (free boson field ϕ). This assumption is not valid on the lines $h = \pm 1$. Although the theory is critical on those lines and can be described by some free field ϕ , the relation between the magnetization and this field is highly nonlocal and one can not apply the simple argument of [8] to the XY model at $h = \pm 1$.

Although EFP at the critical magnetic field does not show a Gaussian behavior, there is an important difference between the asymptotic behavior of EFP on and off the critical lines. Namely, a power-law pre-factor appears on all critical lines. For the XY model it is universal (constant on a given critical line) and takes values $\lambda = 1/4$ for $\gamma = 0$ [1] and $\lambda = 1/16$ on the lines $h = \pm 1$. It would be interesting to understand which operators determine these particular values of “scaling dimensions” of EFP (see the remark at the end of section 3.3.1).

In some parts of the phase diagram we used the so-called *generalized Fisher-Hartwig conjecture* [15] which is not proven yet. However, our numeric calculations support our analytical results, see, e.g., Figure 3. We note that to the best of our knowledge this is the first physically motivated example where the original Fisher-Hartwig conjecture fails and its extended version is necessary. We also suggested that the gFH could be used to find the subleading corrections to the asymptotic behavior, as we did for $h = \pm 1$ in (31,33). From the physical point of view, the oscillations of EFP that we obtain for $h \geq 1$ from the gFH are related to a spontaneous Z_2 symmetry breaking which occurs at $h = 1$.

In conclusion, we notice that it is straightforward to generalize our results

for nonzero temperature. The only modification is that at $T \neq 0$ the thermal correlation functions must be used instead of (9,10). Then, the generating function (17) is non-singular everywhere and we have an exponential decay of $P(n)$ in the whole phase diagram according to the standard Szegö Theorem.

5 Acknowledgments

We greatly benefited from multiple discussions with F. Essler, V.E. Korepin, and B. McCoy. AGA would like to acknowledge the financial support from Alfred P. Sloan foundation.

References

- [1] M. Shiroishi, M. Takahashi, and Y. Nishiyama, J. Phys. Soc. Jap. **70**, 3535 (2001).
Emptiness Formation Probability for the One-Dimensional Isotropic XY Model.
- [2] V.E. Korepin, N.M. Bogoliubov, and A.G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University Press, Cambridge, UK, 1993.
- [3] V.E. Korepin, A.G. Izergin, F.H.L. Essler, and D.B. Uglov, Phys. Lett. **A 190**, 182 (1994).
Correlation functions of the spin-1/2 XXX antiferromagnet.
- [4] Yu. Stroganov, J. Phys. A– Math. Gen. **34**, L179 (2001).
The Importance of being Odd.
A. V. Razumov, and Yu. G. Stroganov, J. Phys. A– Math. Gen. **34**, 3185 (2001).
Spin chains and combinatorics.
- [5] N. Kitanine, J.M. Maillet, N.A. Slavnov and V. Terras, J. Nucl. Phys. **B642**, 433-455 (2002).
Correlation functions of the XXZ spin- $\frac{1}{2}$ Heisenberg chain at free fermion point from their multiple integral representations
N. Kitanine, J.M. Maillet, N.A. Slavnov and V. Terras, J. Phys. A: Math. Gen. **35**, L385-L388 (2002).
Emptiness formation probability of the XXZ spin- $\frac{1}{2}$ Heisenberg chain at $\Delta = \frac{1}{2}$.
- [6] N. Kitanine, J.M. Maillet, N.A. Slavnov, and V. Terras, J. Phys. A: Math. Gen. **35**, L753-L758 (2002).
Large distance asymptotic behaviour of the emptiness formation probability of the XXZ spin- $\frac{1}{2}$ Heisenberg chain
- [7] V.E. Korepin, S. Lukyanov, Y. Nishiyama and M. Shiroishi, to be published in Phys. Lett. A, arXiv:cond-mat/0210140.

Asymptotic Behavior of the Emptiness Formation Probability in the Critical Phase of XXZ Spin Chain.

- [8] A.G. Abanov and V.E. Korepin, Nucl. Phys. **B 647**, 565, (2002).
On the probability of ferromagnetic strings in antiferromagnetic spin chains.
- [9] E. Lieb, T. Schultz, and D. Mattis, Ann. of Phys. **16**, 407-466 (1961)
Two Soluble Models of an Antiferromagnetic Chain
- [10] E. Barouch, and B.M. McCoy, Phys. Rev. **A 3**, 786 (1971).
Statistical Mechanics of the XY Model. II. Spin-Correlation Functions.
- [11] J. Kurmann, H. Thomas, and G. Müller, Physica A **112**, 235 (1982).
Antiferromagnetic Long-Range Order in the Anisotropic Quantum Spin Chain.
G. Müller, and R.E. Shrock, Phys. Rev. **B 32**, 5845 (1985).
Implications of direct-product ground states in the one-dimensional quantum XYZ and XY spin chains.
- [12] T.T. Wu, Phys. Rev. **149**, 380 (1966).
Theory of Toeplitz Determinants and the Spin Correlations of the Two-Dimensional Ising Model. I
- [13] M.E. Fisher and R.E. Hartwig, Adv. Chem. Phys. **15**, 333 (1968).
Toeplitz determinants, some applications, theorems and conjectures.
- [14] H. Widom, Amer. J. Math. **95**, 333 (1973).
Toeplitz Determinants with Singular Generating Functions.
- [15] E.L. Basor, and C.A. Tracy, Phys. **A 177**, 167 (1991).
The Fisher-Hartwig conjecture and generalizations.
E.L. Basor, and K.E. Morrison, Lin. Alg. App. **202**, 129 (1994).
The Fisher-Hartwig Conjecture and Toeplitz Eigenvalues.
- [16] H. Widom, Ind. Univ. Math. J. **21**, 277 (1971).
The Strong Szegö Limit Theorem for Circular Arcs.
- [17] T. Ehrhardt and B. Silbermann, J. Funct. Anal. **148**, 229-256 (1997).
Toeplitz Determinants with One Fisher-Hartwig Singularity.
- [18] T. Ehrhardt, Operator Th: Advances and App. **124**, 217-241 (2001).
A status report on the asymptotic behavior of Toeplitz determinants with Fisher-Hartwig singularities.
- [19] A.G. Abanov and F. Franchini, to be published.
- [20] J.des Cloizeaux and M.L. Mehta, J. Math. Phys. **14**, 1648 (1973).
Asymptotic Behavior of Spacing Distributions for Eigenvalues of Random Matrices.
F. Dyson, Commun. Math. Phys. **47**, 171 (1976).
Fredholm Determinants and Inverse Scattering Problems.