

Equilibrium current temperature quasi-oscillations in a ferromagnetic loop

R.G. Mints^a

^aSchool of Physics and Astronomy, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel

Equilibrium persistent current carried by a small ferromagnetic metal loop is considered. This current is shown to be quasiperiodic in temperature at low temperatures. The quasiperiod is determined mainly by the temperature dependence of the magnetization.

PACS: 73.23.Ra; 75.75.+a

Keywords: Persistent current; Ferromagnetic metal loop

1. INTRODUCTION

An equilibrium persistent current arising in a static magnetic field in a single normal metal loop results in a magnetic response, which periodically oscillates with the magnetic flux threading the loop [1,2]. These oscillations have a fundamental period given by the flux quantum $\Phi_0 = hc/2e$ and exist if the electron phase coherence is preserved. The Josephson-type magnetic response of an isolated normal metal loop subjected to a static external magnetic field was predicted theoretically [1] and studied experimentally for a variety of mesoscopic systems: an array of about 10^7 isolated mesoscopic Cooper rings [3]; a single, isolated micron-size gold loop [4]; a GaAs-AlGaAs single mesoscopic ring [5]; an array of about 10^5 GaAs-AlGaAs single mesoscopic rings [6]; and an array of 30 gold mesoscopic rings [7].

In this paper we consider an equilibrium persistent current carried by an isolated ferromagnetic metal ring in the absence of an external magnetic field. We show that at low temperatures this persistent current is quasiperiodic in temperature. The quasiperiod T is determined mainly by the temperature dependence of the equilibrium magnetization of the ferromagnet.

2. FERROMAGNET-METAL RING

Consider a ferromagnetic metal ring at a certain temperature $T < T_c$, where T_c is the Curie temperature. Suppose that the external magnetic field is equal to zero. In this case the electrons of a ferromagnetic metal are subjected to the internal magnetic field $B = 4\pi M$, where M is the magnetization. In a small ferromagnetic sample M is uniform as the formation of magnetic domains increases the free energy [8]. Therefore, a magnetic flux is threading a small ferromagnetic metal ring even in the absence of an external magnetic field.

A monotonic variation of the magnetic field $4\pi M$ results in a periodic in the flux equilibrium persistent current oscillations in a ferromagnetic metal ring. This effect is similar to the oscillations in a normal metal ring subjected to an external magnetic field. The flux induced by the field $B = 4\pi M$ can be presented as $\Phi = 4\pi M A_e$; where A_e is an effective area of the ring. The value of A_e depends on the orientation of the magnetization M and the specific geometry of the ring. In particular, if M is parallel to the axis of symmetry, then $A_e = \pi dD$, where d is the thickness and D is the diameter of the ring.

The magnetization $M(T)$ of a ferromagnet is a nonlinear function of the temperature T . A small variation of the temperature $T \rightarrow T + \delta T$ results in a flux variation $\delta\Phi = 4\pi A_e \delta M = dT \chi T$. The equilibrium persistent current is periodic in

Tel.: + 972-3-640-9165; fax: + 972-3-642-2979. E-mail address: mints@ccsg.tau.ac.il

with the period τ_0 . Therefore, the nonlinear dependence $M(T)$ results in an equilibrium current quasi-periodic in temperature. The quasi-period T follows from the relation $\tau_0 = T$, which leads to the following expression

$$T = \frac{\tau_0}{4 A_e \frac{dM}{dT}} : \quad (1)$$

It is worth noting that Eq. (1) is valid if $T \ll T_c$.

At low temperatures the dependence $M(T)$ is given by the Bloch law :

$$M = M_0 \left(1 - \frac{T}{T_c} \right)^{3/2} ; \quad (2)$$

where M_0 is the saturation magnetization and the constant $\tau_0 = 0.2 - 0.5$ depending on the ferromagnet. It follows from the Eq. (2) that

$$\frac{dM}{dT} = \frac{3 M_0}{2 T_c} \frac{T}{T_c}^{1/2} : \quad (3)$$

Combining the Eqs. (1) and (3) we find for the quasi-period T the final expression

$$T = \frac{\tau_0 T_c}{6 A_e M_0} \frac{T_c}{T}^{1/2} : \quad (4)$$

3. SUMMARY

To summarize, we demonstrate that the magnetic response of a single one-domain ring of a ferromagnetic metal is quasi-periodic in temperature even in the absence of an applied magnetic field. To estimate the quasi-period T let us consider a small dysprosium ring. Suppose the temperature $T = 4.2$ K and effective area $A_e = 3 \cdot 10^8$ cm². Using for dysprosium [9] the data $T_c = 89$ K and $M_0 = 0.29$ T and estimating $\tau_0 = 0.3$ we find $T = 0.3$ K. This value seems to be reasonable for an experimental observation.

4. ACKNOWLEDGMENTS

I would like to thank M. A. Zbel, D. K. Hmelitskii and A. Larkin for useful discussions. This research is supported by grant No. 2000-011 from the United States - Israel Binational Science Foundation (BSF), Jerusalem, Israel.

REFERENCES

1. M. Buttiker, I. Imry, and R. Landauer, Phys. Lett. A 96, 365 (1983).
2. Y. Imry, Introduction to mesoscopic physics (second edition), Oxford University Press, Oxford, 2002.
3. L. P. Levy, G. Dolan, J. Dunsmuir, and H. Bouchiat, Phys. Rev. Lett. 64, 2074 (1990).
4. V. Chandrasekhar, R. A. Webb, M. J. Brady, M. B. Ketchen, W. J. Gallagher, and A. Kleinsasser, Phys. Rev. Lett. 67, 3578 (1991).
5. D. Mailly, C. Chapelier, and A. Benoit, Phys. Rev. Lett. 70, 2020 (1993).
6. B. Reulet, M. Ramoin, H. Bouchiat, and D. Mailly, Phys. Rev. Lett. 75, 124 (1995).
7. E. M. Q. Jariwala, P. Mohanty, M. B. Ketchen, and R. A. Webb, Phys. Rev. Lett. 86, 1594 (2001).
8. L. D. Landau and E. M. Lifshitz, Electrodynamics of Continuous Media, Pergamon, Oxford, p. 157 (1984).
9. Ferromagnetic Materials, edited by E. P. Wohlfarth, North-Holland Publishing Company, Amsterdam (1980).