

Thermodynamic stability conditions for nonadditive composable entropies

Wada Tatsuaki¹

Department of Electrical and Electronic Engineering, Ibaraki University, Hitachi, Ibaraki, 316-8511, Japan

Received: date / Accepted: date

Abstract. The thermodynamic stability conditions (TSC) of nonadditive and composable entropies are discussed. Generally the concavity of a nonadditive entropy with respect to internal energy is not necessarily equivalent to the corresponding TSC. It is shown that both the TSC of Tsallis' entropy and that of the q -generalized Boltzmann entropy are equivalent to the positivity of the standard heat capacity.

Key words: Thermodynamic stability, Nonadditive entropy, Composability

PACS: 05.20.-y, 05.70.-a, 05.90.+m

1 Introduction

It is well known that the conventional thermodynamic stability condition (TSC) [1] is the concavity of the Boltzmann-Gibbs (BG) entropy S^{BG} with respect to internal energy U , i.e.,

$$\frac{\partial^2 S^{BG}}{\partial U^2} < 0; \quad (1)$$

which is also equivalent to the positivity of heat capacity $C = \partial U / \partial T$, since the temperature T is defined by

$$\frac{1}{T} = \frac{\partial S^{BG}}{\partial U}; \quad (2)$$

and

$$\frac{1}{C} = \frac{\partial T}{\partial (1/T)} = \frac{\partial (1/T)}{\partial U} = -T^2 \frac{\partial^2 S^{BG}}{\partial U^2}. \quad (3)$$

The conventional thermodynamic stability arguments are based on both the extremum principle and additivity of the BG entropy. Nowadays nonadditive entropies have been extensively considered in the literature since the generalization based on Tsallis' nonadditive entropy [2,3] has been successfully applied to a variety of complex systems. The Tsallis entropy is a generalization of the BG entropy by one real parameter q ,

$$S_q = -\frac{1}{1-q} \sum_i^P p_i^q, \quad (4)$$

where p_i stands for a probability of i -th state. For the sake of simplicity, we set the Boltzmann constant to unity throughout this paper. S_q reduces to the BG entropy in the limit of $q \rightarrow 1$.

During the early stages of the development of Tsallis' thermodynamics, Ramshaw [4] pointed out that the concavity of Tsallis' entropy with respect to the standard expectation energy $U_1 = \sum_i p_i E_i$

is not sufficient to guarantee the thermodynamic stability for all values of q because S_q is nonadditive. He derived the TSC of the nonadditive S_q from the TSC of the additive Renyi entropy [5]

$$S_q^R = \frac{\ln \sum_i p_i^q}{1-q}; \quad (5)$$

by utilizing the nonlinear functional relation between both entropies

$$S_q^R = \frac{\ln [1 + (1-q)S_q]}{1-q}; \quad (6)$$

The present author [6] further studied the TSC of S_q by directly utilizing the nonadditivity Eq. (9) of S_q , and shown that the resultant TSC is equivalent to the positivity of the standard heat capacity C .

In this paper we discuss the TSCs of nonadditive composable entropies, and show that the TSC of such a nonadditive entropy is not equivalent to the concavity of the entropy with respect to internal energy. In the next section, after the explanation of the concept of composability, we briefly review the standard TSC of an additive entropy. We then discuss the TSCs of two different kinds of nonadditive composable entropies. The subsection 2.1 deals with the TSC of the Tsallis q -entropy, and in the subsection 2.2 we consider the TSC of the q -generalized Boltzmann entropy. It is shown that the TSCs of the both nonadditive entropies are equivalent to the positivity of the standard heat capacity. The final section is devoted to the conclusions.

2 Thermodynamic stability conditions of composable entropies

The concept of composability [7] is important and quite useful when we consider a thermodynamic system which consists of independent subsystems. Suppose a total system consists of two independent subsystems A and B . If the total entropy of any sort $S(A;B)$ is a bivariate and symmetric function f of the subsystem entropies $S(A)$ and $S(B)$, i.e.,

$$S(A;B) = f(S(A);S(B)) = f(S(B);S(A)); \quad (7)$$

we say S is composable. This means that the total entropy $S(A;B)$ can be built of just the two macroscopic quantities, $S(A)$ and $S(B)$. Hence we can further discuss the thermodynamic properties of any composable system without knowing its underlying microscopic dynamics. For example, the zeroth law of thermodynamics [8,9,10,11] for Tsallis' entropy has been discussed by assuming the composability. It is well known that the BG entropy is composable and additive,

$$S^{BG}(A;B) = S^{BG}(A) + S^{BG}(B); \quad (8)$$

Tsallis' entropy is also composable but it's nonadditive, i.e., so-called pseudo-additive [2,3]

$$S_q(A;B) = S_q(A) + S_q(B) + (1-q)S_q(A)S_q(B); \quad (9)$$

Let us now review the TSC of any additive entropy $S(U)$. We denote the maximum entropy of a thermodynamic system with an internal energy U by $S(U)$. For the sake of simplicity, the internal energy is assumed to be additive, i.e., $U(A;B) = U(A) + U(B)$, in this paper. It is however worth noting that Wang [11] has discussed the zeroth and first laws of thermodynamics by utilizing nonadditive energy expectation value within the framework of nonextensive statistical mechanics.

The essence of conventional thermodynamic stability [1] lies in the entropy maximum principle and additivity of $S(U)$. It is known that the relation between the concavity of $S(U)$ and the standard TSC is straightforward as follows. If one transfer an amount of energy U from one of two identical subsystems to the other subsystem, the total entropy changes from its initial value of $S(U;U) = 2S(U)$ to $S(U+U;U-U) = S(U+U) + S(U-U)$. The entropy maximum principle demands that the resultant value of the entropy is not larger than the initial one, i.e.,

$$2S(U) \geq S(U+U) + S(U-U); \quad (10)$$

which is the TSC of the additive entropy $S(U)$. In the limit of $U \rightarrow 0$, Eq. (10) reduces to the concavity of $S(U)$ with respect to U ,

$$\frac{\partial^2 S(U)}{\partial U^2} \leq 0; \quad (11)$$

Thus the TSC and the concavity of $S(U)$ with respect to U are equivalent each other when $S(U)$ is additive. However this is not the case for a nonadditive entropy as we will see in the subsequent subsections.

2.1 Tsallis' entropy

Let us now turn focus on the TSC of Tsallis' entropy [6]. Taking account of the pseudo-additivity Eq. (9), the TSC of S_q can be written by

$$2S_q(U) + (1-q)[S_q(U)]^2 \geq S_q(U+U) + S_q(U-U) + (1-q)S_q(U+U)S_q(U-U): \quad (12)$$

The physical meaning of this inequality is same as that of the conventional TSC Eq. (10). If we transfer a small amount of energy U from one of two identical subsystems to the other subsystem, then the total entropy changes from its initial value $S_q(U;U)$ of the left-hand-side to the resultant value $S_q(U+U;U-U)$ of the right-hand-side in Eq. (12). The principle of the maximum Tsallis entropy demands that the resultant entropy should not be larger than the initial one. The remarkable difference between Eq. (12) and Eq. (10) is the presence of the nonlinear term proportional to $1-q$, which originally arises from the pseudo-additivity of Eq. (9).

In the limit of $U \rightarrow 0$, Eq. (12) reduces to the differential form

$$\frac{\partial^2 S_q(U)}{\partial U^2} + (1-q)S_q(U)\frac{\partial^2 S_q(U)}{\partial U^2} - \left(\frac{\partial S_q(U)}{\partial U}\right)^2 \geq 0: \quad (13)$$

Now we introduce the q -generalized temperature T_q defined by

$$\frac{1}{T_q} = \frac{\partial S_q}{\partial U}; \quad (14)$$

and the q -generalized heat capacity C_q defined by

$$\frac{1}{C_q} = \frac{\partial T_q}{\partial U} = -T_q^2 \frac{\partial^2 S_q}{\partial U^2}: \quad (15)$$

Then Eq. (13) becomes

$$\frac{1 + (1-q)S_q}{C_q} + (1-q) \geq 0: \quad (16)$$

We thus see that only the positivity of C_q , or equivalently the concavity of $S_q(U)$, is not sufficient to satisfy the TSC of the nonadditive Tsallis entropy. When $q \rightarrow 1$, however, the positivity of C_q guarantees to satisfy Eq. (16) since $1 + (1-q)S_q = \sum_i p_i^q$ is always positive.

Next we shall consider the relation between the TSC and the positivity of the standard heat capacity. By maximizing the total Tsallis entropy $S_q(A;B) = 0$ under the constraint of the total energy conservation $U(A;B) = 0$, we obtain the equilibrium condition for a composite system described by Tsallis' entropy,

$$T_q(A)f_1 + (1-q)S_q(A)g = T_q(B)f_1 + (1-q)S_q(B)g; \quad (17)$$

which should be equal to the intensive temperature T of the total system [8], i.e.,

$$T = f_1 + (1-q)S_q g = T_q: \quad (18)$$

By differentiating the both sides of Eq. (18) with respect to U , we obtain the relation between the standard heat capacity $C = \partial U / \partial T$ and the q -generalized heat capacity C_q as

$$\frac{1}{C} = \frac{1 + (1-q)S_q}{C_q} + 1 - q: \quad (19)$$

The TSC condition of Eq. (16) is thus equivalent to the positivity of the conventional heat capacity,

$$C \geq 0: \quad (20)$$

2.2 The q -generalized Boltzmann entropy

Next we turn focus on another nonadditive composable entropy. Kaniadakis [12,13] has introduced another type of one parameter deformations of the exponential and logarithmic functions

$$\exp_{f_g}(x) = \frac{1}{1 - \frac{x}{i_1}}; \quad (21)$$

$$\ln_{f_g}(x) = \frac{x}{2}; \quad (22)$$

where i_1 is a real parameter with $1 < i_1 < \infty$. In the limit of $i_1 \rightarrow \infty$, the $\exp_{f_g}(x)$ and $\ln_{f_g}(x)$ reduce to the standard exponential and logarithmic functions, respectively. The q -deformed exponential and logarithmic functions have the following properties

$$\exp_{f_g}(x) \exp_{f_g}(y) = \exp_{f_g}(x + y); \quad (23)$$

$$\ln_{f_g}(xy) = \ln_{f_g}(x) + \ln_{f_g}(y); \quad (24)$$

where $+$ -sum is defined by

$$x + y = x \frac{1}{1 - \frac{y}{i_1}} + y \frac{1}{1 - \frac{x}{i_1}}; \quad (25)$$

Kaniadakis has defined the q -deformed entropy as

$$S^K = - \sum_i p_i \ln_{f_g} p_i; \quad (26)$$

to construct a generalized statistical mechanics in the context of special relativity [14]. S^K reduces to the Shannon-BG entropy in the limit of $i_1 \rightarrow \infty$. The q -additivity defined by Eq. (25) is the additivity of relativistic momenta as shown in Ref. [14].

We shall here consider the TSC of the q -generalized Boltzmann entropy defined by

$$S^B(U) = \ln_{f_g} W(U); \quad (27)$$

where W is the number of microcanonical configurations of a thermodynamic system. $S_0^B(U) = \ln W(U)$ is the standard Boltzmann entropy. Note that this S^B is different from the Kaniadakis q -entropy S^K , which is an entropy à la Gibbs, whereas the S^B is one à la Boltzmann. The reason of using S^B , instead of S^K , is that it has the composability (the q -additivity),

$$S^B(A;B) = S^B(A) + S^B(B); \quad (28)$$

By utilizing this composition rule we readily write down the expression of the TSC of S^B ,

$$\frac{\partial^2 S^B(U)}{\partial U^2} = \frac{\partial^2 S^B(U;U)}{\partial U^2} = \frac{\partial^2 S^B(U+U;U)}{\partial U^2} = \frac{\partial^2 S^B(U)}{\partial U^2} + \frac{\partial^2 S^B(U)}{\partial U^2} = 2 \frac{\partial^2 S^B(U)}{\partial U^2}; \quad (29)$$

After some algebra, in the limit of $U \rightarrow \infty$, Eq. (29) becomes

$$\frac{1 + 2 \frac{\partial^2 S^B(U)}{\partial U^2}}{1 + \frac{\partial^2 S^B(U)}{\partial U^2}} = \frac{\partial^2 S^B(U)}{\partial U^2} = \frac{\partial S^B(U)}{\partial U} = \frac{1}{1 + \frac{\partial^2 S^B(U)}{\partial U^2}} = 0; \quad (30)$$

Since the terms in front of the brace is always positive, the differential form of the TSC of S^B becomes

$$\frac{\partial^2 S^B}{\partial U^2} = \frac{\partial S^B}{\partial U} = \frac{1}{1 + \frac{\partial^2 S^B}{\partial U^2}} = 0; \quad (31)$$

Furthermore since $W(U)$ is much larger than unity for almost every energy U in conventional thermodynamics situations, $S^B(U) = \ln_{f_g} W(U)$ is positive for such an energy. Thus the concavity of $S^B(U)$ guarantees to satisfy the TSC of Eq. (31) unlike the case of the TSC of Tsallis' entropy. Interestingly the concavity $\partial^2 S^B(U) = \partial^2 U < 0$ is the sufficient, but not necessary, condition in order to satisfy the TSC of S^B . For example, it is possible to satisfy Eq. (31) even if S^B is convex (or C_{fg} is negative).

Next we turn to the relation between the TSC and the standard heat capacity. We introduce the α -generalized temperature T_{f-g} defined by

$$\frac{1}{T_{f-g}} = \frac{\partial S^B(U)}{\partial U} : \quad (32)$$

Following the similar arguments of the equilibrium condition (or thermodynamic zeroth law) for Tsallis' entropy in the previous subsection, we obtain the relation between T_{f-g} and the corresponding intensive temperature $T = T_{f0g}$ as

$$T = T_{f-g} \frac{1}{1 + \alpha^2 [S^B(U)]^2} \quad (33)$$

Alternatively we can obtain this relation from the definition Eq. (32) of T_{f-g} ,

$$\frac{1}{T_{f-g}} = \frac{\partial}{\partial U} \left(\frac{W}{2} \right) = \frac{\partial \ln W}{\partial U} \frac{W}{2} + \frac{W}{2} = \frac{1}{T_{f0g}} \frac{1}{1 + \alpha^2 [S^B]^2} : \quad (34)$$

By differentiating the both sides of Eq.(33) with respect to U , we obtain the relation between the standard heat capacity C and the α -generalized heat capacity $C_{f-g} = \partial U / \partial T_{f-g}$ as follows.

$$\begin{aligned} \frac{1}{C} &= \frac{1}{T_{f-g}^2} \frac{\partial}{\partial U} \left(\frac{1}{1 + \alpha^2 [S^B]^2} \right) = \frac{1}{T_{f-g}^2} \left(-\frac{2\alpha^2 S^B}{1 + \alpha^2 [S^B]^2} \right) \\ &= -\frac{2\alpha^2 S^B}{T_{f-g}^2} \frac{1}{1 + \alpha^2 [S^B]^2} = -\frac{2\alpha^2 S^B}{T_{f-g}^2} \frac{\partial S^B / \partial U}{1 + \alpha^2 [S^B]^2} ; \end{aligned} \quad (35)$$

where we used Eq. (32) and the relation $1/C_{f-g} = \partial T_{f-g} / \partial U = -\partial S^B / \partial U^2$. By comparing Eq (35) with Eq. (31), we finally find that the TSC of S^B is also equivalent to the positivity of standard heat capacity $C > 0$.

3 Conclusions

We have discussed the thermodynamic stability conditions (TSC) of the two nonadditive composable entropies, which are the Tsallis and α -generalized Boltzmann entropies. Unlike the TSC of an additive entropy, the concavity of a nonadditive composable entropy is not necessarily equivalent to the corresponding TSC in general. It is shown that in addition to the TSC of Tsallis' entropy, the TSC of the α -generalized Boltzmann entropy is also equivalent to the positivity of the standard heat capacity. It is interesting to further study whether it is common feature to composable entropy.

Nonadditive entropy can provide a basic ingredient in order to describe non-separable or correlated systems whose thermodynamic behaviors are complex and anomalous. It is worth noting that the we have derived the TSCs with resort to the composability of the nonadditive entropies. However composability of systems restricts our consideration to separable systems, which are divisible into independent subsystems. For non-composable entropies we have not yet known the expression of the corresponding TSCs.

Acknowledgments

The author greatly thanks Prof. M. Sugiyama for giving the opportunity to reconsider the thermodynamic stability of the nonextensive entropies. He also thanks G. Kaniadakis and S. Abe for valuable discussions.

References

1. Callen H.B Thermodynamics and an Introduction to Thermostatistics 2nd ed. (Wiley New York 1985) Chap. 8
2. Tsallis C (1988) Possible generalization of Boltzmann-Gibbs statistics. J. Stat. Phys. 52 479

3. Kaniadakis G, Lissia M, Rapisarda A (editors) (2001) Proc. of the international school and workshop on non extensive thermodynamics and physical applications (NEXT 2001) Physica A 305 Nos. 1-2
4. Ramshaw JD (1995) Thermodynamic stability conditions for the Tsallis and Renyi entropies. Phys. Lett. 198:119-121
5. Renyi A Probability Theory (North-Holland, Amsterdam 1970)
6. Wada T (2002) On the thermodynamic stability conditions of Tsallis' entropy. Phys. Lett. 297:334-337
7. Hotta M and Joichi I (1999) Compossibility and generalized entropy Phys. Lett. A 262 302
8. Abe S (2001) General pseudoadditivity of composable entropy prescribed by the existence of equilibrium. Phys. Rev. E 63:061105
9. Martinez S, Pennini F, Plastino A (2001) Thermodynamics' zeroth law in a nonextensive scenario Physica A 295:416-424
10. Abe S (2001) Heat and entropy in nonextensive thermodynamics: transmutation from Tsallis theory to Renyi-entropy-based theory Physica A 300:417-423
11. Wang Q A, Le梅haute A (2003) Unnormalized nonextensive expectation value and zeroth law of thermodynamics Chaos, Solitons & Fractals 15:537-541
12. Kaniadakis G (2001) Non-linear kinetics underlying generalized statistics. Physica A 296:405-425
13. Kaniadakis G, Scarfone AM (2002) A new one-parameter deformation of the exponential function Physica A 305:69-75
14. Kaniadakis G (2002) Statistical mechanics in the context of special relativity. Phys. Rev. E 66:056125