# Space-time Engineering with *Lasetron* Pulses

M. Kozlowski<sup>1)</sup>, J. Marciak-Kozlowska<sup>2)</sup>

<sup>1)</sup>Institute of Experimental Physics and Science Teachers College of Warsaw University, Hoża 69, 00-681 Warsaw, Poland, <sup>2)</sup>Institute of Electron Technology, Al Lotników 32/46, 02-668 Warsaw, Poland

#### Abstract

The LASETRON project (*Phys. Rev. Lett.*, **88**, (2002), p. 074801-1) offers the possibility of the investigation of electron-positron structure of the space-time. Following our results (*From Quarks to Bulk Matter*, Hadronic Press, 2001) we analyze theoretically possibility of the penetration of zeptosecond laser pulses  $(10^{-21} \text{ s})$  through space-time.

**Key words:** Zeptosecond laser pulses; *LASETRON*; Electron-positron pairs; Space-time.

### 1. Introduction

In the paper [1] the theoretical project of the *LASETRON* was described. As was shown in paper [1] the  $10^{-21}$  s (zeptosecond) laser pulses can be generated using petawatt lasers, while already available terawatt lasers may generate subattosecond laser pulses of  $10^{-19}$  s. The pulses will be radiated by ultrarelativistic electrons driven by circularly polarized high-intensity laser fields. *LASETRON* pulses can be achieved by placing a solid particle or a piece of wire of subwavelength cross section in the focal plane of a superpowerful laser.

In the book [2] it was shown that the lifetime of the electron-positron pair is of the order of  $10^{-17}$ s. Strictly speaking the lifetime of the order of the relaxation time  $\tau$ 

$$\tau = \frac{\hbar}{m_e \alpha^2 c^2} \approx 10^{-17} \,\mathrm{s} \,. \tag{1}$$

In formula (1)  $m_e = m_p$  is electron, positron mass,  $\alpha$  is the fine structure constant and *c* is the vacuum light velocity. As can be concluded from formula (1) the *zitterbegung* or *tremor* of space-time can be investigated with the *LASETRON* pulses, for the latter are shorter than the characteristic relaxation time.

For the time period  $\Delta t < \tau$ , i.e. for *LASETRON* pulse the vacuum of space-time is filled with the gas of electron-positron pairs with life-time  $\approx 10^{-17}$  s. In that case the propagation of *LASETRON* pulse will be described by the Heaviside equation;

$$\frac{\partial^2 E}{\partial t^2} + \frac{\sigma}{\varepsilon_0} \frac{\partial E}{\partial t} = c_\gamma^2 \frac{\partial^2 E}{\partial x^2}.$$
 (2)

In equation (1)  $\varepsilon_0$  – permittivity of free space-time, and  $\sigma$  is the conductivity of free space-time,  $c_{\gamma}$  is the photon velocity<sup>\*</sup> in space-time and *E* is electric field (in one dimension).

In the seminal paper [4] F. Calogero described the cosmic origin of quantization. In paper [4] the *tremor* of the cosmic particles is the origin of the quantization and the characteristic acceleration of these particles  $a \approx 10^{-10}$  m/s<sup>2</sup> was calculated. In our earlier paper [5] the same value of the acceleration was obtained and compared to the experimental value of the measured space-time acceleration [6]. In this

 $<sup>{}^{*}</sup>C_{\gamma}$  is the photon speed in space-time filled with electron-positron pairs. The

electron-positron fluctuations can change the c to  $c_{\gamma}$ , i.e  $c_{\gamma} \neq c$  [3].

paper we define the cosmic force — *Planck* force,  $F_{\text{Planck}} = M_P a_{\text{Planck}} (a_{\text{Planck}} \approx a)$  and study the history of Planck force as the function of the age of the Universe.

Masses introduce a curvature in space-time, light and matter are forced to move according to space-time metric. Since all the matter is in motion, the geometry of space is constantly changing. A Einstein relates the curvature of space to the mass/energy density:

$$\mathbf{G} = k\mathbf{T} \,, \tag{3}$$

)

**G** is the Einstein curvature tensor and **T** the stress-energy tensor. The proportionality factor *k* follows by comparison with Newton's theory of gravity:  $k = G/c^4$  where *G* is the Newton's gravity constant and *c* is the vacuum velocity of light; it amounts to about  $2.10^{-43}$  N<sup>-1</sup>, expressing the *rigidity* of space-time.

In paper [5] the model for the acceleration of space-time was developed. Prescribing the -G for space-time and +G for matter the acceleration of space-time was obtained:

$$a_{\text{Planck}} = -\frac{1}{2} \left(\frac{\pi}{4}\right)^{\frac{1}{2}} \frac{\left(N + \frac{3}{4}\right)^{\frac{1}{2}}}{M^{\frac{3}{2}}} A_P$$
(4)

where  $A_p$ , *Planck* acceleration equal, viz.:

$$A_{P} = \left(\frac{c^{7}}{\hbar G}\right)^{\frac{5}{2}} = \frac{c}{\tau_{P}} \cong 10^{51} \text{ms}^{-2}.$$
 (5)

As was shown in paper [5] the  $a_{\text{Planck}}$  for  $N = M = 10^{60}$  is of the order of the acceleration detected by Pioneer spacecrafts [6].

Considering  $A_p$  it is quite natural to define the *Planck* force  $F_{Planck}$ ,

$$F_{\text{Planck}} = M_P A_P = \frac{c^4}{G} = k^{-1}, \qquad (6)$$

where

$$M_P = \left(\frac{\hbar c}{G}\right)^{\frac{1}{2}}.$$

From formula (6) we conclude that  $F_{\text{Planck}}^{-1}$  = rigidity of the space-time. The *Planck* force,  $F_{\text{Planck}} = c^4 / G = 1.2 \cdot 10^{44} N$  can be written in units which characterize the microspace-time, i.e. GeV and fm. In that units

$$k^{-1} = F_{\text{Planck}} = 7.6 \cdot 10^{38} \text{GeV} / \text{fm}$$

## 2. The Planck, Yukawa and Bohr forces

As was shown in paper [5] the present value of *Planck* force equal

$$F_{\text{Planck}}^{\text{Now}}(N=M=10^{60}) \cong -\frac{1}{2} \left(\frac{\pi}{4}\right)^{\frac{1}{2}} 10^{-60} \frac{c^4}{G} = -10^{-22} \frac{\text{GeV}}{\text{fm}}.$$
 (7)

In papers [7, 8] the *Planck* time  $\tau_P$  was defined as the relaxation time for space-time

$$\tau_P = \frac{\hbar}{M_P c^2}.$$
(8)

Considering formulae (6) and (8)  $F_{\text{Planck}}$  can be written as

$$F_{\text{Planck}} = \frac{M_P c}{\tau_P},\tag{9}$$

where c is the velocity for gravitation propagation. In papers [7, 8] the velocities and relaxation times for thermal energy propagation in atomic and nuclear matter were calculated:

$$v_{\text{atomic}} = \alpha_{em} c, \tag{10}$$
$$v_{\text{nuclear}} = \alpha_{s} c,$$

where  $\alpha_{em} = e^2 / (\hbar c) = 1/137$ ,  $\alpha_s = 0.15$ . In the subsequent we define atomic and nuclear accelerations:

$$a_{\text{atomic}} = \frac{\alpha_{em}c}{\tau_{\text{atomic}}},$$

$$a_{\text{nuclear}} = \frac{\alpha_{s}c}{\tau}.$$
(11)

 $\tau_{\text{nuclear}}$ Considering that  $\tau_{\text{atomic}} = \hbar/(m_e \alpha_{em}^2 c^2)$ ,  $\tau_{\text{nuclear}} = \hbar/(m_N \alpha_s^2 c^2)$  one obtains from formula (11)

$$a_{\text{atomic}} = \frac{m_e c^3 \alpha_{em}^3}{\hbar},$$

$$a_{\text{nuclear}} = \frac{m_N c^3 \alpha_s^3}{\hbar}.$$
(12)

We define, analogously to *Planck* force the new forces:  $F_{\rm Bohr}$ ,  $F_{\rm Yukawa}$ 

$$F_{\text{Bohr}} = m_e a_{\text{atomic}} = \frac{\left(m_e c^2\right)^2}{\hbar c} \alpha_{em}^3 = 5 \cdot 10^{-13} \frac{\text{GeV}}{\text{fm}},$$

$$F_{\text{Yukawa}} = m_N a_{\text{nuclear}} = \frac{\left(m_N c^2\right)^2}{\hbar c} \alpha_s^3 = 1.6 \cdot 10^{-2} \frac{\text{GeV}}{\text{fm}}.$$
(13)

Comparing formulae (7) and (13) we conclude that gradients of *Bohr* and *Yukawa* forces are much large than  $F_{\text{Planck}}^{\text{Now}}$ , i.e.:

$$\frac{F_{\text{Bohr}}}{F_{\text{Planck}}^{\text{Now}}} = \frac{5 \cdot 10^{-13}}{10^{-22}} \cong 10^9,$$

$$\frac{F_{\text{Yukawa}}}{F_{\text{Planck}}^{\text{Now}}} = \frac{10^{-2}}{10^{-22}} \cong 10^{20}.$$
(14)

The formulae (14) guarantee present day stability of matter on the nuclear and atomic levels.

As the time dependence of  $F_{Bohr}$  and  $F_{Yukawa}$  are not well established, in the subsequent we will assumed that  $\alpha_s$  and  $\alpha_{em}$  [9] do not dependent on time. Considering formulae (9) and (12) we obtain

$$\frac{F_{\text{Yukawa}}}{F_{\text{Planck}}} = \frac{1}{\left(\frac{\pi}{A}\right)^{\frac{1}{2}}} \frac{\left(m_N c^2\right)^2}{M_P c^2} \frac{\alpha_s^3}{\hbar} T,$$
(15)

$$\frac{F_{\rm Bohr}}{F_{\rm Planck}} = \frac{1}{\left(\frac{\pi}{4}\right)^{\frac{1}{2}}} \frac{\left(m_e c^2\right)^2}{M_P c^2} \frac{\alpha_{em}^3}{\hbar} T.$$
 (16)

As can be realized from formulae (15), (16) in the past  $F_{\text{Planck}} \approx F_{\text{Yukawa}}$  (for T = 0.002 s) and  $F_{\text{Planck}} \approx F_{\text{Bohr}}$  (for  $T \approx 10^8$  s), T = age of universe. The calculated ages define the limits for instability of the nuclei and atoms.

### 3. The Planck, Yukawa and Bohr particles

In 1900 M. Planck [10] introduced the notion of the universal mass, later on called the *Planck* mass

$$M_P = \left(\frac{\hbar c}{G}\right)^{\frac{1}{2}}.$$
 (17)

Considering the definition of the Yukawa force (13)

$$F_{\rm Yukawa} = \frac{m_N v_N}{\tau_N} = \frac{m_N \alpha_{\rm strong} c}{\tau_N}, \qquad (18)$$

the formula (18) can be written as:

$$F_{\rm Yukawa} = \frac{m_{\rm Yukawa}c}{\tau_N},\tag{19}$$

where

$$m_{\rm Yukawa} = m_N \alpha_{\rm strong} \cong 147 \frac{MeV}{c^2} \sim m_{\pi} \,.$$
 (20)

From the definition of the *Yukawa* force we deduced the mass of the particle which mediates the strong interaction – pion mass postulated by Yukawa in [11].

Accordingly for *Bohr* force:

$$F_{\rm Bohr} = \frac{m_e v}{\tau_{\rm Bohr}} = \frac{m_e \alpha_{em} c}{\tau_{\rm Bohr}} = \frac{m_{\rm Bohr} c}{\tau_{\rm Bohr}}, \qquad (21)$$

$$m_{\rm Bohr} = m_e \alpha_{em} = 3.7 \frac{\rm keV}{c^2}.$$
 (22)

For the Bohr particle the range of interaction is

$$\gamma_{\rm Bohr} = \frac{\hbar}{m_{\rm Bohr}c} \approx 0.1\,\rm nm\,, \qquad (23)$$

which is of the order of atomic radius.

Considering the electromagnetic origin of the mass of the *Bohr* particle, the planned sources of hard electromagnetic field *LASETRON* [1] are best suited to the investigation of the properties of the *Bohr* particles.

# 4. Possible interpretation of $F_{\text{Planck}}$ , $F_{\text{Yukawa}}$ and $F_{\text{Bohr}}$

In an important work, published already in 1951 J. Schwinger [12] demonstrated that in the background of a static uniform electric field, the QED space-time is unstable and decayed with spontaneous emission of  $e^+e^-$  pairs. In the paper [12] Schwinger calculated the critical field strengths  $E_s$ :

$$E_s = \frac{m_e^2 c^3}{e\hbar}.$$
 (24)

Considering formula (23) we define the Schwinger force:

$$F_{\rm Schwinger}^{e} = eE_{S} = \frac{m_{e}^{2}c^{3}}{\hbar}.$$
 (25)

Formula (24) can be written as:

$$F_{\rm Schwinger}^{e} = \frac{m_{e}c}{\tau_{Sch}},$$
(26)

where

$$\tau_{sch} = \frac{\hbar}{m_e c^2} \tag{27}$$

is *Schwinger* relaxation time for the creation of  $e^+e^-$  pair. Considering

formulae (13) the relation of  $F_{Yukawa}$  and  $F_{Bohr}$  to the *Schwinger* force can be established

$$F_{\text{Yukawa}} = \alpha_s^3 \left(\frac{m_N}{m_e}\right)^2 F_{\text{Schwinger}}^e, \quad \alpha_s = 0.15,$$

$$F_{\text{Bohr}} = \alpha_{em}^3 F_{\text{Schwinger}}^e, \quad \alpha_{em} = \frac{1}{137},$$
(28)

and for *Planck* force

$$F_{Planck} = \left(\frac{M_P}{m_e}\right)^2 F_{\text{Schwinger}}^e .$$
<sup>(29)</sup>

In Table 1 the values of the  $F_{\text{Schwinger}}^{e}$ ,  $F_{\text{Planck}}$ ,  $F_{\text{Yukawa}}$  and  $F_{\text{Bohr}}$  are presented, all in the same units GeV/fm. As in those units the forces span the range  $10^{-13}$  to  $10^{38}$  it is valuable to recalculate the *Yukawa* and *Bohr* forces in the units natural to nuclear and atomic level. In that case one obtains:

$$F_{Y_{ukawa}} \approx 16 \frac{\text{MeV}}{\text{fm}}$$
 (30)

It is quite interesting that  $a_v \approx 16 \text{ MeV}$  is the volume part of the binding energy of the nuclei (droplet model).

Table 1: Schwinger, Planck, Yukawa and Bohr forces

$F^{e}_{ m Schwinger}$	$F_{\rm Planck}$	$F_{ m Yukawa}$	$F_{ m Bohr}$
[GeV/fm]	[GeV/fm]	[GeV/fm]	[GeV/fm]
≈10 <sup>-6</sup>	$\approx 10^{38}$	$\approx 10^{-2}$	$\approx 10^{-13}$

For the *Bohr* force considering formula (13) one obtains:

$$F_{\rm Bohr} \approx \frac{50 \,\mathrm{eV}}{0.1 \,\mathrm{nm}} \,. \tag{31}$$

Considering that the *Rydberg* energy  $\approx 27$  eV and *Bohr* radius  $\approx 0.1$  nm formula (31) can be written as

$$F_{\rm Bohr} \approx \frac{\rm Rydberg\ energy}{\rm Bohr\ radius}.$$
 (32)

### 5. Concluding remarks

In this paper the forces: *Planck*, *Yukawa* and *Bohr* were defined. It was shown that the present value of the *Planck* force (which is the source of the universe acceleration)  $\approx 10^{-22}$  GeV/fm is much smaller than the *Yukawa* ( $\approx 10^{-2}$  GeV/fm) and *Bohr* ( $10^{-13}$  GeV/fm) forces respectively. This fact guarantees the stability of the matter in the present. However in the past for *T* (age of the universe), *T* < 0.002 s,  $F_{\text{Yukawa}} < F_{\text{Planck}}$  (0.002 s) and  $F_{\text{Bohr}} < F_{\text{Planck}}$  ( $10^{8}$  s). In this paper the relation of the *Schwinger* force (for the vacuum creation of the *e^+e^-* pairs to the *Planck*, *Yukawa* and *Bohr* force was obtained. With the *LASETRON* pulses the electron-positron structure of space-time can be investigated.

#### References

- Kaplan A. F., Shkolnikov P., *Phys. Rev. Lett.*, 88, (2002), p. 074801-1.
- [2] Kozlowski M., Marciak-Kozlowska J., *From Quarks to Bulk Matter*, Hadronic Press, 2001.
- [3] Gharibyam V., arXiv/hep-ex/03030/0.
- [4] Calogero. F, Phys. Letters, A228, (1997) p. 335.
- [5] Kozlowski M., Marciak-Kozlowska J., *Nuovo Cimento*, vol, 116B, (2001) p. 821.
- [6] Anderson J. D. et al., *Phys. Rev. Lett.*, **81** (1998) p. 2858.
- [7] Marciak-Kozlowska J., Kozlowski M., *Foundations of Physics Letters*, vol. 9, (1996) p. 235.
- [8] Kozlowski M., Marciak-Kozlowska J., *Foundations of Physics Letters*, vol. 10, (1997) p. 295.
- [9] Kozlowski M., Marciak-Kozlowska J., arXiv/astro-ph/0307168.
- [10] Planck M. *The theory of heat radiation*, Dover Publications 1959, p. 173.
- [11] Yukawa H, Proc. Phys.-Math. Soc. Japan, 17, (1935) p. 48.
- [12] Schwinger J., Phys. Rev., 82 (1951) p. 664.